

SCIENCE'S COMPASS

I find that graduate students in the sciences, at one time myself included, tend not to realize just how well they are being prepared for any conceivable career choice. It seems that the trauma of the graduate school-postdoctoral daze must first be given time to subside before most can appreciate the universality of their training. Some self-inspection is necessary, as Gale implies, to realize that academic appointments are just one among the many rewarding ways to translate these increasingly valuable skills into a livelihood.

T. J. Murphy

Department of Pharmacology, Emory University School of Medicine, Atlanta, GA 30322, USA. E-mail: tmurphy@pharm.emory.edu

Environmental Health: Nickel-and-Diming It

Researchers in environmental toxicology should endorse the argument advanced by B. M. Lester, L. L. LaGasse, and R. Seifer (Policy Forum, *Science's Compass*, 23 Oct., p. 633) on how to interpret the outcome of cocaine abuse during pregnancy. A predicted epidemic of "crack babies" never materialized. Instead, they note, the offspring exhibit subtle deficiencies such as IQ reductions of about 3%. Although

small magnitude, it is a gap with vital public health and policy implications that even many scientists fail to appreciate.

I urged such a perspective for neurotoxicity risk assessment some time ago (1) and for grasping the consequences of maternal drug abuse (2). The definition of excessive lead exposure in children is now based largely on shifts in the population distribution of IQ scores (3), as are many of the economic benefits flowing from the removal of lead from gasoline (4). The debate about the health risks of methyl mercury in fish is essentially a debate about similar shifts in measures of neurobehavioral development (*ScienceScope*, 18 Sept., p. 1779). The health risks of polychlorinated biphenyls (PCBs) and endocrine disruptors can be viewed from the same vantage point (5). Environmental health protection, so to speak, is a nickel-and-dime business.

Bernard Weiss

Department of Environmental Medicine, University of Rochester School of Medicine and Dentistry Rochester, NY 14642, USA. E-mail: weiss@envmed.rochester.edu

References

1. B. Weiss, *Trends Pharmacol. Sci.* **9**, 59 (1988).
2. ——— in *Cocaine Mothers and Cocaine Babies: The Role of Toxins in Development*, M. Lewis and M. Bendersky, Eds. (Erlbaum, Hillsdale, NJ, 1995), pp. 41–55.
3. Centers for Disease Control, *Preventing Lead Poisoning*

in *Young Children* (U.S. Department of Health and Human Services, Washington, DC, 1991).

4. *The Benefits and Costs of the Clean Air Act, 1970 to 1990* (U.S. Environmental Protection Agency, Washington, DC, 1996).
5. B. Weiss, *Neurotoxicology* **18**, 581, (1997); J. L. Jacobson and S. W. Jacobson, *N. Engl. J. Med.* **335**, 783 (1996).

An Early Snowball Earth?

In their article "A Neoproterozoic snowball Earth" (*Reports*, 28 Aug., p. 1342), Paul F. Hoffman *et al.* report that global ice-house conditions existed during the Proterozoic, as inferred from negative carbon isotopes in carbonate rocks from Namibia. These conditions are said to have led to the near termination of life on Earth. In summary, the hypothesis suggests that global glaciation existed until volcanic outgassing increased carbon dioxide (CO₂) concentrations to 120,000 parts per million of volume, at which time the global ice-house conditions collapsed. Several issues of geology and climate, however, remain unresolved.

First, how did global glacial conditions come about? Reduced solar forcing could not have been the cause, because the solar constant was lower before the breakup of the Rodinia supercontinent. A positive ice-albedo feedback triggered by reduced CO₂


MRC Presents: SUPER AGAROSE

Combining Convenience and Performance

Convenience • First agarose suspension available
• No weighing of powdered agarose

Performance • Exceptional resolution of DNA and RNA
• Recommended for Southern & northern blotting, PCR, RPA and RFLP

Procedure • Dilute with buffer
• Boil and pour gel



Markers and genomic DNA in 1% Super Agarose

Markers and PCR product in 2% Super Agarose

Markers and total RNA in 1% Super Agarose

MRC
MOLECULAR RESEARCH CENTER, INC
5645 Montgomery Rd.
Cincinnati, OH 45212
Phone: 513-841-0900
888-841-0900
Fax: 513-841-0080
E-mail: info@mrcgene.com
Visit our website at:
<http://www.mrcgene.com>
For international distributors,
contact MRC.

SCIENCE'S COMPASS

concentrations or high obliquity has been suggested (1). However, there are problems with both of these hypotheses.

It would have been difficult for the ocean to completely freeze because of its large thermal capacity and the wind-driven and thermohaline circulation. The equatorial ocean would have been buffered from sea ice by the high, year-round incident solar radiation at the surface, and therefore very thick ice would have needed to be present. Early studies of energy balance models suggest that a 5% reduction could generate a snowball Earth; however, more recent studies suggest that the solar forcing must be reduced by more than 10% to produce global glaciation by means of the ice-albedo feedback mechanism when large expanses of ocean exist (2). As for a high obliquity, there is no definitive evidence to suggest that Earth's obliquity was very different from that of the present or that high-latitude temperatures were warmer than those of low latitudes during the Neoproterozoic. Finally, there is evidence of low-latitude carbonates and evaporites along with glacial deposits during the late Proterozoic (3).

It would not have been possible for an idealized tropical supercontinent with fixed sea-surface temperatures (SSTs) (4) to maintain subfreezing temperatures on land, primarily because of the large quantity of solar radiation reaching the surface. Further snow-albedo feedback would not have been possible because moisture cannot penetrate the interiors of the supercontinent that would have allowed snow to accumulate, even if surface temperatures had been cold enough. It is possible that the results of Hoffman *et al.* indicate regional effects. The Congo Craton underwent significant changes between 700 and 600 million years ago as a result of a collision with East and West Gondwanaland (5). The Pan-African orogeny, which may have been of Himalayan proportions (6), could have produced alpine glaciers in the tropics. One possible hypothesis is that the collision led to a rapid draw down of CO₂ as a result of intense weathering because of its location in the deep tropics. Alpine glaciers then migrated to lower elevations.

Gregory S. Jenkins

Department of Meteorology, Pennsylvania State University, University Park, PA 16803, USA. E-mail: osei@essc.psu.edu

Christopher R. Scotese

Department of Geology, University of Texas, Post Office Box 19049, Arlington, TX 76019, USA

References and Notes

1. G. E. Williams, *Earth Sci. Rev.* **34**, 1 (1993).
2. G. S. Jenkins, *Global Planet. Change* **7**, 321 (1993); B. Longdoz and L. M. François, *ibid.* **14**, 97 (1997).
3. N. M. Chumakov and D. P. Elson, *Episodes* **12**, 150 (1989); A. Ronov, V. Khain, K. Seslavinsky, *Atlas of Lithological-Paleogeographical Maps of the World* (Mingeo SSSR, Leningrad, 1984), Late Precambrian and Paleozoic maps of the continents.
4. G. S. Jenkins and L. A. Frakes, *Geophys. Res. Lett.* **25**, 3525 (1998).
5. R. J. Stern, *Annu. Rev. Earth Planet. Sci.* **22**, 319 (1994).
6. C. R. Scotese, *J. African Earth Sci.* **27**, 171 (1998).

Response

Jenkins and Scotese question the existence of a Neoproterozoic snowball Earth on the grounds of climate model simulations (1, 2) and suggest that the evidence from Namibia can be reinterpreted in terms of alpine glaciation associated with the tectonic assembly of Gondwana. Climate model results depend on prescribed boundary conditions.

Two of the studies (1) cited by Jenkins and Scotese are aimed at the "faint young sun" climatic paradox for the Archean Earth and prescribe a global ocean without continents. The other study (2), which appeared after our report in *Science*, addresses the problem of Neoproterozoic low-latitude glaciation on a supercontinent with a general circulation model (GCM). All of the models share a common trait, as does an earlier attempt

to model Neoproterozoic low-latitude glaciation (3)—that changes in tropical SST are very small relative to changes in insolation (which are believed to have been lower in the past because of stellar evolution). In order to simulate the effect of a 5% reduction in the solar constant, Jenkins and Frakes (2) artificially reduced present-day monthly SSTs by 2°C at all latitudes. Given that tropical SSTs were actually 3° to 5°C colder than present at the last glacial maximum (21,000 years ago) (4), when the difference in insolation was less than 1%, there is good reason to question the prescribed response of SSTs to solar forcing in the models: in the Quaternary, the climatic response to orbital forcing was strongly amplified by changes in the levels of greenhouse gases (5). As the causal link between orbital forcing and greenhouse gas concentrations remains elusive, there are no grounds for its exclusion from Neoproterozoic climate change. In any event, it is hardly surprising that low-latitude glaciation does not occur in a GCM that has tropical SSTs prescribed only 2°C lower than present (2). Jenkins and

Science

Notice to Readers

On 23 November 1998, *Science's* European headquarters moved to the following new address:


**Science International
Bateman House
82-88 Hills Road
Cambridge CB2 1LQ
United Kingdom**

Tel: +(44)(0) 1223 326500
Fax: +(44)(0) 1223 326501
E-mail: science@science-int.co.uk



Filter in half the time with the Stericup™ vacuum filter unit. Our patented fast-flow, low protein-binding Millipore Express™ membrane gives you faster filtration with bacterial retention. Features include a unique no-tip, easy-grip design; a recessed bottom that allows stacking for convenient storage; and a tab inside the funnel that holds a prefilter securely in place.

Volume filtered through Stericup vs. Brand N cup in 75 seconds, DME with 10% serum (FBS)




To place an order in the US, call Fisher Scientific at 800-766-7000 (800-234-7437 in Canada). In Europe fax +33 3.88.38.91.95. In Japan call (03) 5442-9716. In Asia call (852) 2803-9111. For more information call Technical Service at 800-MILLIPORE. Circle No. 24 on Readers' Service Card www.millipore.com/sterile


Carbohydrate Analysis

As easy as...


1.



2.



3.



Glyko's FACE (Fluorophore Assisted Carbohydrate Electrophoresis) technology includes everything you need to work with and analyze complex carbohydrates:

- **KITS:** precast gels, gel buffer, all reagents and manuals.
- **IMAGERS:** 14-bit camera, FACE software, enhanced band-finding and resolution of minor bands (sensitivity of 5 pmols). Also use for protein and DNA gel imaging.
- **SYSTEMS:** Pentium MMX 166MHz computer with a 2GB hard drive, 32MB RAM, internal Zip drive, fax/modem, CD-ROM, color ink jet printer and FACE software.
- **EXPERT TECHNICAL HELP**
- **OTHER PRODUCTS AND SERVICES:**
 - Recombinant glycosidases
 - Analytical service for general FACE applications
 - Clinical lab service for Lysosomal storage diseases, plasma/urine levels of heparin
 - QC validation with FACE

1 800 33 GLYKO (334 5956) U.S. only
Phone: 1 415 382 6653, Fax: 1 415 382 7889
www.glyko.com



©1998 GLYKO, INC.
FACE is a registered trademark of Glyko, Inc. PENTIUM is a registered trademark of Intel Corp.

SCIENCE'S COMPASS

Frakes acknowledge (2) that "by prescribing SSTs we eliminate any possibility of "ice-house" conditions through a positive ice-albedo feedback at oceanic grid points." In their view, this constraint is justified "by the geologic rock record which does not indicate global glaciation during the Proterozoic or at any other period in Earth's history." We read the rock record differently.

An alpine glacial interpretation can be ruled out for reasons given in our original report: there were no mountains in the region at the time of the Ghaub glaciation. The glacial deposits occur within a broadly conformable succession of shelf carbonates, extending for more than 400 kilometers landward of the shelf edge where our study was conducted (6). Virtually all the glacial debris is from sources on the carbonate shelf. Mountains did form during the assembly of Gondwana, which involved the closure of oceans flanking the Congo Craton. Our study area, located on the southwestern salient of the craton, was directly affected by closure of the ocean to the west and south (present-day coordinates). The first deformation related to this closure [less than 550 million years ago (7)] folded the entire shelf carbonate suc-

cession enclosing the glacial deposits and clearly postdates the glaciation, which occurred 700 to 750 million years ago. The eastern ocean closed in two stages—680 million years ago and less than 550 million years ago (7), and the mountain belt related to its closure was located 1900 kilometers east of our study area. The dramatic rise in the ratio of strontium-87 to strontium-86 in the Neoproterozoic ocean that is commonly taken to signal an erosion flux associated with the Pan-African orogeny occurs only after the last of the proposed snowball glaciations—580 million years ago (8).

The snowball Earth hypothesis explains the most salient features of the Neoproterozoic sedimentary record, including low-latitude glaciations, banded iron-formations, cap carbonates, and carbon isotopic excursions, but Jenkins and Scotese offer no alternative explanation.

Paul F. Hoffman

Daniel P. Schrag

Galen P. Halverson

Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138, USA

J. Alan Kaufman

Department of Geology, University of Maryland, College Park, MD 20742, USA

References and Notes

1. G. S. Jenkins, *Global Planet. Change* **7**, 321 (1993); B. Longdoz and L. M. François, *ibid.* **14**, 97 (1997).
2. G. S. Jenkins and L. A. Frakes, *Geophys. Res. Lett.* **25**, 3525 (1998).
3. T. J. Crowley and S. K. Baum, *J. Geophys. Res.* **98**, 16723 (1993).
4. T. Van der Hammen, *J. Biogeogr.* **1**, 3 (1974); S. C. Porter, *Quat. Res.* **12**, 161 (1979); M. Stute, P. Schlosser, J. F. Clark, W. S. Broecker, *Science* **256**, 1000 (1992); T. P. Guilderson, R. G. Fairbanks, J. L. Rubenstein, *ibid.* **263**, 663 (1994); D. P. Schrag, G. Hampt, D. W. Murray, *ibid.* **272**, 1930 (1996).
5. J. M. Bamola, D. Raynaud, Y. S. Korotkevich, C. Lorius, *Nature* **297**, 391 (1987); D. Raynaud, J. Chappellaz, J. M. Barnola, Y. S. Korotkevich, C. Lorius, *ibid.* **333**, 655 (1988).
6. R. M. Miller, in *African Basins*, vol. 3, *Sedimentary Basins of the World*, R. C. Selley, Ed. (Elsevier, Amsterdam, 1997), pp. 237–268.
7. J. G. Meert and R. Van der Voo, *Geodynamics* **23**, 223 (1997).
8. A. J. Kaufman, A. H. Knoll, G. M. Narbonne, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 6600 (1997).

CORRECTIONS AND CLARIFICATIONS

In the 13 November NetWatch (p. 1223), the URL for the Left Handed DNA Hall of Fame should have been given as "www.lecb.ncifcrf.gov/~toms/LeftHanded.DNA.html."

In the third line of the legend of table 1 (p. 738) in the report "Rapid identification of subtype-selective agonists of the somatostatin receptor through combinatorial chemistry" by S. P. Rohrer *et al.* (23 Oct., p. 737), "(in nanomoles)" should have read, "(nM)." In the first footnote in the legend of table 2 (p. 739), "(in M)" should have read, "(nM)."

1999 AAAS WILLIAM D. CAREY ANNUAL SCIENCE AWARD

The AAAS William D. Carey Annual Science Award was established in 1989 with funds donated by Bill Golden and David Hamburg to encourage advanced students in science or engineering to participate in the AAAS Annual Meeting. This award provides up to \$300 support for students to attend the 1999 AAAS Annual Meeting. In addition, AAAS will furnish registration fees and a year's subscription to *Science*.

Graduate students are invited to apply for the 1999 William D. Carey Annual Science Award. Consideration for this award will be based on academic achievement and other accomplishments related to the advancement of science. Applicants should submit a curriculum vitae and a 200–300 word description of their research topics to:

AAAS Carey Annual Science Award
1200 New York Ave., NW
Washington, DC 20005
202-326-6670
FAX: 202-371-9849

Deadline is December 18, 1998.