



## Review

## Gene therapy for brain tumors: Basic developments and clinical implementation

Hikmat Assi<sup>a,b,c</sup>, Marianela Candolfi<sup>d</sup>, Gregory Baker<sup>a,b,c</sup>, Yohei Mineharu<sup>c</sup>,  
Pedro R. Lowenstein<sup>a,b</sup>, Maria G. Castro<sup>a,b,\*</sup>

<sup>a</sup> Department of Neurosurgery, University of Michigan Medical School, MSRBII, Rm 4570, 1150 Medical Center Drive, Ann Arbor, MI 48109, United States

<sup>b</sup> Department of Cell and Developmental Biology, University of Michigan Medical School, MSRBII, Rm 4570, 1150 Medical Center Drive, Ann Arbor, MI 48109, United States

<sup>c</sup> Department of Molecular and Medical Pharmacology, David Geffen School of Medicine, UCLA, Los Angeles, CA, United States

<sup>d</sup> Instituto de Investigaciones Biomedicas, Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina

## HIGHLIGHTS

- ▶ GBM models that reproduce the salient features of the human disease still need to be developed.
- ▶ TLR2 activation is critical for initiating anti-brain tumor specific adaptive immune responses.
- ▶ Combined cytotoxic and immune-stimulatory approaches are most effective against brain tumors.
- ▶ Viral vectors are safe to administer to human patients diagnosed with GBM.

## ARTICLE INFO

## Article history:

Received 2 May 2012

Accepted 3 August 2012

## Keywords:

Gene therapy  
Glioblastoma  
Cytotoxic  
Immunotherapy  
Viral vectors  
TK  
Flt3L

## ABSTRACT

Glioblastoma multiforme (GBM) is the most common and deadliest of adult primary brain tumors. Due to its invasive nature and sensitive location, complete resection remains virtually impossible. The resistance of GBM against chemotherapy and radiotherapy necessitate the development of novel therapies. Gene therapy is proposed for the treatment of brain tumors and has demonstrated pre-clinical efficacy in animal models. Here we review the various experimental therapies that have been developed for GBM including both cytotoxic and immune stimulatory approaches. We also review the combined conditional cytotoxic immune stimulatory therapy that our lab has developed which is dependent on the adenovirus mediated expression of the conditional cytotoxic gene, Herpes Simplex Type 1 Thymidine Kinase (TK) and the powerful DC growth factor Fms-like tyrosine kinase 3 ligand (Flt3L). Combined delivery of these vectors elicits tumor cell death and an anti-tumor adaptive immune response that requires TLR2 activation. The implications of our studies indicate that the combined cytotoxic and immunotherapeutic strategies are effective strategies to combat deadly brain tumors and warrant their implementation in human Phase I clinical trials for GBM.

© 2012 Elsevier Ireland Ltd. All rights reserved.

## Contents

1. Introduction .....	72
2. Experimental glioblastoma models .....	72
2.1. Transplantable murine GBM models .....	72
2.2. Genetically engineered murine GBM models .....	73
3. Cytotoxic gene therapy .....	73
4. Immune mediated gene therapy: stimulating anti-brain tumor immunity .....	74
5. The clinical scenario .....	75
Funding .....	76
References .....	76

\* Corresponding author at: MSRB II, Room 4570, University of Michigan School of Medicine, 1150 West Medical Center Drive, Ann Arbor, MI 48109-5689, United States.  
Tel.: +1 734 764 0850; fax: +1 734 764 7051.

E-mail address: [maricacas@umich.edu](mailto:maricacas@umich.edu) (M.G. Castro).

## 1. Introduction

Glioblastoma multiforme (GBM) is a highly malignant brain tumor of astrocytic origin. In the United States, GBM accounts for over 50% of all gliomas and carries an annual incidence rate of 3.2 cases per 100,000 persons, making it both the most common and lethal primary brain tumor in adults [14]. The World Health Organization classifies GBM as a grade IV malignant neuroepithelial cancer of the central nervous system with two distinct variants: giant cell glioblastoma and gliosarcoma [47]. Although histological differences exist between its various forms, GBM is by definition a highly mitotic and diffusely infiltrating glial anaplasia that demonstrates marked nuclear and cytoplasmic pleomorphism. Other defining histological features of GBM include prominent glomeruloid microvascular hyperplasia and central areas of tumor necrosis that are often, but not necessarily, associated with perinecrotic nuclear pseudopalisading [71]. Genetically, GBM displays heterogeneous alterations in a number of key bio-molecular pathways implicated in processes of cellular proliferation, survival, invasion and angiogenesis [28]. The current gold standard of therapy for GBM consists of surgical resection when feasible followed by chemo and radiotherapy [74]. The oral alkylating agent temozolomide, introduced in 2005, is now the cytotoxic drug of choice for patients with newly diagnosed GBM due to its additional role as a radio-sensitizer when administered concurrently with radiotherapy [70].

GBM presents a number of significant drawbacks to therapy. Widespread tumor cell migration occurring predominantly in association with white matter tracts and the basement membrane of brain microvasculature preclude complete surgical resection and invariably leads to tumor recurrence, frequently within 2–3 cm of the resection cavity [31]. Another significant drawback to therapy is the inherent resistance of GBM cells to cytotoxic therapies. Constitutive up-regulation of DNA repair enzymes such as O6-methylguanine-DNA methyltransferase (MGMT) and activation of anti-apoptotic regulators such as Bcl2-like 12 (Bcl2L12) allow resistant tumor cells to persist in the face of chemo and radiotherapy [34,69]. Clinically, it has been determined that GBM patients with high levels of MGMT promoter methylation survive significantly longer than those without evidence of MGMT promoter methylation [44].

The rampant vascular endothelial proliferation seen in high-grade gliomas coupled with the phenomenon of tumor dormancy, the inability of solid tumors to grow larger than a few mm<sup>3</sup> in the absence of sprouting angiogenesis [26,27], served as the impetus for the use of anti-angiogenic agents in the treatment of GBM. However, clinical studies show that its use in the adjuvant setting along with current first-line chemotherapies only modestly increases progression-free survival and provides no additional survival benefit in patients with GBM [32,46,56,57]. Furthermore, the FDA has recently announced that it has removed bevacizumab's indication for HER2-negative metastatic breast cancer, stating that a detailed analysis of the clinical data revealed a lack of therapeutic efficacy. The effects of anti-angiogenic therapy are likely attributed to those of vascular normalization, a phenomenon that temporarily reduces peri-tumoral vasogenic edema and leads to improved patient symptoms and quality of life [40]. It has also been suggested that vascular normalization acts by allowing for better penetration of systemically administered chemotherapeutic agents into the tumor bed, but only over a relatively narrow window of time [40].

Despite recent advances made in the field of neuro-oncology, GBM remains a uniformly lethal disease with a dismal prognosis. Even with optimal therapeutic intervention, median patient survival continues to be 12–15 months post-diagnosis [74]. GBMs aggressive growth and invasion, coupled with its inherent

resistance to cytotoxic therapy, make it a formidable opponent even for today's optimal therapeutic modalities. Thus the field is in great need for new and novel tumor-specific therapies.

## 2. Experimental glioblastoma models

One of the most important tools in the development of translational therapeutics is the availability of adequate animal models to preclinically test the efficacy and toxicity of novel therapies. The ideal animal model to test anti-glioma therapies comprises the following features: (1) histopathological resemblance to human GBM, i.e. similar invasion pattern; (2) biochemical resemblance to human GBM, i.e. similar genetic lesions; (3) intact tumor–host biochemical and immunological interactions, i.e. the tumor needs to be non-immunogenic in a host with an intact immune system; (4) intracranial location, so that the tumor is surrounded by normal brain parenchyma; (5) accurate knowledge of tumor location; (6) predictable growth pattern; (7) high reproducibility. Unfortunately, a GBM animal model that comprises all these features does not exist and, thus, experimentation in different tumor models is advised in order to better predict the clinical outcome of translational therapies.

### 2.1. Transplantable murine GBM models

The first brain tumor models were generated by systemic administration of chemical carcinogens to rodents [43]. Although this strategy is no longer utilized, glioma cell lines obtained by this method [6,7] are still extensively used for in vivo tumor models, developed by intracranial or subcutaneous (s.c.) implantation in rodents. Although s.c. GBM models allow to follow tumor growth by daily measurement using a caliper and are a faster and easier alternative to intracranial tumor implantation, the lack of surrounding non-neoplastic brain parenchyma, the absence of a blood–brain barrier and the immune-privilege present in the brain make s.c. models unsuitable to assess the efficacy or the neurotoxicity of anti-glioma therapeutic approaches.

The advantages of intracranially implanted tumor models are their predictable and highly reproducible tumor growth rates, the accurate knowledge of the site of the tumor, the possibility of testing a large cohort of animals, and the relatively fast progression from tumor implantation to death [9], which make *them an invaluable* tool for the preclinical assessment of novel therapies. Syngeneic GBM models are generated by implantation of murine GBM cell lines that are not immunogenic when implanted in animals with an intact immune system [9]. Syngeneic mouse models of GBM are not abundant, and they are constituted by the following cell lines implanted in their corresponding mouse host: GL26 and GL261 GBM cells in C57BL6 mice [9,19], SMA-560 cells in VMDK mice [10] and VM-M3 in VM mice [65]. Amongst the syngeneic rat GBM models, the most extensively used are CNS-1 cells in Lewis rats [9], F98, 9L and RG-2 cells in Fisher rats [5]. The integrity of the immunological interaction between host and GBM makes these models an excellent tool to study antitumor immunity, as well as the efficacy and toxicological profile of immunotherapeutic approaches for GBM [19].

Xenograft models allow assessing the response of human GBM cells in the context of the normal brain, and have been extensively used as preclinical in vivo models. Although the hosts are immune-compromised mice and rats, human GBM xenografts require the injection of much larger number of cells than syngeneic models to growth with reproducible rates [9]. Besides the obvious limitation of xenograft models, which is the lack of an intact immune system, there is an additional question that needs to be addressed when choosing a human GBM xenograft: some of the main genetic

lesions detected in the original GBM specimens, such as EGFR amplification and hypermethylation of the DNA O6-methylguanine methyltransferase (MGMT) promoter, can be lost after prolonged cell culture [13]. This constitutes a concern when using human GBM cell lines that have been maintained in culture for years. In order to address this limitation patient tumor specimens have been implanted directly in the flank of nude mice and maintained by serial transplantation in vivo [13]. These tumors can be cryopreserved or cultured for short periods before injecting the cells into the brain of immune-compromised mice [30]. We have recently employed one of these transplantable human tumors, GBM12, which retains EGFR amplification [30], p53 mutation [30], and expression of IL13R $\alpha$ 2 [3] from the original GBM specimen, to address the efficacy of a targeted toxin delivered using a regulated adenoviral vector [11].

The main limitation of implantation tumor models is that although they resemble the histopathological features of human GBM, they do not replicate exactly their invasive pattern, being less diffuse than their human counterpart [9]. Also, glioma-genesis is artificially achieved and does not resemble the pathogenesis of the human disease. In spite of these shortcomings, implantation models serve as a reliable tool in translational neuro-oncology that allows the preclinical assessment of novel therapies.

## 2.2. Genetically engineered murine GBM models

Genetically engineered murine GBM models mimic gliomagenesis more accurately and exhibit the histological and molecular hallmarks of human GBM. Transgenic mouse models have been constructed by introducing genetic alterations known to be present in human gliomas. Although the alteration of a single tumor suppressor gene or overexpression of an oncogene is insufficient to induce high-grade gliomas with good penetrance, the introduction of more than one genetic lesion found in human GBM leads to murine glioma models that resemble the main histological features of their human counterparts.

Amplification of growth factors and their receptors, i.e. PDGF and EGFR, or silencing of tumor suppressor genes, i.e. p53 and PTEN, are detected in the majority of human GBM specimens [52,62,64,77]. The intracranial delivery of these genes into the brain of pre-natal or adult rodents using gene therapy vectors that integrate into the host genome has been used to generate endogenous brain tumors with variable success. Delivery of retroviral vectors that encode for PDGF into the rat adult brain or in the newborn mouse brain leads to the formation of GBM in less than 20 days [4] or 4–12 weeks [66], respectively. Generation of models that harbor combined genetic lesions mimics more closely the clinical scenario [58]. These models have been generated by delivering retroviral vectors encoding for growth factors (bFGF) or their constitutively activated receptors (EGFRvIII) and cyclin dependent kinases (cdk4) in the brain of p53 deficient mice [36]. While mice that harbor single genetic lesions did not develop brain tumors, combination of genetic lesions led to 50% of the mice [36].

Since Ras activation has been involved in gliomagenesis [37], this molecule has been targeted in order to develop novel endogenous mouse models [50,51]. Constitutive Ras activation in neural stem cells leads to the generation of Grade III astrocytomas in mice, which evolve to Grade IV astrocytomas when p53 and p16/p19 are suppressed [51]. Delivery of Ras and AKT to specific areas of the brain has also been achieved using lentiviral vectors [50]. Although administration of single oncogenes did generate tumor in some of the animals, combination of Ras and AKT in p53 KO mice led to tumor formation in 75–100% of the mice injected, depending on the area injected [50], supporting the notion that gliomagenesis requires several genetic abnormalities occurring in definite areas of the brain.


The use of the *Sleeping Beauty* (SB) transposable element also allows integration of known oncogenes in the genome of brain cells [75]. SB is a synthetic transposable element constituted by a transposon DNA substrate and a transposase enzyme. SB transposase mediates the excision and insertion of transposon DNA into the host genome [53]. Delivery of SB-dependent plasmids encoding for AKT, Ras, EGFRvIII, and a p53 shRNA into the brain of neo-natal mice led to brain tumor formation [75]. The combination of Ras, EGFRvIII and p53 shRNA generated tumors in 100% of the mice that had a median survival of 83 days. An advantage of the SB system is that allows integration of large transposons (<10 kb) into the genome of many different strains of mice [75].

The main limitations of genetically engineered GBM models that restrict their use in translational neuro-oncology are: their variable reproducibility, their long tumor latency and the lack of an accurate knowledge of tumor location. Nevertheless, the fast development of novel imaging techniques [22] allows following tumor growth and assessing its exact location, facilitate the use of genetically engineered models in the preclinical testing of translational therapeutics.

## 3. Cytotoxic gene therapy

Cytotoxic agents have been traditionally used to treat cancer [17,63]. However, one of the therapeutic challenges of treating GBM is that pro-apoptotic agents may be cytotoxic to the surrounding brain parenchyma. The neurotoxicity of pro-apoptotic agents can be reduced by targeting the cytotoxic molecule to GBM cells, which can be achieved using different strategies. The use of gene therapy vectors encoding the therapeutic transgene under the control of a tumor-specific promoter allows restricting the expression of cytotoxic agents to tumor cells. Likewise, p53-upregulated modulator of apoptosis (PUMA) induces GBM-cell death when its expression is controlled by the hTERT promoter, without affecting surrounding non-neoplastic tissue [39]. Vectors encoding pro-apoptotic molecules, such as Bax, under the control of hypoxia responsive elements are also useful to target cytotoxic gene expression to GBM cells within the hypoxic tumor microenvironment [55]. Radio-inducible promoters allow temporal and spatial control of cytotoxic transgene expression. Overexpression of caspase 8 or TRAIL under the control of a radiation-inducible promoter (early growth response gene-1, EGR-1) induces GBM cell death and tumor regression only when combined with fractionated radiotherapy [72]. Restriction of transgene expression to GBM cells becomes crucial when delivering powerful pro-apoptotic molecules into the brain, such as FasL or TRAIL which are highly toxic to the non-neoplastic brain parenchyma [12].

Another strategy to restrict the cytotoxicity of pro-apoptotic molecules to GBM cells is to target receptors that are exclusively expressed on tumor cells and are absent in the normal brain. Chimeric proteins conformed by ligands of these receptors, i.e. IL-13R $\alpha$ 2, transferrin receptor, EGFR, fused to highly cytotoxic proteins, such as *Pseudomonas* exotoxin. We constructed a doxycycline-dependent regulatable adenoviral vector (Ad.mhIL-4.TRE.mhIL-13-PE) that encodes a mutated human IL-13 fused to *Pseudomonas* exotoxin (mhIL-13-PE) that specifically binds to IL13R $\alpha$ 2 [11,20], an IL13 receptor that is overexpressed in GBM in most human patients [41,54,76]. Ad.mhIL-4.TRE.mhIL-13-PE also encodes a mutated human IL-4 that binds to the physiological receptor IL4R/IL13R without interacting with IL13R $\alpha$ 2 [20,49], to block any potential binding of mhIL-13-PE to normal brain cells. This therapeutic vector exhibited higher efficacy and negligible neurotoxicity when compared to the protein formulation, i.e. Cintredekin Besudotox [11], which when tested in clinical trials failed to achieve clinical endpoints and revealed severe neurotoxicity



**Fig. 1.** Diagram illustrating the effects of adenoviral mediated TK/Flt3L gene therapy. Intra-tumoral injections of adenovirus expressing TK in combination with adenovirus expressing Flt3L into a brain tumor induce tumor cell death, release of intracellular inflammatory molecules, such as HMGB1 and tumor antigen. Flt3L recruits DCs to the tumor site, where they phagocytose tumor cell remnants and migrate to the dLN followed by priming of a T cell mediated cytotoxic anti-tumor immune response. *Abbreviations:* Tu, tumor; NK, natural killer cell; dLN, draining lymph node; TLR, toll like receptor; GCV, Ganciclovir; TK, thymidine kinase; Flt3L, fms-like tyrosine kinase 3 ligand; HMGB1, high-mobility group box-1.

[45]. Thus, gene therapy vectors emerge as a useful alternative to deliver these cytotoxins into the GBM microenvironment.

#### 4. Immune mediated gene therapy: stimulating anti-brain tumor immunity

To protect itself against damage the mammalian CNS has evolved various mechanisms to restrict the function of the immune system. The brain occupies a niche in the body where the immune system has limited capacity to detect and eliminate foreign antigens. This is referred commonly to as “immune privilege”. Experimentally, it has been significantly more difficult to elicit immune responses towards CNS antigens in comparison to peripheral antigens. A number of key physiological processes that contribute to the suppression of CNS immunity have been identified, i.e. paucity of dendritic cells (DCs) in the brain, lack of lymphatic drainage, production of anti-inflammatory mediators such as TGF-B and NO by cells in CNS including low major-histocompatibility (MHC-II) expression on infiltrating microglia. These mechanisms are ways of protecting vital organs from immune-mediated attack [24].

The immune system is a key determinant of tumor rejection and escape. Therefore one of our research goals was to try and overcome this so-called immune privilege and in the process discover novel immune mediated gene therapeutic approaches for the treatment of GBM. As the brain is virtually absent of any dendritic cells, it is difficult to prime an adaptive immune response against antigens that are exclusive to the brain. Our goal was to generate a tumor micro-environment that is conducive to dendritic cell migration and maturation in the hopes of rescuing animals from lethal models

of GBM. We have succeeded in developing an adenoviral mediated immunotherapy for brain tumors that is dependent on the expression of two genes: *Thymidine Kinase* (TK), phosphorylates the prodrug Ganciclovir which induces DNA crosslinking followed by cell death [2] and *fms-like tyrosine kinase-3 ligand* (Flt3L), a potent DC growth factor that serves to increase the number of infiltrating DCs within the tumor microenvironment [1]. Tumor cell death induced by TK causes the release of tumor antigens, which are phagocytosed by surveying DCs and transported to the draining lymph nodes. Here T cells are primed to elicit an antigen specific cytotoxic anti-tumor immune response (Fig. 1). Combination therapy using the two adenoviruses induces tumor regression, long term survival and immunological memory in several mouse and rat GBM models [2,19].

Toll-like receptor (TLR) signaling; specifically TLR2 in DCs is essential for priming an effective immune response [19]. Using TLR2 KO mice, we demonstrated a defect of DCs to migrate to the tumor microenvironment when treated with adenovirus expressing TK/Flt3L. In addition, DCs lacking TLR2 failed to stimulate the proliferation and activation of tumor antigen specific T cells. The loss of TLR2 signaling abolished the CD8+ cytotoxic T cell mediated response seen in wild type mice [19]. These data highlight the importance of a receptor thought to be involved in innate immunity orchestrating an adaptive anti-brain tumor immune response.

But if this classical danger sensing receptor is required for initiating the immune response then surely there must be a tumor derived ligand. We identified high-mobility-group box 1 (HMGB1), a protein thought to be involved in controlling inflammation and sepsis as the TLR ligand in our model [19]. In viable cells HMGB1 is part of the chromatin structure but in response to

pro-apoptotic stimuli, including TK (plus GCV) mediated gene therapy, radiotherapy and TMZ, GBM cells shed nuclear HMGB1 [19], suggesting that cytotoxic agents may also be suitable adjuvants for immunotherapeutic strategies. HMGB1 binds TLR2 on bone marrow derived DCs stimulating their maturation. Neutralizing HMGB1 through small molecule inhibitors or antibodies leads to a loss of TLR2 stimulation, subsequent DC anergy and loss of therapeutic efficacy [19]. Our data suggest that HMGB1 released from a dying tumor is a critical TLR2 ligand that initiates the anti-tumor immune response [12,19]. Understanding how the TLR2 pathway in DCs is activated might potentially yield new targets for eliciting clinically effective anti-brain tumor immune responses.

An important facet to any adaptive immune response is the proper activation of cytotoxic T lymphocytes (CTLs). As such, several other groups have attempted to develop treatments based on T cell augmentation. In a rat glioma model, recombinant vaccinia virus expressing the cytokines IL-2 or IL-12 resulted in inhibition of tumor growth when injected intra-tumorally [16]. These soluble mitogenic factors mediate the growth survival and differentiation of antigenic T cells. As such combination therapies of IL-2 and IL-12 proved to be more effective in increasing the levels natural killer, Mac-1<sup>+</sup>, and NKT cells in blood as well as increased interferon-gamma, and tumor necrosis factor-alpha expression in tumors. Similar results were obtained with intra-tumoral injection of adenovirus expressing IL-12 or IL-2 in a breast cancer model [23].

Tumor necrosis factor-alpha (TNF $\alpha$ ) and interferon-gamma (IFN $\gamma$ ) are potent immune-stimulatory cytokines capable of inducing tumor cell death that have been used in experimental GBM models. IFN $\gamma$  is a key cytokine that promotes Th-1 polarization [25] and is thought to have anti-angiogenic properties [67]. Aside from facilitating immune cell migration, TNF $\alpha$  has been shown to regulate multiple functions of immune cells including cell growth, inflammation, and autoimmunity [15,73]. TNF $\alpha$  has also been shown to cause direct necrosis in neoplastic cells [8]. In a mouse glioma model the use of adenoviral vectors expressing TNF $\alpha$  or IFN $\gamma$  delivered intra-tumorally induced infiltration of CD4<sup>+</sup> and CD8<sup>+</sup> cells in addition to increasing expression of MHC-I and -II on the tumor cells. Intracranial administration of both vectors led to a statistically significant increase in survival of tumor bearing mice [21]. These studies highlight the various ways the immune system has been harnessed to elicit effective albeit experimental treatments for brain tumors. Combination therapies that elicit immunogenic cell death and are immune stimulatory are predicted to be of most clinical benefit.

## 5. The clinical scenario

The relatively high efficiency of preclinical trials of various different therapeutic modalities gave confidence to clinicians and scientists to move ahead into early clinical trials in human patients. The high experimental therapeutic efficacy of TK+GCV in *in vitro* and *in vivo* models of GBM prompted the use of this conditional cytotoxic gene therapy strategy in human patients. The first clinical studies using suicide gene therapy in GBM patients consisted of intra-tumoral injections of the HSV1-TK vector producing cells (VPCs) alongside the standard treatment of surgery and radiotherapy [60]. These patients had increased numbers of IFN- $\gamma$ -producing T cells and an increase in serum IL-12 and FasL levels compared to patients that received standard treatment alone, suggesting that TK (+GCV)-mediated gene therapy can stimulate a Th1 type immune response [60]. These potentially positive results prompted clinicians to test retroviral vectors encoding the suicide gene HSV1-TK in a large Phase III double blinded, controlled clinical trial in patients with glioblastoma. No statistically

significant efficacy benefits were obtained, and the use of retroviral vectors encoding HSV1-TK has now been abandoned. As an alternative, adenoviral vectors encoding the suicide gene HSV1-TK were also tested in small trials and their administration was deemed safe in patients diagnosed with GBM [29,61,68]. In one particular study a clinically and statistically significant increase in mean survival from  $39.0 \pm 19.7$  (SD) to  $70.6 \pm 52.9$  weeks was reported when patients received injections of AdvHSV-TK into the tumor bed following surgical resection [38]. This prompted progression towards a larger Phase III double blinded and controlled trial. Unfortunately, this Phase III trial failed to live up to early clinical success, and did not provide strong evidence of either clinical or statistically significant benefits. In 248 cases with newly diagnosed and previously untreated Glioblastoma multiforme patients received either standard therapy (surgery resection and radiotherapy) or standard therapy combined with adjuvant gene therapy. Twelve months survival rates were 50% vs. 55% in the gene therapy and control groups respectively [59]. Median survival was also similar across the two groups indicating a low therapeutic effect of TK.

These results demonstrate the challenges of translational medicine, and the poor predictability of preclinical science when tested in the human diseases. Secondly, these challenges possibly will work to highlight the need of using multiple agents simultaneously to elicit tumor cell killing and activation of the immune system. One such approach is the combination of immune-stimulatory strategies, coupled to cytotoxic strategies [33,35].

The pleiotropic cytokine IFN $\beta$  has also been tested in a Phase I trial. In a dose escalation study, patients diagnosed with grade 3 or grade 4 brain tumors received doses of adenovirus expressing IFN $\beta$  up to  $2 \times 10^{11}$  viral particles (VPs) alongside standard resection [18]. Doses up to  $5 \times 10^{10}$  VP were deemed safe and were associated with a dose-dependent induction of apoptosis within the tumors. With the exception of one case in which a patient exhibited post-surgical confusion in the high dose cohort, no other adverse events were reported. The therapeutic benefit of IFN $\beta$  gene therapy remains to be seen, and will require multi-center trials in a large number of patients.

The combined Ad-Flt3L+Ad-TK (GCV) gene therapy strategy for the treatment of GBM has received approval from the FDA, for a Phase I dose escalation study in patients newly diagnosed with GBM. This preliminary clinical study will provide information on the safety and efficacy of adenovirus mediated delivery of cytotoxic TK and immune stimulatory Flt3L [42]. In conclusion, the pre-clinical and clinical findings highlight the importance and synergy of tumor cell killing to be used in combination with stimulation of an effective immune response. It is very likely that the combination of cytotoxic agents and immune stimulatory approaches with synchronous TLR2 activation will result in improved therapies for the treatment of brain tumors in humans. The lack of major clinical success highlights the limitations in developing new treatments for brain tumors [48]. Poor clinical responses are seen with most major new drugs being used, and although the contemporary standard of care has pushed median patient survival to 2 years post-tumor resection, it has been difficult to discover any agents that have made a much larger impact in prolonging patients' life. As we improve the preclinical models, their statistical analysis, and also aim to enhance the design, analysis, and interpretation of clinical trials for brain tumors, the challenge to improve patients' lives remains daunting. Though we trust that the combination of powerful immunotherapies and powerful cytotoxic approaches has the potential to improve patients' lives, it will only be clinically and statistically significant Phase III double blind randomized clinical trials that will give us assurance that real progress towards finding a cure has been made.

## Funding

This work was supported by National Institutes of Health/National Institute of Neurological Disorders & Stroke (NIH/NINDS) Grants 1U01-NS052465, U01-NS052465-04S1, 1R21-NS054143, 1R01-NS 074387 and 1R01-NS 061107 to M.G.C.; NIH/NINDS Grants 1R01-NS 054193 and 1R01-NS 057711 to P.R.L.; the Department of Neurosurgery, University of Michigan School of Medicine and The Phase One Foundation to M.G.C. and P.R.L. M.C. was supported by the National Council of Science and Technology (CONICET, Argentina).

## References

- [1] S. Ali, J.F. Curtin, J.M. Zirger, W. Xiong, G.D. King, C. Barcia, C. Liu, M. Puntel, S. Goverdhan, P.R. Lowenstein, M.G. Castro, Inflammatory and anti-glioma effects of an adenovirus expressing human soluble Fms-like tyrosine kinase 3 ligand (hsFlt3L): treatment with hsFlt3L inhibits intracranial glioma progression, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 10 (2004) 1071–1084.
- [2] S. Ali, G.D. King, J.F. Curtin, M. Candolfi, W. Xiong, C. Liu, M. Puntel, Q. Cheng, J. Prieto, A. Ribas, J. Kupiec-Weglinski, N. van Rooijen, H. Lassmann, P.R. Lowenstein, M.G. Castro, Combined immunostimulation and conditional cytotoxic gene therapy provide long-term survival in a large glioma model, *Cancer Research* 65 (2005) 7194–7204.
- [3] C. Allen, G. Paraskevaku, I. Iankov, C. Giannini, M. Schroeder, J. Sarkaria, R.K. Puri, S.J. Russell, E. Galanis, Interleukin-13 displaying retargeted oncolytic measles virus strains have significant activity against gliomas with improved specificity, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 16 (2008) 1556–1564.
- [4] M. Assanah, R. Lochhead, A. Ogden, J. Bruce, J. Goldman, P. Canoll, Glial progenitors in adult white matter are driven to form malignant gliomas by platelet-derived growth factor-expressing retroviruses, *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 26 (2006) 6781–6790.
- [5] R.F. Barth, B. Kaur, Rat brain tumor models in experimental neuro-oncology: the C6, 9L, T9, RG2, F98, BT4C, RT-2 and CNS-1 gliomas, *Journal of Neuro-Oncology* 94 (2009) 299–312.
- [6] P. Benda, J. Lightbody, G. Sato, L. Levine, W. Sweet, Differentiated rat glial cell strain in tissue culture, *Science* 161 (1968) 370–371.
- [7] P. Benda, K. Sameda, J. Messer, W.H. Sweet, Morphological and immunochemical studies of rat glial tumors and clonal strains propagated in culture, *Journal of Neurosurgery* 34 (1971) 310–323.
- [8] L. Bertazza, S. Mocellin, The dual role of tumor necrosis factor (TNF) in cancer biology, *Current Medicinal Chemistry* 17 (2010) 3337–3352.
- [9] M. Candolfi, J.F. Curtin, W.S. Nichols, A.G. Muhammad, G.D. King, G.E. Pluhar, E.A. McNeil, J.R. Ohlfest, A.B. Freese, P.F. Moore, J. Lerner, P.R. Lowenstein, M.G. Castro, Intracranial glioblastoma models in preclinical neuro-oncology: neuropathological characterization and tumor progression, *Journal of Neuro-Oncology* 85 (2007) 133–148.
- [10] M. Candolfi, J.F. Curtin, K. Yagiz, H. Assi, M.K. Wibowo, G.E. Alzadeh, D. Foulad, A.G. Muhammad, S. Salehi, N. Keech, M. Puntel, C. Liu, N.R. Sanderson, K.M. Kroeger, R. Dunn, G. Martins, P.R. Lowenstein, M.G. Castro, B Cells Are Critical to T-cell-Mediated Antitumor Immunity Induced by a Combined Immune-Stimulatory/Conditionally Cytotoxic Therapy for Glioblastoma, *Neoplasia* 13 (2011) 947–960.
- [11] M. Candolfi, W. Xiong, K. Yagiz, C. Liu, A.K. Muhammad, M. Puntel, D. Foulad, A. Zadmehr, G.E. Ahlzadeh, K.M. Kroeger, M. Tesarfreund, S. Lee, W. Debinski, D. Sareen, C.N. Svendsen, R. Rodriguez, P.R. Lowenstein, M.G. Castro, Gene therapy-mediated delivery of targeted cytotoxins for glioma therapeutics, *Proceedings of the National Academy of Sciences of the United States of America* 107 (2010) 20021–20026.
- [12] M. Candolfi, K. Yagiz, D. Foulad, G.E. Alzadeh, M. Tesarfreund, A.K. Muhammad, M. Puntel, K.M. Kroeger, C. Liu, S. Lee, J.F. Curtin, G.D. King, J. Lerner, K. Sato, Y. Mineharu, W. Xiong, P.R. Lowenstein, M.G. Castro, Release of HMGB1 in response to proapoptotic glioma killing strategies: efficacy and neurotoxicity, *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research* 15 (2009) 4401–4414.
- [13] B.L. Carlson, J.L. Pokorny, M.A. Schroeder, J.N. Sarkaria, Establishment, maintenance and in vitro and in vivo applications of primary human glioblastoma multiforme (GBM) xenograft models for translational biology studies and drug discovery, in: Editorial Board, S.J. Enna (Ed.), *Current Protocols in Pharmacology*, Chapter 14, Unit 14, 2001, pp. 16–23.
- [14] CBTRUS Statistical Report: Primary Brain and Central Nervous System Tumors Diagnosed in the United States in 2004–2008. [http://www.cbtrus.org/2012-NPCR-SEER/CBTRUS\\_Report\\_2004-2008\\_3-23-2012.pdf](http://www.cbtrus.org/2012-NPCR-SEER/CBTRUS_Report_2004-2008_3-23-2012.pdf) (04.04.12).
- [15] I. Chatzidakis, C. Mamalaki, T cells as sources and targets of TNF: implications for immunity and autoimmunity, *Current Directions in Autoimmunity* 11 (2010) 105–118.
- [16] B. Chen, T.M. Timiryasova, M.L. Andres, E.H. Kajioka, R. Dutta-Roy, D.S. Gridley, I. Fodor, Evaluation of combined vaccinia virus-mediated antitumor gene therapy with p53, IL-2, and IL-12 in a glioma model, *Cancer Gene Therapy* 7 (2000) 1437–1447.
- [17] O.L. Chinot, M. Barrie, S. Fuentes, N. Eudes, S. Lancelot, P. Metellus, X. Muracciole, D. Braguer, L. Ouafik, P.M. Martin, H. Dufour, D. Figarella-Branger, Correlation between O6-methylguanine-DNA methyltransferase and survival in inoperable newly diagnosed glioblastoma patients treated with neoadjuvant temozolomide, *Journal of Clinical Oncology* 25 (2007) 1470–1475.
- [18] E.A. Chiocci, K.M. Smith, B. McKinney, C.A. Palmer, S. Rosenfeld, K. Lillehei, A. Hamilton, B.K. DeMasters, K. Judy, D. Kirn, A phase I trial of Ad.hIFN-beta gene therapy for glioma, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 16 (2008) 618–626.
- [19] J.F. Curtin, N. Liu, M. Candolfi, W. Xiong, H. Assi, K. Yagiz, M.R. Edwards, K.S. Michelsen, K.M. Kroeger, C. Liu, A.K. Muhammad, M.C. Clark, M. Ardit, B. Comin-Anduix, A. Ribas, P.R. Lowenstein, M.G. Castro, HMGB1 mediates endogenous TLR2 activation and brain tumor regression, *PLoS Medicine* 6 (2009) e10.
- [20] W. Debinski, D.M. Gibo, N.I. Obiri, A. Kealher, R.K. Puri, Novel anti-brain tumor cytotoxins specific for cancer cells, *Nature Biotechnology* 16 (1998) 449–453.
- [21] M. Ehteshami, K. Samoto, P. Kabos, F.L. Acosta, M.A. Gutierrez, K.L. Black, J.S. Yu, Treatment of intracranial glioma with in situ interferon-gamma and tumor necrosis factor-alpha gene transfer, *Cancer Gene Therapy* 9 (2002) 925–934.
- [22] M.A. Elmeliyeg, A.M. Carcaboso, L. Chow LM, Z.M. Zhang, C. Calabrese, S.L. Throm, F. Wang, S.J. Baker, C.F. Stewart, Magnetic resonance imaging-guided microdialysis cannula implantation in a spontaneous high-grade glioma murine model, *Journal of Pharmaceutical Sciences* (2011).
- [23] P.C. Emtage, Y. Wan, M. Hitt, F.L. Graham, W.J. Muller, A. Zlotnik, J. Gauldie, Adenoviral vectors expressing lymphotactin and interleukin 2 or lymphotactin and interleukin 12 synergize to facilitate tumor regression in murine breast cancer models, *Human Gene Therapy* 10 (1999) 697–709.
- [24] Z. Fabry, H.A. Schreiber, M.G. Harris, M. Sandor, Sensing the microenvironment of the central nervous system: immune cells in the central nervous system and their pharmacological manipulation, *Current Opinion in Pharmacology* 8 (2008) 496–507.
- [25] M. Feili-Hariri, D.H. Falkner, P.A. Morel, Polarization of naive T cells into Th1 or Th2 by distinct cytokine-driven murine dendritic cell populations: implications for immunotherapy, *Journal of Leukocyte Biology* 78 (2005) 656–664.
- [26] J. Folkman, Tumor angiogenesis: therapeutic implications, *New England Journal of Medicine* 285 (1971) 1182–1186.
- [27] J. Folkman, M. Hochberg, Self-regulation of growth in three dimensions, *Journal of Experimental Medicine* 138 (1973) 745–753.
- [28] F.B. Furnari, T. Fenton, R.M. Bachoo, A. Mukasa, J.M. Stommel, A. Stegh, W.C. Hahn, K.L. Ligon, D.N. Louis, C. Brennan, L. Chin, R.A. DePinho, W.K. Cavenee, Malignant astrocytic glioma: genetics, biology, and paths to treatment, *Genes and Development* 21 (2007) 2683–2710.
- [29] I.M. Germano, J. Fable, S.H. Gultekin, A. Silvers, Adenovirus/herpes simplex-thymidine kinase/ganciclovir complex: preliminary results of a phase I trial in patients with recurrent malignant gliomas, *Journal of Neuro-Oncology* 65 (2003) 279–289.
- [30] C. Giannini, J. Sarkaria, A. Saito, J.H. Uhm, E. Galanis, B.L. Carlson, M. Schroeder, C.D. James, Patient tumor EGFR and PDGFRA gene amplifications retained in an invasive intracranial xenograft model of glioblastoma multiforme, *Neuro-oncology* 7 (2005) 164–176.
- [31] A. Giese, R. Bjerkvig, M.E. Berens, M. Westphal, Cost of migration: invasion of malignant gliomas and implications for treatment, *Journal of Clinical Oncology* 21 (2003) 1624–1636.
- [32] S.A. Grossman, X. Ye, S. Piantadosi, S. Desideri, L.B. Nabors, M. Rosenfeld, J. Fisher, Survival of patients with newly diagnosed glioblastoma treated with radiation and temozolomide in research studies in the United States, *Clinical Cancer Research* 16 (2010) 2443–2449.
- [33] K.J. Harrington, M. Hingorani, M.A. Tanay, J. Hickey, S.A. Bhide, P.M. Clarke, L.C. Renouf, K. Thway, A. Sibtain, I.A. McNeish, K.L. Newbold, H. Goldswieg, R. Coffin, C.M. Nutting, Phase I/II study of oncolytic HSV GM-CSF in combination with radiotherapy and cisplatin in untreated stage III/IV squamous cell cancer of the head and neck, *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research* 16 (2010) 4005–4015.
- [34] M.E. Hegi, A.C. Diserens, T. Gorlia, M.F. Hamou, N. de Tribolet, M. Weller, J.M. Kros, J.A. Hainfellner, W. Mason, L. Mariani, J.E. Bromberg, P. Hau, R.O. Mirimanoff, J.G. Cairncross, R.C. Janzer, R. Stupp, MGMT gene silencing and benefit from temozolomide in glioblastoma, *New England Journal of Medicine* 352 (2005) 997–1003.
- [35] A.B. Heimberger, W. Sun, S.F. Hussain, M. Dey, L. Crutcher, K. Aldape, M. Gilbert, S.J. Hassenbusch, R. Sawaya, B. Schmittling, G.E. Archer, D.A. Mitchell, D.D. Bigner, J.H. Sampson, Immunological responses in a patient with glioblastoma multiforme treated with sequential courses of temozolomide and immunotherapy: case study, *Neuro-oncology* 10 (2008) 98–103.
- [36] E.C. Holland, W.P. Hively, R.A. DePinho, H.E. Varmus, A constitutively active epidermal growth factor receptor cooperates with disruption of G1 cell-cycle arrest pathways to induce glioma-like lesions in mice, *Genes and Development* 12 (1998) 3675–3685.
- [37] J.T. Huse, E.C. Holland, Targeting brain cancer: advances in the molecular pathology of malignant glioma and medulloblastoma, *Nature Reviews Cancer* 10 (2010) 319–331.
- [38] A. Immonen, M. Vapalahti, K. Tyynela, H. Hurskainen, A. Sandmair, R. Vanninen, G. Langford, N. Murray, S. Yla-Herttua, AdvHSV-tk gene therapy with intravenous ganciclovir improves survival in human malignant glioma: a

- randomised, controlled study, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 10 (2004) 967–972.
- [39] H. Ito, T. Kanzawa, T. Miyoshi, S. Hirohata, S. Kyo, A. Iwamaru, H. Aoki, Y. Kondo, S. Kondo, Therapeutic efficacy of PUMA for malignant glioma cells regardless of p53 status, *Human Gene Therapy* 16 (2005) 685–698.
- [40] R.K. Jain, E. di Tomaso, D.G. Duda, J.S. Loeffler, A.G. Sorensen, T.T. Batchelor, Angiogenesis in brain tumours, *Nature Reviews Neuroscience* 8 (2007) 610–622.
- [41] J.S. Jarboe, K.R. Johnson, Y. Choi, R.R. Lonser, J.K. Park, Expression of interleukin-13 receptor alpha2 in glioblastoma multiforme: implications for targeted therapies, *Cancer Research* 67 (2007) 7983–7986.
- [42] G.D. King, K.M. Kroeger, C.J. Breese, M. Candolfi, C. Liu, C.M. Manalo, A.K. Muhammad, R.N. Pechnick, P.R. Lowenstein, M.G. Castro, Flt3L in combination with HSV1-TK-mediated gene therapy reverses brain tumor-induced behavioral deficits, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 16 (2008) 682–690.
- [43] P. Kleihues, K.J. Zulch, S. Matsumoto, U. Radke, Morphology of malignant gliomas induced in rabbits by systemic application of N-methyl-N-nitrosourea, *Zeitschrift für Neurologie* 198 (1970) 65–78.
- [44] D. Krex, B. Klink, C. Hartmann, A. von Deimling, T. Pietsch, M. Simon, M. Sabel, J.P. Steinbach, O. Heese, G. Reifenberger, M. Weller, G. Schackert, Long-term survival with glioblastoma multiforme, *Brain* 130 (2007) 2596–2606.
- [45] S. Kunwar, M.D. Prados, S.M. Chang, M.S. Berger, F.F. Lang, J.M. Piepmeyer, J.H. Sampson, Z. Ram, P.H. Gutin, R.D. Gibbons, K.D. Aldape, D.J. Croteau, J.W. Sherman, R.K. Puri, Direct intracerebral delivery of cintredekin besudotox (IL13-PE38QQR) in recurrent malignant glioma: a report by the Cintredekin Besudotox Intraparenchymal Study Group, *Journal of Clinical Oncology* 25 (2007) 837–844.
- [46] A. Lai, A. Tran, P.L. Nghiemphu, W.B. Pope, O.E. Solis, M. Selch, E. Filka, W.H. Yong, P.S. Mischel, L.M. Liau, S. Phuphanich, K. Black, S. Peak, R.M. Green, C.E. Spier, T. Kolevska, J. Polikoff, L. Fehrenbacher, R. Elashoff, T. Cloughesy, Phase II study of bevacizumab plus temozolomide during and after radiation therapy for patients with newly diagnosed glioblastoma multiforme, *Journal of Clinical Oncology* 29 (2011) 142–148.
- [47] D.N. Louis, H. Ohgaki, O.D. Wiestler, WHO Classification of Tumours of the Central Nervous System, International Agency for Research on Cancer (IARC), Lyon, 2007.
- [48] P.R. Lowenstein, M.G. Castro, Uncertainty in the translation of preclinical experiments to clinical trials. Why do most phase III clinical trials fail? *Current Gene Therapy* 9 (2009) 368–374.
- [49] A.B. Madhankumar, A. Mintz, W. Debinski, Interleukin 13 mutants of enhanced avidity toward the glioma-associated receptor, IL13Ralpha2, *Neoplasia* 6 (2004) 15–22.
- [50] T. Marumoto, A. Tashiro, D. Friedmann-Morvinski, M. Scadeng, Y. Soda, F.H. Gage, I.M. Verma, Development of a novel mouse glioma model using lentiviral vectors, *Nature Medicine* 15 (2009) 110–116.
- [51] T. Muraguchi, S. Tanaka, D. Yamada, A. Tamase, M. Nakada, H. Nakamura, T. Hoshii, T. Ooshio, Y. Tadokoro, K. Naka, Y. Ino, T. Todo, J. Kuratsu, H. Saya, J. Hamada, A. Hirao, NKX2.2 suppresses self-renewal of glioma-initiating cells, *Cancer Research* 71 (2011) 1135–1145.
- [52] M. Nagane, H. Lin, W.K. Cavenee, H.J. Huang, Aberrant receptor signaling in human malignant gliomas: mechanisms and therapeutic implications, *Cancer Letters* 162 (Suppl.) (2001) S17–S21.
- [53] J.R. Ohlfest, P.D. Lobitz, S.G. Perkinson, D.A. Largaespada, Integration and long-term expression in xenografted human glioblastoma cells using a plasmid-based transposon system, *Molecular Therapy* 10 (2004) 260–268.
- [54] H. Okada, K.L. Low, G. Kohanbash, H.A. McDonald, R.L. Hamilton, I.F. Pollack, Expression of glioma-associated antigens in pediatric brain stem and non-brain stem gliomas, *Journal of Neuro-Oncology* 88 (2008) 245–250.
- [55] T. Ozawa, J.L. Hu, L.J. Hu, E.L. Kong, A.W. Bollen, K.R. Lamborn, D.F. Deen, Functionality of hypoxia-induced BAX expression in a human glioblastoma xenograft model, *Cancer Gene Therapy* 12 (2005) 449–455.
- [56] W.B. Pope, Q. Xia, V.E. Paton, A. Das, J. Hambleton, H.J. Kim, J. Huo, M.S. Brown, J. Goldin, T. Cloughesy, Patterns of progression in patients with recurrent glioblastoma treated with bevacizumab, *Neurology* 76 (2011) 432–437.
- [57] M. Preusser, S. de Ribaupierre, A. Wohrer, S.C. Erridge, M. Hegi, M. Weller, R. Stupp, Current concepts and management of glioblastoma, *Annals of Neurology* 70 (2011) 9–21.
- [58] B. Purow, D. Schiff, Advances in the genetics of glioblastoma: are we reaching critical mass? *Nature Reviews Neurology* 5 (2009) 419–426.
- [59] N.G. Rainov, A phase III clinical evaluation of herpes simplex virus type 1 thymidine kinase and ganciclovir gene therapy as an adjuvant to surgical resection and radiation in adults with previously untreated glioblastoma multiforme, *Human Gene Therapy* 11 (2000) 2389–2401.
- [60] N.G. Rainov, C.M. Kramm, U. Banning, D. Riemann, H.J. Holzhausen, V. Heidecke, K.J. Burger, W. Burkert, D. Korholz, Immune response induced by retrovirus-mediated HSV-tk/GCV pharmacogene therapy in patients with glioblastoma multiforme, *Gene Therapy* 7 (2000) 1853–1858.
- [61] A.M. Sandmair, S. Loimas, P. Puranen, A. Immonen, M. Kossila, M. Puranen, H. Hurskainen, K. Tyynela, M. Turunen, R. Vanninen, P. Lehtolainen, L. Paljarvi, R. Johansson, M. Vapalahti, S. Yla-Herttuala, Thymidine kinase gene therapy for human malignant glioma, using replication-deficient retroviruses or adenoviruses, *Human Gene Therapy* 11 (2000) 2197–2205.
- [62] M. Sanson, J. Thillet, K. Hoang-Xuan, Molecular changes in gliomas, *Current Opinion in Oncology* 16 (2004) 607–613.
- [63] S. Sathornsumetee, J.N. Rich, Designer therapies for glioblastoma multiforme, *Annals of the New York Academy of Sciences* 1142 (2008) 108–132.
- [64] J.A. Schwartzbaum, J.L. Fisher, K.D. Aldape, M. Wrensch, Epidemiology and molecular pathology of glioma, *Nature Clinical Practice Neurology* 2 (2006) 494–503 (quiz p. 491 following 516).
- [65] L.M. Shelton, P. Mukherjee, L.C. Huysentruyt, I. Urits, J.A. Rosenberg, T.N. Seyfried, A novel pre-clinical in vivo mouse model for malignant brain tumor growth and invasion, *Journal of Neuro-Oncology* 99 (2010) 165–176.
- [66] A.H. Shih, C. Dai, X. Hu, M.K. Rosenblum, J.A. Koutcher, E.C. Holland, Dose-dependent effects of platelet-derived growth factor-B on glial tumorigenesis, *Cancer Research* 64 (2004) 4783–4789.
- [67] Y.A. Sidky, E.C. Borden, Inhibition of angiogenesis by interferons: effects on tumor- and lymphocyte-induced vascular responses, *Cancer Research* 47 (1987) 5155–5161.
- [68] P.S. Smitt, M. Driesse, J. Wolbers, M. Kros, C. Avezaat, Treatment of relapsed malignant glioma with an adenoviral vector containing the herpes simplex thymidine kinase gene followed by ganciclovir, *Molecular Therapy: The Journal of the American Society of Gene Therapy* 7 (2003) 851–858.
- [69] A.H. Stegh, C. Brennan, J.A. Mahoney, K.L. Forloney, H.T. Jeng, J.P. Luciano, A. Protopopov, L. Chin, R.A. Depinho, Glioma oncoprotein Bcl2L12 inhibits the p53 tumor suppressor, *Genes and Development* 24 (2010) 2194–2204.
- [70] R. Stupp, W.P. Mason, M.J. van den Bent, M. Weller, B. Fisher, M.J. Taphoorn, K. Belanger, A.A. Brandes, C. Marosi, U. Bogdahn, J. Curschmann, R.C. Janzer, S.K. Ludwin, T. Gorlia, A. Allgeier, D. Lacombe, J.G. Cairncross, E. Eisenhauer, R.O. Mirimanoff, Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma, *New England Journal of Medicine* 352 (2005) 987–996.
- [71] J.C. Tonn, M. Westphal, J.T. Rutka, *Neuro-Oncology of CNS Tumors*, 2005.
- [72] H. Tsurushima, X. Yuan, L.E. Dillehay, K.W. Leong, Radioresponsive tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) gene therapy for malignant brain tumors, *Cancer Gene Therapy* 14 (2007) 706–716.
- [73] H. Wajant, The role of TNF in cancer, *Results and Problems in Cell Differentiation* 49 (2009) 1–15.
- [74] P.Y. Wen, S. Kesari, Malignant gliomas in adults, *New England Journal of Medicine* 359 (2008) 492–507.
- [75] S.M. Wiesner, S.A. Decker, J.D. Larson, K. Ericson, C. Forster, J.L. Gallardo, C. Long, Z.L. Demorest, E.A. Zamora, W.C. Low, K. SantaCruz, D.A. Largaespada, J.R. Ohlfest, De novo induction of genetically engineered brain tumors in mice using plasmid DNA, *Cancer Research* 69 (2009) 431–439.
- [76] J. Wykosky, D.M. Gibo, C. Stanton, W. Debinski, Interleukin-13 receptor alpha 2, EphA2, and Fos-related antigen 1 as molecular denominators of high-grade astrocytomas and specific targets for combinatorial therapy, *Clinical Cancer Research* 14 (2008) 199–208.
- [77] Y. Zhu, L.F. Parada, The molecular and genetic basis of neurological tumours, *Nature Reviews Cancer* 2 (2002) 616–626.