

An Honest Approach to Parallel Trends*

Ashesh Rambachan[†] Jonathan Roth[‡]

September 10, 2020

Abstract

This paper proposes robust inference methods for difference-in-differences and event-study designs that do not require that the parallel trends assumption holds exactly. Instead, the researcher must only impose restrictions on possible violations of the parallel trends assumption, and our methodology then guarantees uniformly valid (“honest”) inference when the imposed restrictions are satisfied. We first show that fixed length confidence intervals have near-optimal expected length for a practically-relevant class of restrictions. We next introduce a novel inference procedure that accommodates a wider range of restrictions on the possible violations of parallel trends, and is based on the observation that inference in our setting is equivalent to testing a system of moment inequalities with a large number of linear nuisance parameters. The resulting confidence sets are consistent, and have optimal local asymptotic power for many parameter configurations. We recommend researchers conduct sensitivity analyses to show what conclusions can be drawn under various restrictions on the set of possible violations of parallel trends.

Keywords: Difference-in-differences, event-study, parallel trends, sensitivity analysis, robust inference, partial identification.

*We are grateful to Isaiah Andrews, Elie Tamer, and Larry Katz for their invaluable advice and encouragement. We also thank Gary Chamberlain, Raj Chetty, Peter Ganong, Ed Glaeser, Nathan Hendren, Ryan Hill, Ariella Kahn-Lang, Jens Ludwig, Sendhil Mullainathan, Claudia Noack, Frank Pinter, Adrienne Sabety, Pedro Sant’Anna, Jesse Shapiro, Neil Shephard, Jann Spiess, Jim Stock, and seminar participants at Brown, Chicago Booth, Dartmouth, Harvard, Michigan, Microsoft Research, and Princeton for helpful comments, and Dorian Carloni for sharing data. We gratefully acknowledge financial support from the NSF Graduate Research Fellowship under Grant DGE1745303 (Rambachan) and Grant DGE1144152 (Roth).

[†]Harvard University, Department of Economics. Email: asheshr@g.harvard.edu

[‡]Microsoft. Email: Jonathan.Roth@microsoft.com

1 Introduction

Conventional methods for causal inference in difference-in-differences and event-study designs are valid only under the so-called “parallel trends” assumption. However, researchers are often unsure whether the parallel trends assumption holds in applied settings.

This paper develops robust inference methods for difference-in-differences and event-study designs.¹ Instead of imposing that parallel trends holds exactly, our methods only require the researcher to specify that the possible violations of parallel trends are restricted to some known set Δ . The set Δ encodes context-specific knowledge about the possible violations of parallel trends, and the researcher can conduct sensitivity analyses to examine the robustness of their conclusions to different assumptions about such violations. For instance, we show that the intuition underlying the common practice of testing for pre-existing trends can be formalized by setting Δ to impose restrictions on the smoothness of the difference in trends between the pre-treatment and post-treatment periods. Likewise, economic knowledge about simultaneous policy changes or confounding secular trends may imply sign or shape restrictions. These types of restrictions, as well as many others, can be represented by polyhedral Δ 's (i.e., sets defined by a system of linear inequalities), which are the primary focus of our analysis. Under these polyhedral restrictions, the treatment effect of interest is typically set-identified.

To conduct inference on the treatment effect of interest, we first consider optimal fixed length confidence intervals (FLCIs) based on affine estimators, following [Donoho \(1994\)](#). FLCIs have attractive properties for certain types of restrictions – specifically, when Δ is convex and centrosymmetric – which include our baseline smoothness class. For these types of restrictions, results from [Armstrong and Kolesar \(2018, 2020\)](#) imply that the optimal FLCI has near-optimal expected length when in fact parallel trends holds. Unfortunately, however, we show that FLCIs can have poor properties for broader classes of restrictions, such as those that incorporate sign or shape restrictions. We provide a novel characterization of when the optimal FLCI is consistent, meaning that any fixed point outside of the identified set falls outside

¹Throughout the paper, we use the phrase “event-study” to refer to a large class of specifications that estimate dynamic treatment effects along with placebo pre-treatment effects. This includes, but is not limited to, settings with staggered treatment timing; see Related Literature and Section 7 below.

the FLCI with probability approaching one asymptotically. The optimal FLCI is consistent *if and only if* the length of the identified set is its maximum over the parameter space. This condition often fails for several leading restrictions on the class of differential trends, such as those that incorporate sign or shape restrictions. Our (in)consistency result may be of general interest in other applications where FLCIs have been used.

Motivated by this result, we introduce a general inference procedure that delivers desirable asymptotic properties over a larger class of possible restrictions on the class of differential trends. We exploit the novel observation that conducting inference on the treatment effect of interest under such restrictions is equivalent to a moment inequality problem with linear nuisance parameters. A practical challenge is that the number of nuisance parameters scales linearly in the number of post-treatment periods, and thus will often be large (above 10) in typical empirical applications. This renders many moment inequality procedures, which rely on test inversion over a grid for the full parameter vector, computationally infeasible. We overcome this computational challenge by employing the conditional inference procedure developed in [Andrews, Roth and Pakes \(2019, henceforth ARP\)](#), which delivers computationally tractable confidence sets that uniformly control size in our setting.

We then derive two novel results on the asymptotic performance of conditional confidence sets in our setting. First, we show that the conditional confidence sets are consistent for all polyderal Δ . Second, we provide a novel condition under which the conditional confidence sets have local asymptotic power converging to the power envelope (i.e., the upper bound on the power of any procedure that controls size uniformly).² To prove the optimality result, we make use of duality results from linear programming and the Neyman-Pearson lemma to show that both the optimal test and our conditional test converge to a t-test in the direction of the Lagrange multipliers of the linear program that profiles out the nuisance parameters. Both of these asymptotic results are novel, and exploit additional structure in our context not present in the more general setting considered in ARP.

Our results on the local asymptotic power of the conditional confidence sets have

²This condition is implied by the linear independence constraint qualification (LICQ), which has been used recently in the moment inequality settings studied in [Gafarov \(2019\)](#); [Cho and Russell \(2019\)](#); [Flynn \(2019\)](#); [Kaido and Santos \(2014\)](#). Moreover, in contrast to many previous uses of LICQ, we do not require this condition to obtain uniform asymptotic size control. See [Section 4.3](#) for further discussion.

two limitations. First, the condition needed for the conditional confidence sets to have optimal local asymptotic power does not hold for all parameter values, and in particular fails when the parameter of interest is point-identified. Second, our optimality results are under asymptotics where sampling variation grows small relative to the length of the identified set. Our asymptotic power results thus may not translate to good finite-sample power when sampling variation is large relative to the length of the identified set, particularly when there are non-binding moment restrictions that are close to binding.

Therefore, we finally introduce a novel hybrid inference procedure that combines the relative strengths of the conditional confidence sets and FLCIs. The hybrid procedure is consistent, and has near-optimal local asymptotic power when the condition for the optimality of the conditional approach holds. Further, we find in simulations (discussed in more detail below) that hybridization with the FLCIs improves performance in finite sample when the moments are not well-separated. In practice, we recommend the hybrid approach for settings in which the conditions for the consistency of the FLCIs is not guaranteed.

The main text of the paper presents results in the context of a finite-sample normal model, which facilitates exposition and provides a good approximation to the finite-sample distribution of conventional estimators in many settings. In the context of this model, we consider “small- σ asymptotics”, analogous to those in [Kadane \(1971\)](#) and [Moreira and Ridder \(2019\)](#), in which the sampling uncertainty grows small relative to the length of the identified set. In the supplemental materials, we show that these results in the normal model translate to asymptotic results that hold uniformly over a large class of non-normal data-generating processes satisfying some high-level regularity conditions.

To explore the performance of our methods in practice, we conduct simulations calibrated to the 12 recently-published empirical papers surveyed in [Roth \(2019\)](#). As expected based on the theoretical results, the performance of the FLCIs depends greatly on the choice of Δ . The FLCIs perform within 15 percent of the optimal bound for our baseline smoothness class, where the conditions for their finite-sample near-optimality hold. However, excess length for FLCIs can be substantially larger than for our other procedures for simulation designs that incorporate sign or shape restrictions. The conditional confidence sets perform well across a wider range of simulation designs, and often have excess length within a few percent of the optimum.

However, we find they can exhibit poor performance in settings where the binding and non-binding moments are not well-separated relative to the sampling variation in the data, such as when the target parameter is (nearly) point identified. Finally, the hybrid approach performs quite well across a wide range of specifications, with performance typically close to the better of the FLCIs and conditional approach. Based on our Monte Carlo simulations, we recommend the FLCIs for the special settings where the conditions for their consistency and finite-sample near-optimality hold, and otherwise recommend the hybrid approach.

We recommend applied researchers use our methods to conduct sensitivity analyses in which they report confidence sets under varying restrictions on the set of possible violations of parallel trends. For instance, one class of restrictions we consider restricts the extent to which the difference in trends may deviate from linearity, and is governed by a single parameter that determines the degree of possible non-linearity. If the researcher is interested in testing a particular null hypothesis — e.g., the treatment effect in a particular period is zero — then a simple statistic to report is the “breakdown” value of the non-linearity parameter at which the null hypothesis of interest can no longer be rejected.³ The researcher can also report how her conclusions change with the inclusion of additional sign or monotonicity restrictions motivated by context-specific knowledge. Performing such sensitivity analyses makes clear what must be assumed about the possible violations of parallel trends in order to draw specific causal conclusions. We provide an R package, `HonestDiD`, for implementation of our recommended methods.⁴ We conclude by applying our empirical recommendations to two recently published papers.

Related literature: This paper contributes to an active literature on difference-in-differences and event-study designs by developing robust inference methods that allow for possible violations of the parallel trends assumption. Our approach is most closely related to [Manski and Pepper \(2018\)](#), who show that the treatment effect of interest is partially identified under “bounded variation” assumptions that relax the usual exact parallel trends assumption. We consider identification under a broad class of restrictions on the possible violations of parallel trends, which nests as a special case the bounded variation assumptions considered in [Manski and Pepper \(2018\)](#).

³Similar “breakdown” concepts have been proposed in other settings with partial identification ([Horowitz and Manski, 1995](#); [Kline and Santos, 2013](#); [Masten and Poirier, 2020](#)).

⁴The latest version of may be downloaded [here](#).

This broader class of restrictions can be used to formalize a variety of arguments made intuitively in applied work — e.g., that pre-treatment differences in trends are informative about counterfactual post-treatment differences in trends. Importantly, we develop methods for conducting inference on the causal effects of treatment under these assumptions, whereas [Manski and Pepper \(2018\)](#) only consider identification.

Several other recent papers consider various relaxations of the parallel trends assumption. [Keele, Small, Hsu and Fogarty \(2019\)](#) develop techniques for testing the sensitivity of difference-in-differences designs to violations of the parallel trends assumption, but they do not incorporate information from the observed pre-trends in their sensitivity analysis. Empirical researchers commonly adjust for the extrapolation of a linear trend from the pre-treatment periods when there are concerns about violations of the parallel trends assumption, which is valid if the underlying trend is exactly linear (e.g., [Dobkin, Finkelstein, Kluender and Notowidigdo, 2018](#); [Goodman-Bacon, 2018a,b](#); [Bhuller, Havnes, Leuven and Mogstad, 2013](#)). Our methods nest this approach as a special case, but allow for valid inference under less restrictive assumptions about the class of possible differences in trends. [Freyaldenhoven, Hansen and Shapiro \(2019\)](#) propose a method that allows for violations of the parallel trends assumption but requires an additional covariate that is affected by the same confounding factors as the outcome but not by the treatment of interest. [Ye, Keele, Hasegawa and Small \(2020\)](#) consider partial identification of treatment effects when there exist two control groups whose outcomes have a bracketing relationship with the outcome of the treated group.

Our methods address several concerns related to established empirical practice in difference-in-differences and event-study designs. First, common tests for pre-trends may be underpowered against meaningful violations of parallel trends, potentially leading to severe undercoverage of conventional confidence intervals ([Freyaldenhoven et al., 2019](#); [Bilinski and Hatfield, 2018](#); [Kahn-Lang and Lang, 2019](#); [Roth, 2019](#)). Second, statistical distortions from pre-testing for pre-trends may further undermine the performance of conventional inference procedures ([Roth, 2019](#)). Third, parametric approaches to controlling for pre-existing trends may be quite sensitive to functional form assumptions ([Wolfers, 2006](#); [Lee and Solon, 2011](#)). We address these issues by providing tools for inference that do not rely on an exact parallel trends assumption and that make clear the mapping between assumptions on the potential differences in trends and the strength of one’s conclusions.

Our work is complementary to a growing literature on the causal interpretation of event-study coefficients in two-way fixed effects models in the presence of staggered treatment timing or heterogeneous treatment effects (Meer and West, 2016; Borusyak and Jaravel, 2016; Sun and Abraham, 2020; Athey and Imbens, 2018; de Chaisemartin and D’Haultfoeuille, 2018a,b; Goodman-Bacon, 2018a; Kropko and Kubinec, 2018; Callaway and Sant’Anna, 2019; Imai and Kim, 2019; Słoczyński, 2018). A key finding is that regression coefficients from conventional approaches may not produce convex weighted averages of treatment effects even if parallel trends holds. Several alternative estimators have been proposed that consistently estimate sensible causal estimands under a suitable parallel trends assumption. Our methodology can be applied to assess the sensitivity of results obtained using these methods to violations of the corresponding parallel trends assumption; see Section 7 for additional discussion.

More broadly, our approach relates to a large and active literature on sensitivity analysis and misspecification robust inference, including Imbens (2003); Rosenbaum (2005); Altonji, Elder and Taber (2005); Conley, Hansen and Rossi (2012); Kolesar and Rothe (2018); Armstrong and Kolesar (2018); Masten and Poirier (2018); Bonhomme and Weidner (2018); Oster (2019) among many others.

2 General set-up

We introduce the assumptions, target parameter, and inferential goal considered in the paper. We consider a finite-sample normal model, which arises as an asymptotic approximation to a variety of econometric settings of interest.

2.1 Finite sample normal model

Consider the following model

$$\hat{\beta}_n \sim \mathcal{N}(\beta, \Sigma_n), \tag{1}$$

where $\hat{\beta}_n \in \mathbb{R}^{T+\bar{T}}$ and $\Sigma_n = \frac{1}{n}\Sigma^*$ for Σ^* a known, positive-definite $(T + \bar{T}) \times (T + \bar{T})$ matrix. We refer to $\hat{\beta}_n$ as the estimated event-study coefficients, and partition $\hat{\beta}_n$ into vectors corresponding with the pre-treatment and post-treatment periods, $\hat{\beta}_n = (\hat{\beta}'_{n,pre}, \hat{\beta}'_{n,post})'$, where $\hat{\beta}_{n,pre} \in \mathbb{R}^T$ and $\hat{\beta}_{n,post} \in \mathbb{R}^{\bar{T}}$. We adopt analogous notation to partition other vectors that are the same length as $\hat{\beta}_n$.

This finite sample normal model (1) can be viewed as an asymptotic approximation to a wide range of econometric settings. Under mild regularity conditions, a variety of estimation strategies in difference-in-differences and event study designs will yield asymptotically normal estimated event-study coefficients, $\sqrt{n}(\hat{\beta}_n - \beta) \xrightarrow{d} \mathcal{N}(0, \Sigma^*)$.⁵ This convergence in distribution suggests the finite-sample approximation $\hat{\beta}_n \stackrel{d}{\approx} \mathcal{N}(\beta, \Sigma_n)$, where $\stackrel{d}{\approx}$ denotes approximate equality in distribution and $\Sigma_n = \frac{1}{n}\Sigma^*$. We derive results assuming this equality in distribution holds exactly in finite samples. In the supplemental materials, we show that results in the finite sample normal model translate to uniform asymptotic statements for a large class of data-generating processes.

We assume the mean vector β satisfies the following causal decomposition.

Assumption 1. *The parameter vector β can be decomposed as*

$$\beta = \underbrace{\begin{pmatrix} \tau_{pre} \\ \tau_{post} \end{pmatrix}}_{:= \tau} + \underbrace{\begin{pmatrix} \delta_{pre} \\ \delta_{post} \end{pmatrix}}_{:= \delta} \text{ with } \tau_{pre} \equiv 0. \quad (2)$$

The first term, τ , represents the time path of the dynamic causal effects of interest. We assume the treatment has no causal effect prior to its implementation, so $\tau_{pre} = 0$. The second term, δ , represents the difference in trends between the treated and untreated groups that would have occurred absent treatment. The parallel trends assumption imposes that $\delta = 0$. Therefore, under parallel trends, $\beta = \tau$.

Example: Difference-in-differences We observe an outcome Y_{it} for a sample of individuals $i = 1, \dots, N$ for three time periods, $t = -1, 0, 1$. Individuals in the treated population ($D_i = 1$) receive a treatment between period $t = 0$ and $t = 1$.⁶ The observed outcome equals $Y_{i,t} = D_i Y_{i,t}(1) + (1 - D_i) Y_{i,t}(0)$, where $Y_{i,t}(1)$ and $Y_{i,t}(0)$

⁵Examples of estimators that yield asymptotically normal event-study estimates include canonical two-way fixed effects estimators, the GMM procedure proposed by Freyaldenhoven et al. (2019), instrumental variables event-studies (Hudson, Hull and Liebersohn, 2017), the estimation strategies of Sun and Abraham (2020) and Callaway and Sant’Anna (2019) to address issues with non-convex weights on cohort-specific effects in staggered treatment designs, as well as a range of procedures that flexibly control for differences in covariates between treated and untreated groups (e.g., Heckman, Ichimura, Smith and Todd, 1998; Abadie, 2005; Sant’Anna and Zhao, 2020).

⁶For the purposes of this example, we think of the observed sample as consisting of N_1 independent draws from the treated ($D = 1$) population and N_0 independent draws from the control population ($D = 0$) with $N = N_0 + N_1$, as in Abadie and Imbens (2006).

are the potential outcomes for individual i in period t associated with the treatment and control conditions. Assume the treatment has no causal effect prior to implementation, meaning $Y_{i,t}(1) = Y_{i,t}(0)$ for $t < 1$. The causal estimand of interest is the average treatment effect on the treated (ATT), $\tau_{ATT} = \mathbb{E}[Y_{i,1}(1) - Y_{i,1}(0) | D_i = 1]$.

In this setting, researchers commonly estimate the “dynamic event study regression”

$$Y_{it} = \lambda_i + \phi_t + \sum_{s \neq 0} \beta_s \times \mathbb{1}[t = s] \times D_i + \epsilon_{it}. \quad (3)$$

The estimated coefficient $\hat{\beta}_1$ equals the “difference-in-differences” of sample means across treated and untreated groups between period $t = 0$ and $t = 1$, $\hat{\beta}_1 = (\bar{Y}_{1,1} - \bar{Y}_{1,0}) - (\bar{Y}_{0,1} - \bar{Y}_{0,0})$, where $\bar{Y}_{d,t}$ is the sample mean of Y_{it} for treatment group d . The “pre-period” coefficient $\hat{\beta}_{-1}$ can be written as $\hat{\beta}_{-1} = (\bar{Y}_{1,-1} - \bar{Y}_{1,0}) - (\bar{Y}_{0,-1} - \bar{Y}_{0,0})$.

Taking expectations and re-arranging, we see that

$$\begin{aligned} \mathbb{E}[\hat{\beta}_1] &= \tau_{ATT} + \underbrace{\mathbb{E}[Y_{i,1}(0) - Y_{i,0}(0) | D_i = 1] - \mathbb{E}[Y_{i,1}(0) - Y_{i,0}(0) | D_i = 0]}_{\text{Post-period differential trend} := \delta_1}, \\ \mathbb{E}[\hat{\beta}_{-1}] &= \underbrace{\mathbb{E}[Y_{i,-1}(0) - Y_{i,0}(0) | D_i = 1] - \mathbb{E}[Y_{i,-1}(0) - Y_{i,0}(0) | D_i = 0]}_{\text{Pre-period differential trend} := \delta_{-1}}. \end{aligned}$$

The parameter $\beta = \mathbb{E}[\hat{\beta}]$ thus satisfies the decomposition (2), where $\tau_{post} = \tau_{ATT}$ is the ATT, $\delta_{post} = \delta_1$ is the difference in trends in *untreated* potential outcomes between $t = 0$ and $t = 1$, and $\delta_{pre} = \delta_{-1}$ is the analogous difference in untreated potential outcomes between $t = -1$ and $t = 0$. Under suitable regularity conditions, $\hat{\beta}$ will also satisfy a central limit theorem, so that (1) will hold approximately in large samples.

▲

Remark 1. We have motivated our normal model from a sampling-based perspective, which is the most common framework for uncertainty in the difference-in-differences literature. While applicable in many cases, the sampling view may be unnatural in some settings, such as when the unit of observation is a state and all 50 states are observed (Manski and Pepper, 2018). In Rambachan and Roth (2020), we show that the normal model (1) also arises from a design-based model that treats the population as fixed and views the assignment of treatment as the source of randomness in the data. In that setting, δ is a function of the finite-population covariance between treatment probabilities and potential outcomes.

2.2 Target parameter and identification

The parameter of interest is a scalar, linear combination of the post-treatment causal effects, $\theta := l' \tau_{post}$ for some known \bar{T} -vector l . For example, θ equals the t -th period causal effect τ_t when the vector l equals the t -th standard basis vector. Similarly, θ equals the average causal effect across all post-treatment periods when $l = (\frac{1}{T}, \dots, \frac{1}{T})'$. Under the parallel trends assumption ($\delta = 0$), θ is point-identified.

We relax the parallel trends assumption by assuming that δ lies in a set of possible violations of parallel trends Δ , which is specified by the researcher. This nests the usual parallel trends assumption as a special case with $\Delta = \{0\}$. Under the assumption that $\delta \in \Delta \neq \{0\}$, the parameter θ will typically be set-identified. For a given value of β , the set of values θ consistent with β under the assumption $\delta \in \Delta$ is

$$\mathcal{S}(\beta, \Delta) := \left\{ \theta : \exists \delta \in \Delta, \tau_{post} \in \mathbb{R}^{\bar{T}} \text{ s.t. } l' \tau_{post} = \theta, \beta = \delta + \begin{pmatrix} 0 \\ \tau_{post} \end{pmatrix} \right\}, \quad (4)$$

which we refer to as the *identified set*. When Δ is a closed and convex set, the identified set has a simple characterization.

Lemma 2.1. *If Δ is closed and convex, then $\mathcal{S}(\beta, \Delta)$ is an interval in \mathbb{R} , $\mathcal{S}(\beta, \Delta) = [\theta^{lb}(\beta, \Delta), \theta^{ub}(\beta, \Delta)]$, where*

$$\theta^{lb}(\beta, \Delta) := l' \beta_{post} - \underbrace{\left(\max_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = \beta_{pre} \right)}_{=: b^{max}(\beta_{pre}; \Delta)}, \quad (5)$$

$$\theta^{ub}(\beta, \Delta) := l' \beta_{post} - \underbrace{\left(\min_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = \beta_{pre} \right)}_{=: b^{min}(\beta_{pre}; \Delta)}. \quad (6)$$

Proof. Re-arranging terms in (4), the identified set can be equivalently written as $\mathcal{S}(\beta, \Delta) = \{\theta : \exists \delta \in \Delta \text{ s.t. } \delta_{pre} = \beta_{pre}, \theta = l' \beta_{post} - l' \delta_{post}\}$. The result is then immediate. \square

2.3 Possible choices of Δ

The class of possible violations of parallel trends Δ must be specified by the researcher, and the choice of Δ will depend on the economic context. We highlight several choices of Δ that may be reasonable in empirical applications and formalize

intuitive arguments that are commonly made by applied researchers regarding possible violations of parallel trends. We ultimately recommend that researchers conduct sensitivity analysis with respect to a range of Δ 's that may be plausible in their context.

2.3.1 Smoothness restrictions

We begin by introducing a class of restrictions that formalizes the intuition behind the common practice of testing for pre-existing differences in trends (pre-trends). Researchers frequently test for pre-trends as a way of assessing the plausibility of the parallel trends assumption. The logic of these tests is intuitively based on the notion that the pre-trend is informative about the counterfactual difference in trends. In other words, the difference in trends must evolve “smoothly” over time; if not, then the fact that the pre-trend is close to zero would not be informative about the validity of the parallel trends assumption, since the (counterfactual) difference in trends could be close to zero in the pre-treatment period and then jump around the time of treatment.


We formalize this logic by introducing smoothness restrictions on the possible differences in trends. Specifically, we bound the extent to which the difference in trends can deviate from linearity. Deviations from linearity are a natural starting point since applied researchers concerned about possible violations of the parallel trends assumption commonly include treatment-group specific linear trends in their regression specifications.⁷ There are concerns, however, that this linear extrapolation of the pre-trend may not quite be correct (Wolfers, 2006; Lee and Solon, 2011). A natural relaxation is therefore to require only that the difference in trends not deviate “too much” from linearity. We formalize this by bounding the extent to which the slope of the differential trend may change between consecutive periods, requiring that δ lie in the set

$$\Delta^{SD}(M) := \{\delta : |(\delta_{t+1} - \delta_t) - (\delta_t - \delta_{t-1})| \leq M, \forall t\}, \quad (7)$$

⁷That is, researchers may augment specification (3) with group-specific linear trends, an approach Dobkin et al. (2018) refer to as a “parametric event-study.” An analogous approach is to estimate a linear trend using only observations prior to treatment, and then subtract out the estimated linear trend from the observations after treatment (Bhuller et al., 2013; Goodman-Bacon, 2018a,b).

where we adopt the convention that $\delta_0 = 0$.⁸ The parameter $M \geq 0$ governs the amount by which the slope of δ can change between consecutive periods. In the special case where $M = 0$, $\Delta^{SD}(0)$ requires that the difference in trends be exactly linear.

Figure 1: Intuition for $\Delta^{SD}(M)$




Alternatively, one might allow the smoothness of the differential trend to depend on the magnitude of the pre-trend. For instance, applied researchers may have intuition that if the observed pre-period difference in trends is small, then the counterfactual post-period difference in trends would also be small. If the two groups did not follow similar trends in the pre-period, though, it may be more plausible that the difference in trends between the two groups would have changed substantially between the pre- and post-periods. This intuition may be formalized by bounding the percentage change in the slope of the differential trend across periods,

$$\Delta^{RM}(\bar{M}) := \{\delta : |\delta_{t+1} - \delta_t| \leq \bar{M}|\delta_t - \delta_{t-1}|, \forall t\}. \quad (8)$$

Example: Difference-in-differences (continued) In the three-period difference-in-differences model, assuming the differential trend is exactly linear is equivalent to assuming $\Delta = \{\delta : \delta_1 = -\delta_{-1}\}$. In contrast, assuming $\delta \in \Delta^{SD}(M)$ only requires that the linear extrapolation be *approximately* correct, $\delta_1 \in [-\delta_{-1} - M, -\delta_{-1} + M]$. $\Delta^{RM}(\bar{M})$ bounds the magnitude of δ_1 based on the magnitude of δ_{-1} with $\Delta^{RM}(\bar{M}) = \{(\delta_{-1}, \delta_1) : |\delta_1| \leq \bar{M}|\delta_{-1}|\}$. The larger the magnitude of the observed pre-period violation in parallel trends, $|\delta_{-1}|$, the wider the range of possible post-period violations of parallel trends. Figure 2 gives a geometric depiction of Δ^{SD} and Δ^{RM} in this example. ▲

⁸ $\Delta^{SD}(M)$ bounds the discrete analog of the second derivative of δ , and is thus similar to restrictions on the second derivative of the conditional expectation function or density in regression discontinuity settings (Kolesar and Rothe, 2018; Frandsen, 2016; Noack and Rothe, 2019). Smoothness restrictions are also used to obtain partial identification in Kim, Kwon, Kwon and Lee (2018).

Figure 2: Example choices for Δ



Note: Diagrams of potential restrictions Δ on the set of possible violations of parallel trends in the three-period difference-in-differences model. See discussion in Section 2 for further details on each example.

2.3.2 Sign and monotonicity restrictions

Context-specific knowledge may sometimes further imply sign or monotonicity restrictions on the differential trend. For instance, there may be simultaneous, confounding policy changes that we expect to have a positive effect on the outcome of interest, in which case we might restrict the post-period bias to be positive, $\delta \in \Delta^{PB} := \{\delta : \delta_t \geq 0 \forall t \geq 0\}$. Likewise, in some empirical settings, there may be secular trends that we expect would have continued following the treatment date.⁹ We may then wish to impose that the differential trend be increasing, $\delta \in \Delta^I := \{\delta : \delta_t \geq \delta_{t-1} \forall t\}$. Such sign and monotonicity restrictions may also be combined with smoothness restrictions. For example, define $\Delta^{SDPB}(M) := \Delta^{SD}(M) \cap \Delta^{PB}$ and $\Delta^{RMI}(\bar{M}) := \Delta^{RM}(\bar{M}) \cap \Delta^I$. Figure 2 gives a geometric depiction of Δ^{SDPB} and Δ^{RMI} in the three-period difference-in-differences model.

2.3.3 Polyhedral restrictions

The smoothness, shape, and sign restrictions discussed so far will be applicable in a variety of economic contexts. However, in some cases researchers may have context-specific knowledge that imply other types of restrictions. Throughout the paper, we therefore consider the broader class of Δ 's that take a polyhedral form, i.e. sets which

⁹Monotone violations of parallel trends are often discussed in applied work. For example, [Lovenheim and Willen \(2019\)](#) argue that violations of parallel trends cannot explain their results because “pre-[treatment] trends are either zero or in the wrong direction (i.e., opposite to the direction of the treatment effect).” [Greenstone and Hanna \(2014\)](#) estimate upward-sloping pre-existing trends and argue that “if the pre-trends had continued” their estimates would be upward biased.

can be expressed as a series of linear restrictions on δ .

Assumption 2 (Polyhedral shape restriction). *The class Δ takes the form $\Delta = \{\delta : A\delta \leq d\}$ for some known matrix A and vector d , where the matrix A has no all-zero rows.*

This class of restrictions encompasses nearly all of the aforementioned examples, as well as many others. Indeed, it is immediate from Figure 2 that with two dimensions, Δ^{SD} , Δ^{SDPB} , and Δ^{RMI} are all polyhedra, and the geometric intuition from the two-period case extends to higher dimensions.¹⁰ The one exception is Δ^{RM} , which is not convex and thus not a polyhedron. However, Δ^{RM} can be expressed as the union of polyhedra. One can thus form a confidence set for Δ^{RM} by taking the union of the confidence sets we develop below for each of the polyhedra that compose Δ^{RM} .

Remark 2 (Bounded variation assumptions). Manski and Pepper (2018) consider identification of treatment effects under so-called “bounded variation assumptions.” These assumptions can be expressed in the polyhedral form introduced in Assumption 2. Within the context of our ongoing difference-in-differences example, MP’s “bounded difference-in-differences variation” assumption corresponds directly with placing a bound on the magnitude of $|\delta_1|$ when $\hat{\beta}_1$ is the coefficient from specification (3). MP also consider “bounded time” and “bounded state” variation assumptions, which correspond with bounds on the magnitudes of $|\mu_{11} - \mu_{10}|$ and $|\mu_{11} - \mu_{01}|$, where $\mu_{dt} := \mathbb{E}[Y(0)|D = d, t = t]$. These restrictions can be accommodated by augmenting the vector $\hat{\beta}$ to include the sample means corresponding with estimates of the differences in outcomes for the appropriate treatment-group by time period cells.¹¹ ■

Remark 3 (Ashenfelter’s dip). When applying difference-in-differences to job training programs, one might worry that people who enroll in the program choose to do so in response to negative transitive shocks to earnings. If so, then the counterfactual difference in trends will exhibit the so-called Ashenfelter’s dip (Ashenfelter, 1978), in

¹⁰In the case with one pre-period and one post-period, $\Delta^{SD}(M) = \{\delta : A^{SD}\delta \leq d^{SD}\}$ for $A^{SD} = \begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix}$ and $d^{SD} = \begin{pmatrix} M \\ M \end{pmatrix}$. This generalizes naturally when there are multiple pre-periods and multiple post-periods.

¹¹After augmenting the vector for the event-study coefficients, equation (2) needs to be re-written to replace $(0, \tau'_{post})'$ with $M\tau_{post}$, where M is a matrix that accounts for the fact that elements of τ enter both the event-study coefficients and the augmented terms. Our proposed methods and results do not rely on the structure that $M = (0, I)'$ and thus easily accommodate this modification.

which earnings groups for the treated group trend downwards (relative to control) before treatment and upwards afterwards. In this type of setting, a researchers might naturally use a polyhedral Δ to impose i) restrictions on the signs of the pre-treatment and post-treatment biases, as well as ii) restrictions on the magnitude of the rebound effect relative to the pre-treatment shock.

2.4 Inferential Goal

Given a particular choice of Δ , our goal is to construct confidence sets that are uniformly valid for all parameter values θ in the identified set. We construct confidence sets \mathcal{C}_n satisfying

$$\inf_{\delta \in \Delta, \tau} \inf_{\theta \in \mathcal{S}(\Delta, \delta + \tau)} \mathbb{P}_{(\delta, \tau, \Sigma_n)}(\theta \in \mathcal{C}_n) \geq 1 - \alpha. \quad (9)$$

We subscript the probability operator by (δ, τ, Σ_n) to make explicit that the distribution of $\hat{\beta}_n$ (and hence \mathcal{C}_n) depends on these parameters. In the supplemental materials, we show that the coverage requirement (9) in the normal model translates to uniform asymptotic coverage over a large class of data-generating processes.

In practice, we recommend that applied researchers tailor their choice of confidence set \mathcal{C}_n based on their choice of Δ . If the researcher selects $\Delta^{SD}(M)$ or related Δ satisfying particular properties described below, we recommend that the researcher use the optimal fixed length confidence interval (FLCI) described in Section 3. Otherwise, we recommend that the researcher use the conditional-FLCI hybrid confidence set, which is introduced in Section 5. Our practical recommendations are based on the theoretical properties of these confidence sets that we derive in Sections 3-5 and the simulation results in Section 6. An applied reader interested in applying our methods but not their theoretical properties may wish to skip ahead to Sections 7-8, in which we provide further details on our practical recommendations and illustrate them in two applications to recently published empirical papers.

3 Inference using Fixed Length Confidence Intervals

We first consider fixed length confidence intervals (FLCIs) based on affine estimators. We show that FLCIs deliver finite-sample guarantees for certain choices of Δ , including our baselines smoothness class Δ^{SD} , but may perform poorly for other types of restrictions.

3.1 Constructing FLCIs

Following [Donoho \(1994\)](#) and [Armstrong and Kolesar \(2018, 2020\)](#), we consider fixed length confidence intervals based on an affine estimator for θ ,

$$\mathcal{C}_{\alpha,n}(a, v, \chi) = \left(a + v' \hat{\beta}_n \right) \pm \chi, \quad (10)$$

where a and χ are scalars and $v \in \mathbb{R}^{T+\bar{T}}$. We wish to minimize the half-length of the confidence interval χ subject to the constraint that $\mathcal{C}_{\alpha,n}(a, v, \chi)$ satisfies the coverage requirement (9).

To do so, note that $a + v' \hat{\beta}_n \sim \mathcal{N}(a + v' \beta, v' \Sigma_n v)$, and hence $|a + v' \hat{\beta}_n - \theta| \sim |\mathcal{N}(b, v' \Sigma_n v)|$, where $b = a + v' \beta - \theta$ is the bias of the affine estimator $a + v' \hat{\beta}$ for θ . Observe further that $\theta \in \mathcal{C}_n(a, v, \chi)$ if and only if $|a + v' \hat{\beta}_n - \theta| \leq \chi$. For fixed values a and v , the smallest value of χ that satisfies (9) is therefore the $1 - \alpha$ quantile of the $|\mathcal{N}(\bar{b}, v' \Sigma_n v)|$ distribution, where \bar{b} is the worst-case bias of the affine estimator,

$$\bar{b}(a, v) := \sup_{\delta \in \Delta, \tau_{post} \in \mathbb{R}^{\bar{T}}} |a + v'(\delta + M_{post} \tau_{post}) - l' \tau_{post}|. \quad (11)$$

Let $cv_\alpha(t)$ denote the $1 - \alpha$ quantile of the folded normal distribution $|\mathcal{N}(t, 1)|$.¹² For fixed a and v , the smallest value of χ satisfying the coverage requirement (9) is

$$\chi_n(a, v; \alpha) = \sigma_{v,n} \cdot cv_\alpha(\bar{b}(a, v)/\sigma_{v,n}), \quad (12)$$

where $\sigma_{v,n} := \sqrt{v' \Sigma_n v}$.

The minimum-length FLCI is then constructed by choosing the values of a and v to minimize (12). This minimization optimally trades off bias and variance, since the half-length $\chi_n(a, v; \alpha)$ is increasing in both the worst-case bias \bar{b} and the variance $\sigma_{v,n}^2$ (assuming $\alpha \in (0, 0.5]$). When Δ is convex, this minimization can be solved as a nested optimization problem, where both the inner and outer minimizations are convex ([Low, 1995](#); [Armstrong and Kolesar, 2018, 2020](#)). We denote by $\mathcal{C}_{\alpha,n}^{FLCI}$ the $1 - \alpha$ level FLCI with the shortest length,

$$\mathcal{C}_{\alpha,n}^{FLCI} = \left(a_n + v_n' \hat{\beta}_n \right) \pm \chi_n, \quad (13)$$

¹²If $t = \infty$, we define $cv_\alpha = \infty$.

where $\chi_n := \inf_{a,v} \chi_n(a, v; \alpha)$ and a_n, v_n are the optimal values in the minimization.

Example: $\Delta^{SD}(M)$ Suppose $\theta = \tau_1$. For $\Delta^{SD}(M)$, the affine estimator used by the optimal FLCI takes the form

$$a + v' \hat{\beta}_n = \hat{\beta}_{n,1} - \sum_{s=-T+1}^0 w_s \left(\hat{\beta}_{n,s} - \hat{\beta}_{n,s-1} \right), \quad (14)$$

where the weights w_s sum to one (but may be negative). It takes the post-period event-study coefficient for period 1 and subtracts out a weighted sum of the estimated slopes between consecutive pre-periods. Intuitively, since Δ^{SD} restricts the changes in the slope of the underlying trend across periods, but not the slope of the trend itself, an affine estimator with finite bias must subtract out an estimate of the slope of the trend between $t = 0$ and $t = 1$ using the observed slopes in the pre-period. The worst-case bias will be smaller if more weight is placed on pre-periods closer to the treatment date, but it may reduce variance to place more weight on earlier pre-periods. The weights w_s are optimally chosen to balance this tradeoff. \blacktriangle

Example: $\Delta^{SDPB}(M)$ and $\Delta^{SDI}(M)$ Suppose $\theta = \tau_1$. The affine estimator used by the optimal FLCI for both $\Delta^{SDPB}(M)$ and $\Delta^{SDI}(M)$ is the same as that used for $\Delta^{SD}(M)$. This is because the worst-case bias for an affine estimator is the same across all three Δ 's. To see why, suppose that the vector δ maximizes the bias for an affine estimator (a, v) over $\Delta^{SD}(M)$. Notice that the vector that adds a constant slope to δ , say $\tilde{\delta}_c = \delta + c \cdot (-T, \dots, \bar{T})'$, also lies in $\Delta^{SD}(M)$. For c sufficiently large, $\tilde{\delta}_c$ will lie in $\Delta^{SDPB}(M)$ and $\Delta^{SDI}(M)$ as well. However, we argued in the previous example that if (a, v) produces finite worst-case bias over Δ^{SD} , it must subtract out a weighted average of the pre-treatment slopes, and the same argument applies for Δ^{SDPB} and Δ^{SDI} . It follows that the worst-case bias under δ is the same as the worst-case bias under $\tilde{\delta}_c$, as we wished to show. Therefore, the optimal FLCI is unable to adapt to additional sign or monotonicity restrictions. \blacktriangle

Example: $\Delta^{RMI}(\bar{M})$ Suppose $\theta = \tau_1$. If $\Delta = \Delta^{RMI}(\bar{M})$ and $\bar{M} > 0$, then all affine estimators for τ_1 have infinite worst-case bias, and the FLCI is thus the entire real line. To see this, observe that since τ_{post} is unrestricted in (11), any affine estimator with finite bias must set $v_{post} = e_1$. For $v_{post} = e_1$, the bias of the affine estimator

$a + v'\hat{\beta}_n$ equals $|a + v'_{pre}\delta_{pre} + \delta_1|$ regardless of the value of τ . By the triangle inequality, $\max\{|a + v'_{pre}\delta_{pre} + \tilde{\delta}_1|, |a + v'_{pre}\delta_{pre} + \bar{\delta}_1|\} \geq \frac{1}{2}|\tilde{\delta}_1 - \bar{\delta}_1|$ for any two values $\tilde{\delta}_1$ and $\bar{\delta}_1$. As seen in Figure 2 for the two-period case, the range of values of δ_1 that are feasible given a pre-period trend δ_{pre} can be made arbitrarily large by setting δ_{pre} such that $|\delta_{-1}|$ is large. Hence $\bar{b}(a, v) = \infty$ for all choices of (a, v) . \blacktriangle

3.2 Finite-sample near optimality

In particular cases of interest, such as when $\Delta = \Delta^{SD}(M)$, the FLCIs introduced above have near-optimal expected length in the finite-sample normal model. The following result, which is an immediate consequence of results in [Armstrong and Kolesar \(2018, 2020\)](#), bounds the ratio of the expected length of the shortest possible confidence interval that controls size relative to the length of the optimal FLCI.

Assumption 3. Assume i) Δ is convex and centrosymmetric (i.e. $\delta \in \Delta$ implies $-\delta \in \Delta$), and ii) $\delta_A \in \Delta$ is such that $(\delta - \delta_A) \in \Delta$ for all $\delta \in \Delta$.

Proposition 3.1. Suppose δ_A and Δ satisfy Assumption 3. Then, for any $\tau_A \in \mathbb{R}^T$, Σ^* positive definite, and $n > 0$,

$$\frac{\inf_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_{\alpha}(\Delta, \Sigma_n)} \mathbb{E}_{(\delta_A, \tau_A, \Sigma_n)} [\lambda(\mathcal{C}_{\alpha, n})]}{2\chi_n} \geq \frac{z_{1-\alpha}(1-\alpha) - \tilde{z}_{\alpha}\Phi(\tilde{z}_{\alpha}) + \phi(z_{1-\alpha}) - \phi(\tilde{z}_{\alpha})}{z_{1-\alpha/2}},$$

where $\lambda(\cdot)$ denotes the length of an interval and $\tilde{z}_{\alpha} = z_{1-\alpha} - z_{1-\alpha/2}$.

Part i) of Assumption 3 is satisfied for Δ^{SD} but not for our other ongoing examples. For example, Δ^{SDPB} and Δ^{RMI} are convex but not centrosymmetric, and Δ^{RM} is neither convex nor centrosymmetric. Part ii) of Assumption 3 is always satisfied when parallel trends holds ($\delta_A = 0$). It also holds whenever δ_A is a linear trend for the case of $\Delta^{SD}(M)$. For $\alpha = 0.05$, the lower bound in Proposition 3.1 evaluates to 0.72, so the expected length of the shortest confidence possible interval that satisfies the coverage requirement (9) is at most 28% shorter than the length of the optimal FLCI.¹³ A nice feature of this result is that it places no restrictions on Σ , and thus

¹³Additionally, as noted in [Armstrong and Kolesar \(2020\)](#), the results in [Joshi \(1969\)](#) imply that under part i) of Assumption 3, the FLCI achieves minimax expected length, so any procedure that outperforms it somewhere must also be inferior at some other point in the parameter space.

allows the sampling variation in the data to be large (or small) relative to the length of the identified set.

3.3 (In)Consistency of FLCIs

The finite-sample guarantees discussed above do not apply for several types of restrictions Δ of importance, including those that incorporate sign and shape restrictions. We now show that the FLCIs can perform poorly under such restrictions. Specifically, we consider asymptotics wherein the sampling uncertainty grows small relative to the length of the identified set, and show that the FLCIs include points bounded away from the identified set with non-vanishing probability.

Recall from Lemma 2.1 that the identified set $\mathcal{S}(\beta, \Delta)$ is an interval when Δ is convex, with length equal to $\theta^{ub}(\beta, \Delta) - \theta^{lb}(\beta, \Delta) = b^{max}(\beta_{pre}, \Delta) - b^{min}(\beta_{pre}, \Delta)$. Since the length of the identified set only depends on Δ and β_{pre} , denote it by $LID(\beta_{pre}, \Delta)$. Our next result shows that $\mathcal{C}_{\alpha, n}^{FLCI}$ is consistent if and only if Assumption 4 holds, provided that the identified set is not the entire real line (in which case any procedure is trivially consistent).

Assumption 4 (Identified set maximal length and finite). *Suppose δ_{pre} is such that $LID(\delta_{pre}, \Delta) = \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\tilde{\delta}_{pre}, \Delta) < \infty$.*

Proposition 3.2. *Suppose Δ is convex and $\alpha \in (0, .5]$. Fix $\delta_A \in \Delta$ and $\tau_A \in \mathbb{R}^T$, and suppose $\mathcal{S}(\delta_A + \tau_A, \Delta) \neq \mathbb{R}$. Then (δ_A, Δ) satisfy Assumption 4 if and only if $\mathcal{C}_{\alpha, n}^{FLCI}$ is consistent, meaning that*


$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)}(\theta_{out} \in \mathcal{C}_{\alpha, n}^{FLCI}) = 0 \text{ for all } \theta_{out} \notin \mathcal{S}_\theta(\delta_A + \tau_A, \Delta).$$

Thus, if Assumption 4 fails, then $\mathcal{C}_{\alpha, n}^{FLCI}$ is inconsistent in the strong sense that it includes fixed points outside of the identified set with non-vanishing probability. Moreover, we show in Lemma B.26 in the Appendix that the conditions of Proposition 3.1 imply that Assumption 4 holds. Thus, the FLCIs obtain finite sample near-optimality in only a subset of the cases where they are consistent.

Remark 4. In the three-period difference-in-differences example, the length of the identified set corresponds with the height of Δ in Figure 3, and so Assumption 4 holds if and only if Δ achieves its maximal height at δ_{-1} . As shown in Figure 3, the

assumption holds everywhere for Δ^{SD} , for values of δ where the sign restrictions do not bind for Δ^{SDPB} , and nowhere for Δ^{RMI} . The restrictiveness of Assumption 4 thus depends greatly on Δ . ■

Figure 3: Diagram of where Assumptions 3 and 4 hold. The values of δ are colored red (neither holds), light green (Assumption 4 only), and dark green (both hold).



Remark 5. Proposition 3.2 implies that FLCIs can potentially be inconsistent when Δ is convex and centrosymmetric if $\delta \neq 0$. For example, if $\Delta = \{\delta \in \Delta^{SD}(M) \mid \delta_1 \leq M\}$, then the FLCI is inconsistent whenever $\delta_{-1} \neq 0$, even though Proposition 3.1 implies that the FLCI is near-optimal for $\delta = 0$. As discussed above, however, such inconsistency does not arise for our baseline smoothness class $\Delta^{SD}(M)$.

Remark 6. In Appendix A.1, we further show that if Assumption 4 along with an additional condition (Assumption 5 introduced below) hold, then the FLCI also has local asymptotic power approaching the power envelope under the same asymptotics considered in Proposition 3.2. ■

The results in this section establish that when certain conditions on Δ are satisfied, the FLCIs are consistent and have desirable finite-sample guarantees on excess length relative to the optimal confidence interval. These conditions hold for our baseline smoothness class Δ^{SD} , but fail for choices of Δ that may be of interest in empirical applications such as those that incorporate sign and monotonicity restrictions. This motivates us to next consider an alternative method for inference in this setting that can accommodate a larger range of restrictions.

4 Inference using Conditional Confidence Sets

In this section, we introduce a more general procedure for inference that has good asymptotic properties over a large class of possible restrictions Δ . We show that inference on the partially identified parameter $\theta = l'\tau_{post}$ in this setting is equivalent to testing a system of moment inequalities with a potentially large number of nuisance parameters that enter linearly. We then apply the conditional approach developed in ARP to obtain computationally tractable inference. We derive novel results on the asymptotic properties of the conditional test in our context, exploiting additional structure in our setting not found in ARP.

4.1 Representation as a moment inequality problem with linear nuisance parameters

Consider the problem of testing the null hypothesis, $H_0 : \theta = \bar{\theta}, \delta \in \Delta$ when $\Delta = \{\delta : A\delta \leq d\}$. We now show that testing H_0 is equivalent to testing a system of moment inequalities with linear nuisance parameters.

The model (1) implies $\mathbb{E}_{(\delta, \tau, \Sigma_n)} [\hat{\beta}_n - \tau] = \delta$, and hence $\delta \in \Delta$ if and only if $\mathbb{E}_{(\delta, \tau, \Sigma_n)} [A\hat{\beta}_n - A\tau] \leq d$. Defining $Y_n = A\hat{\beta}_n - d$ and $M_{post} = [0, I]'$ to be the matrix such that $\tau = M_{post}\tau_{post}$, it is immediate that the null hypothesis H_0 is equivalent to the composite null

$$H_0 : \exists \tau_{post} \in \mathbb{R}^{\bar{T}} \text{ s.t. } l'\tau_{post} = \bar{\theta} \text{ and } \mathbb{E}_{(\delta, \tau, \Sigma_n)} [Y_n - AM_{post}\tau_{post}] \leq 0. \quad (15)$$

In this equivalent form, $\tau_{post} \in \mathbb{R}^{\bar{T}}$ is a vector of nuisance parameters that must satisfy the linear constraint $l'\tau_{post} = \bar{\theta}$.

By applying a change of basis, we can further re-write H_0 as an equivalent composite null hypothesis with an unconstrained nuisance parameter. Re-write the expression $AM_{post}\tau_{post}$ as $\tilde{A} \begin{pmatrix} \theta \\ \tilde{\tau} \end{pmatrix}$, where \tilde{A} is the matrix that results from applying a suitable change of basis to the columns of AM_{post} , and $\tilde{\tau} \in \mathbb{R}^{\bar{T}-1}$.¹⁴ The null H_0 is

¹⁴Specifically, let Γ be a square matrix with the vector l' in the first row and remaining rows chosen so that Γ has full rank. Define $\tilde{A} := AM_{post}\Gamma^{-1}$. Then $AM_{post}\tau = \tilde{A}\Gamma\tau_{post} = \tilde{A} \begin{pmatrix} \theta \\ \underbrace{\Gamma_{(-1, \cdot)}\tau_{post}}_{:=\tilde{\tau}} \end{pmatrix}$.

If $\bar{T} = 1$, then $\tilde{\tau}$ is 0-dimensional and should be interpreted as 0.

then equivalent to

$$H_0 : \exists \tilde{\tau} \in \mathbb{R}^{\bar{T}-1} \text{ s.t. } \mathbb{E} \left[\tilde{Y}_n(\bar{\theta}) - \tilde{X} \tilde{\tau} \right] \leq 0, \quad (16)$$

where $\tilde{Y}(\bar{\theta}) = Y_n - \tilde{A}_{(\cdot,1)} \bar{\theta}$ and $\tilde{X} = \tilde{A}_{(\cdot,-1)}$. Since $\tilde{Y}_n(\bar{\theta})$ is normally distributed with covariance matrix $\tilde{\Sigma}_n = A \Sigma_n A'$ under the finite-sample normal model (1), testing $H_0 : \theta = \bar{\theta}, \delta \in \Delta$ is equivalent to testing a set of moment inequalities with linear nuisance parameters.

Remark 7. The hypothesis (16) is a special case of the testing problem studied in ARP, which focuses on testing null hypotheses of the form $H_0 : \exists \tau \text{ s.t. } \mathbb{E} [Y(\theta) - X\tau | X] \leq 0$, almost surely. Our setting is a special case of this framework in which: i) the variable X takes the degenerate distribution $X = \tilde{X}$, and ii) $Y(\theta) = \tilde{Y}(\theta)$ is linear in θ . The first feature plays an important role in developing our novel consistency and local asymptotic power results presented later in this section: if i) fails and X is continuously distributed, then the tests proposed by ARP will generally not be consistent, as they do not allow for the number of moments to grow with n . The current proof of the optimal local asymptotic results also exploit the geometry of feature ii), although we conjecture that this could be relaxed to allow $Y(\theta)$ to vary smoothly in θ . ■

4.2 Constructing conditional confidence sets

An important practical consideration for testing hypotheses of the form (16) is that the dimension of the nuisance parameter $\tilde{\tau} \in \mathbb{R}^{\bar{T}-1}$ grows linearly with the number of post-periods \bar{T} and may be large in practice. For instance, in Section 8 we apply our methodology to a recent paper in which $\bar{T} = 23$. Moreover, 5 of the 12 recent event-study papers reviewed in Roth (2019) have $\bar{T} > 10$. This renders many moment inequality methods, especially those which rely on test inversion over a grid for the full parameter vector, practically infeasible in this context. We now show how the conditional approach of ARP, which directly exploits the linear structure of the hypothesis (16), can be applied to obtain computationally tractable and powerful tests even when the number of post-periods \bar{T} is large.¹⁵

¹⁵Other moment inequality methods have been proposed for subvector inference, but typically do not exploit the linear structure of our setting — see, e.g, Chen, Christensen and Tamer (2018); Bugni, Canay and Shi (2017); Kaido, Molinari and Stoye (2019a); Chernozhukov, Newey and Santos (2015); Romano and Shaikh (2008). Gafarov (2019), Cho and Russell (2019), and Flynn (2019)

Suppose we wish to test (16) for some fixed $\bar{\theta}$. The conditional testing approach considers tests based on the test statistic

$$\hat{\eta} := \min_{\eta, \tilde{\tau}} \eta \text{ s.t. } \tilde{Y}_n(\bar{\theta}) - \tilde{X}\tilde{\tau} \leq \tilde{\sigma}_n \cdot \eta, \quad (17)$$

where $\tilde{\sigma}_n = \sqrt{\text{diag}(\tilde{\Sigma}_n)}$. This linear program selects the value of the nuisance parameters $\tilde{\tau} \in \mathbb{R}^{\tilde{T}-1}$ that produces the most slack in the maximum studentized moment. Duality results from linear programming (e.g. Schrijver (1986), Section 7.4) imply that the value $\hat{\eta}$ obtained from the primal program (17) equals the optimal value of the dual program,¹⁶

$$\hat{\eta} = \max_{\gamma} \gamma' \tilde{Y}_n(\bar{\theta}) \text{ s.t. } \gamma' \tilde{X} = 0, \gamma' \tilde{\sigma}_n = 1, \gamma \geq 0. \quad (18)$$

If a vector γ_* is optimal in the dual problem above, then it is a vector of Lagrange multipliers for the primal problem. We denote by \hat{V}_n the set of optimal vertices of the dual program.¹⁷

To derive critical values, we analyze the distribution of $\hat{\eta}$ conditional on the event that a vertex γ_* is optimal in the dual problem. Lemma 9 of ARP shows that conditional on the event $\gamma_* \in \hat{V}_n$ and a sufficient statistic S_n for the nuisance parameters, the test statistic $\hat{\eta}$ follows a truncated normal distribution with

$$\hat{\eta} | \{\gamma_* \in \hat{V}_n, S_n = s\} \sim \xi | \xi \in [v^{lo}, v^{up}], \quad (19)$$

where $\xi \sim \mathcal{N}\left(\gamma'_* \tilde{\mu}, \gamma'_* \tilde{\Sigma}_n \gamma_*\right)$, $\tilde{\mu} = \mathbb{E}\left[\tilde{Y}_n(\bar{\theta})\right]$, $S_n = \left(I - \frac{\tilde{\Sigma}_n \gamma_* \gamma'_*}{\gamma'_* \tilde{\Sigma}_n \gamma_*}\right) \tilde{Y}_n(\bar{\theta})$, and v^{lo}, v^{up} are known functions of $\tilde{\Sigma}_n, s, \gamma_*$.¹⁸ All quantiles of the conditional distribution of $\hat{\eta}$

also provide methods for subvector inference with linear moment inequalities, but in contrast to our approach require a linear independence constraint qualification (LICQ) assumption for size control.

¹⁶Technically, the duality results require that $\hat{\eta}$ be finite. However, one can show that $\hat{\eta}$ is finite with probability 1, unless the span of \tilde{X} contains a vector with all negative entries, in which case the identified set for θ is the real line. We therefore trivially define our test never to reject if $\hat{\eta} = -\infty$.

¹⁷In general, there may not be a unique solution to the dual program. However, Lemma 11 of ARP shows that conditional on any one vertex of the dual program's feasible set being optimal, every other vertex is optimal with either probability 0 or 1. It thus suffices to condition on the event that a vector $\gamma_* \in \hat{V}$.

¹⁸The cutoffs v^{lo} and v^{up} are the maximum and minimum of the set $\{x : x = \max_{\gamma \in F_n} \gamma'(s + \frac{\tilde{\Sigma}_n \gamma_*}{\gamma'_* \tilde{\Sigma}_n \gamma_*} x)\}$ when $\gamma'_* \tilde{\Sigma}_n \gamma_* \neq 0$, where F_n is the feasible set of the dual program (18). When $\gamma'_* \tilde{\Sigma}_n \gamma_* = 0$, we define $v^{lo} = -\infty$ and $v^{up} = \infty$, so the conditional test rejects if and only if $\hat{\eta} > 0$.

in the previous display are increasing in $\gamma'_* \tilde{\mu}$,¹⁹ and the null hypothesis (16) implies $\gamma'_* \tilde{\mu} \leq 0$.

We therefore select the critical value for the conditional test to be the $1 - \alpha$ quantile of the truncated normal distribution $\xi | \xi \in [v^{lo}, v^{up}]$ under the worst-case assumption that $\gamma'_* \tilde{\mu} = 0$. Let $\psi_\alpha^C(\tilde{Y}_n(\bar{\theta}), \tilde{\Sigma}_n)$ denote an indicator for whether the conditional test rejects at the $1 - \alpha$ level. The conditional test is defined as

$$\psi_\alpha^C(\tilde{Y}_n(\bar{\theta}), \tilde{\Sigma}_n) = 1 \iff F_{\xi | \xi \in [v^{lo}, v^{up}]}(\hat{\eta}; \gamma'_* \tilde{\Sigma}_n \gamma_*) > 1 - \alpha, \quad (20)$$

where $F_{\xi | \xi \in [v^{lo}, v^{up}]}(\cdot; \sigma^2)$ is the CDF of $\xi \sim \mathcal{N}(0, \sigma^2)$ truncated to $[v^{lo}, v^{up}]$. It follows immediately from Proposition 6 in ARP that the conditional test controls size,

$$\sup_{\delta \in \Delta, \tau} \sup_{\theta \in \mathcal{S}(\Delta, \delta + \tau)} \mathbb{E}_{(\delta, \tau, \Sigma_n)} \left[\psi_\alpha^C(\tilde{Y}_n(\theta), \tilde{\Sigma}_n) \right] \leq \alpha. \quad (21)$$

A confidence set satisfying the uniform coverage criterion (9) can be constructed by test inversion for the scalar parameter θ . The conditional confidence set is given by

$$\mathcal{C}_{\alpha, n}^C := \{\bar{\theta} : \psi_\alpha^C(\tilde{Y}_n(\bar{\theta}), \tilde{\Sigma}_n) = 0\}. \quad (22)$$

Remark 8. For each value of $\bar{\theta}$, the test statistic $\hat{\eta}$ can be computed by solving the linear program (17). To form the confidence set $\mathcal{C}_{\alpha, n}^C$, one only needs to perform test inversion over a grid of values for the scalar parameter $\bar{\theta}$, and thus the problem remains highly tractable even when \bar{T} is large. Moreover, the commonly-used dual simplex algorithm for linear programming returns a vertex to the dual solution (18), so an optimal dual vertex γ_* can be obtained from standard solvers without further calculation. ■

Remark 9. Consider the simple setting in which we have one post-period ($\bar{T} = 1$) and are interested in the treatment effect in the first period, $\theta = \tau_1$. In this case, there are no nuisance parameters, and the form of the conditional test simplifies substantially. The test statistic $\hat{\eta}$ is the maximum of the studentized moments, $\hat{\eta} = \max_j \tilde{Y}_{n,j} / \tilde{\sigma}_{n,j}$, where $\tilde{\sigma}_{n,j}$ is the standard deviation of $\tilde{Y}_{n,j}$. The conditional test rejects in this case if and only if $\frac{\Phi(\hat{\eta}) - \Phi(v^{lo})}{1 - \Phi(v^{lo})} > 1 - \alpha$. Moreover, if the moments \tilde{Y}_n are uncorrelated

¹⁹This follows from the fact that the truncated normal distribution $\xi | \xi \in [v^{lo}, v^{up}]$ has the monotone likelihood ratio property in its mean (see, e.g. Lemma A.1 in Lee, Sun, Sun and Taylor (2016)).

with each other, then v^{lo} is the maximum of the non-binding studentized moments, $v^{lo} = \max_{j \neq \hat{j}} \tilde{Y}_{n,j} / \tilde{\sigma}_{n,j}$, where \hat{j} denotes the location of the maximum studentized moment. ■

4.3 Consistency and optimal local asymptotic power of conditional confidence sets

We now present our two novel results on the asymptotic power of the conditional test in our setting. We first show that the conditional test is consistent, meaning that any fixed point outside of the identified set is rejected with probability approaching one as the sample size $n \rightarrow \infty$.

Proposition 4.1. *The conditional test is consistent. For any $\delta_A \in \Delta$, $\tau_A \in \mathbb{R}^T$, and $\theta^{out} \notin \mathcal{S}(\delta_A + \tau_A, \Delta)$,*

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} (\theta^{out} \notin \mathcal{C}_{\alpha, n}^C) = 1.$$

We next consider the local asymptotic power of the conditional test. We provide a condition under which the power of the conditional test converges to the power envelope in a $n^{-\frac{1}{2}}$ -neighborhood of the identified set. This condition guarantees that the binding and non-binding moments are sufficiently well-separated at points close to the boundary of the identified set.

Assumption 5. *Let $\Delta = \{\delta : A\delta \leq d\}$ and fix $\delta_A \in \Delta$. Consider the optimization:*

$$b^{max}(\delta_{A,pre}) = \max_{\delta} l' \delta_{post} \text{ s.t. } A\delta \leq d, \delta_{pre} = \delta_{A,pre},$$


and assume it has a finite solution. For δ^ a maximizer to the above problem, let $B(\delta^*)$ index the set of binding inequality constraints, so that $A_{(B(\delta^*), \cdot)} \delta^* = d_{B(\delta^*)}$ and $A_{(-B(\delta^*), \cdot)} \delta^* - d_{-B(\delta^*)} = -\epsilon_{-B(\delta^*)} < 0$. Assume that there exists a maximizer δ^* to the problem above such that the rank of $A_{(B(\delta^*), post)}$ is equal to $|B(\delta^*)|$. Analogously, assume that there is a finite solution to the analogous problem that replaces max with min, and that there is a minimizer δ^{**} such that $A_{(B(\delta^{**}), post)}$ has rank $|B(\delta^{**})|$.*

Assumption 5 considers the problem of finding the differential trend $\delta \in \Delta$ that is consistent with the pre-trend identified from the data $(\delta_{A,pre})$ and causes $l' \hat{\beta}_{post}$ to be maximally biased for $\theta := l' \tau_{post}$. It requires that the “right” number of moments bind when we do this optimization.

Remark 10. Assumption 5 is closely related to, but slightly weaker than, linear independence constraint qualification (LICQ). LICQ has been used recently in the moment inequality settings of Gafarov (2019); Cho and Russell (2019); Flynn (2019) and Kaido and Santos (2014); see Kaido, Molinari and Stoye (2019b) for a synthesis. We show in Appendix A.2 that LICQ is equivalent to a modified version of Assumption 5 that replaces “there exists a maximizer δ^* ” with “for every maximizer δ^* ” (and analogously for the minimizer δ^{**}). Thus, LICQ is equivalent to Assumption 5 when the optimizations considered in Assumption 5 have unique solutions, but potentially weaker when there are multiple solutions. We note that many of the mentioned papers require LICQ for asymptotic size control, whereas we only impose Assumption 5 for our results on local asymptotic power. ■

Remark 11. In the special case with one pre-period ($\bar{T} = 1$) and one post-period ($\bar{T} = 1$), Assumption 5 has a simple graphical interpretation. It is satisfied whenever Δ has non-empty interior and δ is not vertically aligned with a vertex. Figure 4 shows the areas at which Assumption 5 holds/fails for three of our ongoing examples. The assumption holds everywhere for Δ^{SD} when $M > 0$, and Lebesgue almost everywhere for Δ^{SDPB} and Δ^{RMI} when $M > 0$ and $\bar{M} > 0$. The sets $\Delta^{SD}(0)$, $\Delta^{SDPB}(0)$, $\Delta^{RMI}(0)$ all have empty interior, and so Assumption 5 fails for these cases (in which θ is point-identified). More generally, one can show that Assumption 5 does not hold if θ is point identified. ■

Figure 4: Diagram of where Assumption 5 holds. The assumption holds (fails) for values of δ plotted in green (red).



Let $\mathcal{I}_\alpha(\Delta, \Sigma_n)$ denote the class of confidence sets that satisfy the finite sample

coverage criterion in (9) at the $1 - \alpha$ level. Under Assumption 5, the power of the conditional test converges to the optimum over $\mathcal{I}_\alpha(\Delta, \Sigma_n)$ in a $n^{-1/2}$ -neighborhood of the identified set as $n \rightarrow \infty$.

Proposition 4.2. *Fix $\delta_A \in \Delta, \tau_A$, and suppose Σ^* is positive definite. Let $\theta_A^{ub} = \sup_\theta \mathcal{S}(\delta_A + \tau_A, \Delta)$ be the upper bound of the identified set. Suppose Assumption 5 holds. Then, for any $x > 0$,*

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta_A^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n}^C \right) &= \lim_{n \rightarrow \infty} \sup_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_\alpha(\Delta, \Sigma_n)} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta_A^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n} \right) \\ &= \Phi(c^*x - z_{1-\alpha}), \end{aligned}$$

for a positive constant c^* .²⁰ The analogous result holds replacing $\theta_A^{ub} + \frac{1}{\sqrt{n}}x$ with $\theta_A^{lb} - \frac{1}{\sqrt{n}}x$, for θ_A^{lb} the lower bound of the identified set (although the constant c^* may differ).

We now provide an outline of the proof of our local asymptotic optimality result, the details of which are contained in the appendix. The proof proceeds in two parts: the first part characterizes the asymptotically optimal test, and the second shows that our conditional test converges to this optimal test.

We first show that under Assumption 5, the local asymptotic power of any test that controls size is bounded above by that of a particular one-sided t-test. Specifically, Assumption 5 implies that there is a unique set of Lagrange multipliers $\bar{\gamma}$ in the “population version” of the test statistic $\hat{\eta}(\theta^{ub})$ that replaces $\tilde{Y}(\theta^{ub})$ with its expectation $\tilde{\mu}(\theta^{ub})$ in (17). We then show that testing $H_0 : \theta = \theta^{ub}, \delta \in \Delta$ against a local alternative can be represented as a test of a convex null against a point alternative in the normal location model. Applying the Neyman-Pearson lemma, we show that the optimal test is a one-sided t-test in the direction of $\bar{\gamma}$ for alternatives sufficiently close to θ^{ub} .

We next show that the conditional test converges in probability to the optimal one-sided t-test discussed above. Since Assumption 5 implies that $\bar{\gamma}$ is the unique dual solution to the “population version” of $\hat{\eta}$, it follows that $\bar{\gamma}$ will be optimal in the dual problem for $\hat{\eta}$ with probability approaching one as $\Sigma_n \rightarrow 0$. Thus, with

²⁰In particular, letting $B = B(\delta^{**})$ as defined in Assumption 5, $c^* = -\bar{\gamma}'_B \tilde{A}_{(B,1)} / \sigma_B$, where $\sigma_B = \sqrt{\bar{\gamma}'_B A_{(B,\cdot)} \Sigma^* A'_{(B,\cdot)} \bar{\gamma}_B}$ and $\bar{\gamma}_B$ is a non-zero vector such that $\bar{\gamma}'_B \tilde{A}_{(B,-1)} = 0, \bar{\gamma}_B \geq 0$. The vector $\bar{\gamma}_B$ is unique up to scale.

probability approaching one the test statistic for the conditional test will be $\hat{\eta} = \bar{\gamma}'\tilde{Y}$, which corresponds with that of the one-sided t-test. Finally, recall that the critical value of the conditional test is based on the $1 - \alpha$ quantile of the distribution of $\gamma'_*\tilde{Y}$ conditional on γ_* being optimal. However, since $\bar{\gamma}$ is optimal with probability approaching 1, the distribution of $\bar{\gamma}'\tilde{Y}$ conditional on $\bar{\gamma}$ being optimal approaches its unconditional distribution, which is normal. Thus, the critical value of the conditional test approaches the $1 - \alpha$ quantile of the normal distribution.

Remark 12. Equation (21) and Proposition 4.2 together show that the conditional approach uniformly controls size over all values of $\delta \in \Delta$, and is asymptotically efficient when δ further satisfies Assumption 5. In general, we cannot guarantee that the bounds of the identified set will be differentiable as a function of $\beta = \delta + \tau$, and the impossibility results in Hirano and Porter (2012) imply that no regular estimators of the identified set bounds exist when differentiability fails. However, one can show that if Assumption 5 holds, then the identified set bounds are differentiable in β . Proposition 4.2 therefore implies that although the conditional test controls size uniformly for all values of $\delta \in \Delta$, this does not come at the expense of efficiency in cases where Assumption 5 holds. Since researchers often do not know ex ante whether Assumption 5 is satisfied, this “robustness” property is desirable. Our results are thus somewhat analogous to results in the weak identification literature that show that certain procedures control size under weak identification but are efficient under strong identification (e.g., Moreira (2003)).

Remark 13 (Relationship to other methods). We are not aware of results analogous to Proposition 4.2 for any other moment inequality procedure that controls size in the finite sample normal model. Observe that if Assumption 5 holds, then it also holds if Δ is augmented to include a moment that is non-binding at both endpoints of the identified set. Hence, for Proposition 4.2 to hold, the local asymptotic power of the test needs to be unaffected by the inclusion of such slack moments. Although relatively insensitive to the inclusion of slack moments, the procedures of Romano, Shaikh and Wolf (2014) and Andrews and Barwick (2012) are still affected by the inclusion of slack moments via the changes to the first-stage critical value and size-adjustment factor, respectively. The method of Cox and Shi (2019) is strongly insensitive to the inclusion of slack moments, but it does not converge to a one-sided t-test as the other moments become slack and thus does not approach the power envelope.

Remark 14 (Finite sample power of the conditional test). Note that the argument above for the optimality of the conditional approach relies on a unique vector of Lagrange multipliers $\bar{\gamma}$ being dual-optimal with probability approaching 1 asymptotically. The asymptotic guarantees of Proposition 4.2 thus may not translate to good finite-sample performance in settings where multiple vectors of Lagrange multipliers are optimal with high probability. Since a vector of Lagrange multipliers corresponds with a set of active moments in the primal problem (17), this will tend to occur in cases where the set of binding and non-binding moments are not “well-separated” relative to the sampling variation in the data. Such a situation will tend to arise when Assumption 5 is “close” to being violated.²¹ ■

5 Conditional-FLCI Hybrid Confidence Sets

Taken together, the results in Section 3 show that the FLCIs have attractive finite-sample properties for particular classes Δ of interest, but they may perform poorly even asymptotically for other types of restrictions. On the other hand, the conditional tests have good asymptotic properties for a wider range of restrictions, but they may perform poorly in finite-sample in settings where the binding and non-binding moments are not well-separated relative to the sampling variation in the data. To address these differing strengths, we propose a novel confidence set that hybridizes the conditional test with the optimal FLCI based on an affine estimator. This conditional-FLCI hybrid confidence set achieves similar desirable asymptotic properties as the conditional confidence set for a wide range of Δ s. In simulations, it leads to substantial improvements in the power of the conditional test in a variety of cases where the moments are not well-separated.

The conditional-FLCI hybrid confidence set is constructed by first testing whether a candidate parameter value lies within the optimal level- $(1 - \kappa)$ FLCI, and then applying a conditional test to all parameter values that lie within the optimal FLCI. In the second stage, we use a modified version of the conditional test that i) adjusts size to account for the first-stage test, and ii) conditions on the event that the first-stage test fails to reject.

²¹Indeed, in the supplementary material we prove a uniform version of the optimality result in Proposition 4.2 under a modified version of Assumption 5 that requires the non-binding moments to be uniformly bounded away from zero.

Formally, suppose that $0 < \kappa < \alpha$.²² Consider the level $(1 - \kappa)$ optimal FLCI, $\mathcal{C}_{\kappa,n}^{FLCI} = a_n + v'_n \hat{\beta}_n \pm \chi_n$. Lemma A.3 shows that the distribution of the test statistic $\hat{\eta}$ defined in (17) follows a truncated normal distribution conditional on the parameter value $\bar{\theta}$ falling within the level $(1 - \kappa)$ optimal FLCI. With this result, the construction of the second-stage of the conditional-FLCI hybrid test is analogous to the construction of the conditional test, except it uses the modified size $\tilde{\alpha} = \frac{\alpha - \kappa}{1 - \kappa}$ to account for the first-stage test. The conditional-FLCI hybrid test ψ_{C-FLCI} equals

$$\psi_{\kappa,\alpha}^{C-FLCI}(\hat{\beta}_n, \bar{\theta}, \tilde{\Sigma}_n) = 1 \iff \theta \notin \mathcal{C}_{\kappa,n}^{FLCI}, \text{ OR } F_{\xi|\xi \in [v_{C-FLCI}^l, v_{C-FLCI}^u]}(\hat{\eta}) > 1 - \tilde{\alpha},$$

where $F_{\xi|\xi \in [v_{C-FLCI}^l, v_{C-FLCI}^u]}(\cdot)$ denotes the CDF of the truncated normal distribution derived in Lemma A.3.

Since the FLCI controls size, the first stage test rejects with probability at most κ under the null that $\theta = \bar{\theta}$. The second-stage test rejects with probability at most $\tilde{\alpha} = \frac{\alpha - \kappa}{1 - \kappa}$ conditional on $\theta \in \mathcal{C}_{\kappa,n}^{FLCI}$. Together, these results imply that the conditional-FLCI hybrid test controls size,

$$\sup_{\delta \in \Delta, \tau} \sup_{\theta \in \mathcal{S}(\Delta, \delta + \tau)} \mathbb{E}_{(\delta, \tau, \Sigma_n)} \left[\psi_{\kappa,\alpha}^{C-FLCI}(\hat{\beta}_n, \bar{\theta}, \tilde{\Sigma}_n) \right] \leq \alpha. \quad (23)$$

We therefore construct a conditional-FLCI hybrid confidence set for the parameter θ that satisfies (9) by inverting the conditional-FLCI test,

$$\mathcal{C}_{\kappa,\alpha,n}^{C-FLCI} = \{\theta : \psi_{\kappa,\alpha}^{C-FLCI}(\hat{\beta}_n, \bar{\theta}, \tilde{\Sigma}_n) = 0\}. \quad (24)$$

In Appendix A.3, we show that the conditional-FLCI hybrid confidence set inherits some desirable asymptotic properties from the conditional approach: it is asymptotically consistent, and under the same conditions as Proposition 4.2, the conditional-FLCI hybrid test has local asymptotic power at least as good as the optimal $\frac{\alpha - \kappa}{1 - \kappa}$ test.

²²In practice, we set $\kappa = \alpha/10$ following Romano et al. (2014) and ARP, although the optimal choice of κ is an interesting question for future research.

6 Simulation study

In this section, we present a series of simulations illustrating the performance of the confidence sets discussed above across a range of relevant data-generating processes. We find good size control for all three of our proposed procedures, and therefore focus in the main text on a comparison of power. In the supplementary material, we present results on size control and other additional simulation results.

6.1 Simulation Design

Our simulations are calibrated using the estimated covariance matrix from the 12 recently-published papers surveyed in Roth (2019).²³ For any given paper in the survey, we denote by $\hat{\Sigma}$ the estimated variance-covariance matrix from the event-study in the paper, calculated using the clustering scheme specified by the authors. We then simulate event-study coefficients $\hat{\beta}_s$ from a normal model under the assumption of parallel trends and zero treatment effects, $\hat{\beta}_s \sim \mathcal{N}(0, \hat{\Sigma})$.²⁴ In simulation s , we construct nominal 95% confidence intervals for the parameter of interest θ using the pair $(\hat{\beta}_s, \hat{\Sigma})$ for each proposed procedure. The parameter of interest is the causal effect in the first post-period ($\theta = \tau_1$).²⁵

For a given choice of Δ , we compute the identified set $\mathcal{S}(0, \Delta)$ and calculate the expected excess length for each of the proposed confidence sets. The excess length of a confidence set $\mathcal{C}(\hat{\beta})$ is the length of the part of the confidence set that falls outside of the identified set, defined as $EL(\mathcal{C}; \hat{\beta}) = \lambda(\mathcal{C}(\hat{\beta}) \setminus \mathcal{S}(0, \Delta))$. We benchmark the expected excess length of our proposed procedures relative to the optimal bound over confidence sets that satisfy the uniform coverage requirement (9).²⁶ For each

²³Roth (2019) systematically reviewed the set of papers containing an event-study plot published in the *American Economic Review*, *AEJ: Applied Economics* and *AEJ: Economic Policy* between 2014 and mid-2018. Section 4.1 of Roth (2019) discusses the sample selection criteria for this survey of published event studies.

²⁴We focus on the normal simulations in the main text since it allows for a tractable computation of the optimal excess length of procedures that control size. In the supplementary material, we show that our procedures perform similarly in simulations based on the empirical distribution in the original paper.

²⁵In the supplementary material, we provide simulation results in which the parameter of interest is the average causal effect in the post-periods ($\theta = \bar{\tau}_{post}$), and find the results are qualitatively unchanged.

²⁶A formula for this optimal bound follows from Theorem 3.2 of Armstrong and Kolesar (2018), which provides an expression for the optimal expected length of a confidence set that satisfies the uniform coverage requirement.

paper, we conduct 1000 simulations and compute the optimal bound and the average excess length for the conditional confidence set, FLCI, and conditional-FLCI hybrid confidence set. The excess length efficiency of a given procedure equals the ratio of the optimal bound to the simulated expected excess length. We report the efficiency ratios for the median paper in the survey.

We consider three choices of Δ to highlight the performance of the confidence sets across a range of conditions: $\Delta^{SD}(M)$, $\Delta^{SDPB}(M)$ and $\Delta^{SDI}(M)$. Table 1 summarizes which of our theoretical results hold for each of the simulation designs.


	Δ^{SD}	Δ^{SDPB}	Δ^{SDI}
Conditional / Hybrid			
Consistent	✓	✓	✓
Asymptotically (near-)optimal	✓	✓	×
FLCI			
Consistent	✓	×	×
Asymptotically (near-)optimal	✓	×	×
Finite-sample near-optimal	✓	×	×

Table 1: Summary of expected properties for each simulation design

In all simulations, the units of the parameter M are standardized to equal the standard error of the first post-period event study coefficient (σ_1). We show results for a variety of choices of M/σ_1 . All of the procedures and the optimal bounds are invariant to scale, meaning the confidence set using $(M, \frac{1}{n}\Sigma^*)$ is $\frac{1}{n}$ times the confidence set using (nM, Σ^*) . Therefore, simulation results on excess length as M/σ_1 grows large are isomorphic to our asymptotic results presented earlier in which $n \rightarrow \infty$ for $\Sigma_n = \frac{1}{n}\Sigma^*$ and $M > 0$ is fixed. The simulation results also have a finite-sample interpretation, illustrating how our results change as we allow the set of underlying trends to be more non-linear, holding Σ^* constant.

The supplementary materials present results from several alternative simulation exercises. We find similar results using the empirical distribution from the first paper in Roth (2019)'s survey rather than the calibrated normal model studied in the main text. We also find qualitatively similar results when using the average of the post-treatment causal effects as the target parameter.

Figure 5: Simulation results: Median efficiency ratios for proposed procedures.



Note: Median efficiency ratios for our proposed confidence sets. The efficiency ratio for a procedure is defined as the optimal bound divided by the procedure's expected excess length. The results for the FLCI are plotted in green, the results for the conditional-FLCI hybrid confidence interval in red and the results for the conditional confidence interval in blue. Results are averaged over 1000 simulations for each of the 12 papers surveyed, and the median across papers is reported here.

6.2 Simulation Results

Results for $\Delta^{SD}(M)$: The top panel of Figure 5 plots the efficiency ratio for each procedure as a function of M/σ_1 when $\Delta = \Delta^{SD}$. All procedures perform well as M/σ_1 grows large with efficiency ratios approaching 1, illustrating our asymptotic (near-)optimality results. The FLCIs also perform quite well for smaller values of M/σ_1 , including the point-identified case where $M = 0$, illustrating the finite-sample near-optimality results for the FLCIs when Assumption 3 holds. Although the conditional confidence sets have efficiency approaching the optimal bound for M/σ_1 large, their efficiency when $M/\sigma_1 = 0$ is only about 50%. This reflects the fact that when $M = 0$, the parameter is point-identified and Assumption 5 fails. The conditional-FLCI hybrid substantially improves efficiency for small values of M/σ_1 , while still retaining near-optimal performance as M/σ_1 grows large.

Results for $\Delta^{SDPB}(M)$: The middle panel of Figure 5 plots the efficiency ratio for each procedure as a function of M/σ_1 when $\Delta = \Delta^{SDPB}$. The efficiency ratios for the conditional and hybrid confidence sets are (near-)optimal as M/σ_1 grows large, highlighting our asymptotic (near-)optimality results for these procedures in this simulation design. However, the efficiency ratios for the FLCIs steadily decrease as M/σ_1 increases, which reflects the fact that the FLCIs are not consistent in this simulation design when $M > 0$. We again see that the conditional-FLCI hybrid improves efficiency when M/σ_1 is small, while retaining near-optimal performance as M/σ_1 grows large.

Results for Δ^{SDI} The bottom panel of Figure 5 plots the efficiency ratio for each procedure as a function of M/σ_1 when $\Delta = \Delta^{SDI}$. As summarized in Table 1, the conditions for asymptotic (near-)optimality do not hold for any of our procedures in this simulation design. Nonetheless, the conditional and hybrid procedures still perform quite well for large values of M/σ_1 , with efficiency approaching about 90%. This evidence is encouraging as it shows that they may perform well asymptotically in cases where Assumption 5 fails. The efficiency of the FLCIs degrades as M/σ_1 grows, reflecting that the FLCIs are inconsistent under this simulation design when $M > 0$. Once again, the conditional-FLCI hybrid improves efficiency when M/σ_1 is small, while retaining similar performance to the conditional approach as M/σ_1 grows large.

7 Practical Guidance

We now provide practical guidance on how these methods may be used to assess the robustness of conclusions in difference-in-differences and event-study designs. We recommend that applied researchers conduct the following steps to assess the robustness of their conclusions in difference-in-differences and related designs.

- 1) Estimate an “event-study”-type specification that produces a vector of estimates $\hat{\beta}$, consisting of “pre-period” coefficients $\hat{\beta}_{pre}$ and “post-period” coefficients $\hat{\beta}_{post}$, where the post-period coefficients have a causal interpretation under a suitable parallel trends assumption.
- 2) Perform a sensitivity analysis where inference is conducted under different assumptions about the set of possible violations of parallel trends Δ .
- 3) Provide economic benchmarks for evaluating the different choices of Δ . This involves using context-specific knowledge about potential confounding factors or using information from pre-treatment periods and placebo groups.

We recommend researchers select either the optimal FLCI or the conditional-FLCI hybrid conditional confidence set based upon the properties of their specified choice of Δ . For cases (e.g. $\Delta = \Delta^{SD}(M)$) where the conditions for the consistency of the FLCIs are non-restrictive and the conditions for finite-sample near-optimality under parallel trends hold, we recommend that the researchers use the optimal FLCI. Outside of these special cases – e.g., when context-specific knowledge motivates sign or shape restrictions – we recommend that researchers use the conditional-FLCI hybrid confidence set. Our R package, `HonestDiD`, implements our methods and chooses the recommended procedure by default.

7.1 When to Use Our Methods

The methods in this paper can be applied in most empirical settings in which researchers use an “event-study plot” to evaluate pre-existing trends. Our methods require the researcher to use an estimator $\hat{\beta}_n$ with an asymptotic normal limit, $\sqrt{n}(\hat{\beta}_n - \beta) \rightarrow_d \mathcal{N}(0, \Sigma^*)$, and that the reduced-form parameter β satisfies the causal decomposition in Assumption 1. We now address two considerations that commonly arise in practice: i) staggered treatment timing, and ii) anticipatory effects.

First, a recent literature has shown that the coefficients from standard two-way fixed effect models may not be causally interpretable in the presence of staggered treatment timing and heterogeneous treatment effects across cohorts. To address these issues, [Sun and Abraham \(2020\)](#) and [Callaway and Sant’Anna \(2019\)](#) provide alternative strategies for estimating weighted averages of cohort-specific treatment effects at a fixed lag (or for placebo analysis, lead) relative to treatment, which yield consistent estimates under a suitable parallel trends assumption.²⁷

These estimates are asymptotically normal under mild regularity conditions, and so our recommended sensitivity analysis can be applied to gauge sensitivity to violations of the needed parallel trends assumption. In empirical settings with staggered treatment timing and heterogeneous treatment effects, we therefore recommend that researchers first use the methods of [Sun and Abraham \(2020\)](#) or [Callaway and Sant’Anna \(2019\)](#) for estimation and then apply our results to conduct sensitivity analysis.

Next, in some cases, there may be changes in behavior in anticipation of the policy of interest, and therefore, β_{pre} may reflect the anticipatory effects of the policy (e.g., see [Malani and Reif \(2015\)](#)). This violates Assumption 1, which assumes pre-treatment coefficients do not reflect causal effects. A simple solution is available if one is willing to assume that anticipatory effects only occur in a fixed window prior to the policy change. Under such an assumption, the researcher may re-normalize the definition of the “pre-treatment” period to be the period prior to when anticipatory effects can occur, in which case β_{pre} is determined only based on untreated potential outcomes.

7.2 Sensitivity Analysis

We recommend that researchers report confidence sets under different assumptions about the set of possible violations of parallel trends Δ . This allows the reader to evaluate what assumptions need to be imposed in order to obtain informative inference.

For instance, in many cases a reasonable baseline choice for Δ may be $\Delta^{SD}(M)$,

²⁷The literature on staggered treatment timing considers generalizations of the parallel trends assumption that impose that untreated potential outcomes for each treated cohort move in parallel to those for some control group; possibilities for the control group include never-treated units, not-yet-treated units, or the last cohort to be treated. See, e.g., Assumption 2 of [Callaway and Sant’Anna \(2019\)](#) or Section 4.2 in [Sun and Abraham \(2020\)](#).

which relaxes the assumption of linear differences in trends by imposing that the slope of the differential trend can change by no more than M in consecutive periods. By reporting robust confidence sets for different values of M , the researcher may evaluate the extent to which their conclusions change as we allow for the possibility of greater non-linearities in the underlying trend. If there is a particular null hypothesis of interest (e.g. the treatment effect is zero), the researcher can report the “breakdown” value of M at which the null hypothesis can no longer be rejected.²⁸ The researcher may also assess how these conclusions change if they impose further sign or shape restrictions motivated by context-specific economic knowledge — e.g. the bias is weakly positive, or the trend is monotone.

7.3 Evaluating a choice of Δ

When conducting sensitivity analysis over possible choices of Δ , researchers may wonder how to evaluate whether a particular choice of Δ is reasonable. For instance, suppose a researcher is considering sets of the form $\Delta^{SD}(M)$. How, then, should she evaluate what value of possible non-linearity M is reasonable? For ease of exposition, we focus on how to evaluate the choice of M , although analogous remarks apply to conducting sensitivity for other forms of Δ as well.

Nothing in the data itself can place an upper bound on the parameter M . This is because M bounds the magnitude of possible non-linearity in the counterfactual difference in trends δ_{post} and thus is not identified. Applied researchers therefore must use domain-specific knowledge to evaluate whether a particular choice of M is plausible in a given empirical setting. To help applied researchers do so, we discuss two approaches that benchmark possible values of M using knowledge of potential confounds and using untreated groups or periods.

Benchmarking using knowledge of potential confounds. Applied researchers may calibrate M by considering the types of economic mechanisms that would produce a violation of the parallel trends assumption, and benchmark M using knowledge of the likely magnitudes of those mechanisms. This suggestion echoes [Kahn-Lang and Lang \(2019\)](#), who advocate for the use of domain-specific knowledge in justifying the

²⁸Formally, one can show that if $\mathcal{C}(M)$ satisfies (9) for $\Delta^{SD}(M)$, then the sample breakdown value, $\hat{M}_{BE} := \inf\{M : \theta_0 \in \mathcal{C}(M)\}$, is a valid $(1 - \alpha)$ -level lower bound for the population breakdown value, $M_{BE} := \inf\{M : \theta_0 \in \mathcal{S}(\Delta^{SD}(M), \beta)\}$. That is, $\mathbb{P}(\hat{M}_{BE} \leq M_{BE}) \geq 1 - \alpha$.

parallel trends assumption. We provide an example of this type of reasoning in our application to [Lovenheim and Willen \(2019\)](#) in Section 8. ■

Benchmarking using untreated groups or periods. In some settings, researchers may observe untreated (placebo) groups who they think are “at least as dissimilar” as the groups used in their main analysis. In this case, it may be sensible to form a confidence interval for the largest change in slope between periods among the placebo groups, and use the upper bound of that interval as a benchmark for M .²⁹ Along similar lines, one could also construct an upper bound on the largest change in slope in the pre-period for the groups used in the main event-study of interest. One could then benchmark M in terms of multiples of the largest value observed in the pre-period. ■

Finally, while an upper bound on the parameter M may not be learned from the data, the observed pre-period data may provide an informative *lower bound* on possible values of M . Since M is an upper limit on how much the slope of δ can change between periods, the largest second difference in δ_{pre} must be no larger than M in absolute value. As δ_{pre} equals the expected value of the pre-period coefficients $\hat{\beta}_{pre}$, we can test this hypothesis using the pre-period coefficients.³⁰ We caution, however, that this exercise provides only a lower bound on the possible value of M . The fact that the pre-period second differences in δ are small does not on its own place a bound on how large the second differences in the post-period could have been under the counterfactual. Nonetheless, this test may be useful in diagnosing situations where the researcher-imposed bound is implausible.

²⁹Such an interval may be formed as follows: Let $\hat{\beta}^{placebo}$ be the vector of estimated event-study coefficients for the placebo groups, and $\delta^{placebo}$ its expectation. We form a lower one-sided CI for $M^{placebo} := \max_j (A^{SD} \delta^{placebo})_j$, where A^{SD} is the matrix that creates second differences. Letting $ub_j = (A^{SD} \hat{\beta}^{placebo})_j + \sigma_j z_{1-\alpha}$ and σ_j is the standard deviation of $(A^{SD} \hat{\beta}^{placebo})_j$, define $ub := \max_j ub_j$. Then $(-\infty, ub]$ is a level $(1 - \alpha)$ CI for $M^{placebo}$. Proper coverage of this CI follows from results by [Berger and Hsu \(1996\)](#) on the “union-intersection” approach.

³⁰We can reject a value of M if we reject the null hypothesis $H_0 : \mathbb{E} [A^{SD} \hat{\beta}_{pre}] \leq M$, where A^{SD} is the matrix that produces the positive and negative second differences of $\hat{\beta}_{pre}$. This is a system of linear moment inequalities.

8 Empirical Applications

We now apply our methods to two recently published papers, illustrating how empirical researchers may use these methods to conduct formal sensitivity analyses in difference-in-differences and event-study designs.

8.1 The effect of duty-to-bargain laws on long-run student outcomes

Lovenheim and Willen (2019, henceforth LW) study the impact of state-level public sector duty-to-bargain (DTB) laws, which mandated that school districts bargain in good faith with teachers' unions. LW examine the impacts of these laws on the adult labor market outcomes of people who were students around the time that these laws were passed, comparing individuals across different states and different birth cohorts to exploit the differential timing of the passage of DTB laws across states. The authors estimate the following regression specification separately for men and women, using data from the American Community Survey (ACS),

$$Y_{sct} = \sum_{r=-11}^{21} D_{scr} \beta_r + X'_{sct} \gamma + \lambda_{ct} + \phi_s + \epsilon_{sct}. \quad (25)$$

Y_{sct} is an average outcome for the cohort of students born in state s in cohort c in ACS calendar year t . D_{scr} is an indicator for whether state s passed a DTB law r years before cohort c turned age 18.³¹ The coefficients $\{\hat{\beta}_r\}$ estimate the dynamic treatment effect (or placebo effect) r years after DTB passage.³² The remaining terms include time-varying controls, birth-cohort-by-ACS-year fixed effects, and state fixed effects. Standard errors are clustered at the state level as in LW, and we normalize the coefficient β_{-2} to 0.³³ We focus on the results where the outcome is employment.

³¹ $D_{sc,-11}$ is set to 1 if state s passed a law 11 years or more after cohort c turned 18. Likewise, $D_{sc,21}$ is set to 1 if state s passed a law 21 or more years before cohort c turned 18.

³²Treatment timing in LW is staggered, and therefore the results in Sun and Abraham (2020) imply that β_r can be interpreted as a sensible weighted average of causal effects under parallel trends only if treatment effects are homogeneous across adoption cohorts. For simplicity, we focus on the robustness of the results to violations of parallel trends using the original specification in LW, which is valid under the assumption of homogeneous treatment effects. As discussed in Section 7.1, our sensitivity analysis can also be applied to estimators that are robust to treatment effect heterogeneity.

³³LW normalize event time -1 to 0, but discuss how cohorts at event time -1 may have been partially treated, since LW impute the year that a student starts school with error. Since our robust

Figure 6: Event-study coefficients $\{\beta_r\}$ for employment, estimated using the event-study specification in (25).




Figure 6 plots the estimated event-study coefficients $\{\hat{\beta}_r\}$ from specification (25). In the event-study for men (left panel), the pre-period coefficients are relatively close to zero, whereas the longer-run post-period coefficients are negative. By contrast, the results for women (right panel) suggest a downward-sloping pre-existing trend. LW therefore interpret the results for men causally, and “urge caution in lending a causal interpretation” to the results for women.

Figure 7: Sensitivity analysis for $\theta = \tau_{15}$ using $\Delta = \Delta^{SD}(M)$





Figure 7 reports results of a sensitivity analysis for the treatment effect on employ-

confidence sets assume that there is no causal effect in the pre-period ($\tau_{pre} = 0$), we instead treat event-time -2 as the reference period in our analysis.

ment for the cohort 15 years after the passage of a DTB law (as in Table 2 of LW), constructing robust confidence sets under varying assumptions on the class of possible violations of parallel trends. In blue, we plot the original OLS confidence intervals for $\hat{\beta}_{15}$ from specification (25). In red, we plot FLCIs when $\Delta = \Delta^{SD}(M)$ for different values of M . In the sensitivity analysis for men (left panel), we see that the FLCIs are similar to those from OLS when allowing for violations of parallel trends that are approximately linear ($M \approx 0$), but become wider as we allow for more non-linearity; the breakdown value for a significant effect is $M \approx 0.01$. For women (right panel), the original OLS estimates are negative and the confidence interval rules out 0. When we allow for linear violations of parallel trends ($M = 0$), however, the picture changes substantially owing to the pre-existing downward trend that is visible in Figure 6. Indeed, for $M < 0.01$ the robust confidence set contains only positive values. Thus, if we were to impose the same smoothness restrictions for men as for women, we would either have to reconcile significant effects of opposite signs by gender (if $M < 0.01$) or we would not be able to rule out null effects for both genders ($M \geq 0.1$).

Figure 8: Sensitivity analysis with and without imposing monotonicity



LW attribute the downward-sloping pre-trend in employment for women to “secular trends” affecting female labor supply, motivating us to introduce the shape restriction that δ is monotonically decreasing. Figure 8 shows how the sensitivity analysis for women changes when we incorporate this shape restriction, comparing the earlier results for $\Delta = \Delta^{SD}(M)$ to those using $\Delta = \Delta^{SDD} := \Delta^{SD}(M) \cap \{\delta : \delta_t \leq$

$\delta_{t-1}, \forall t\}$. The confidence sets imposing the shape restriction are constructed using the conditional-FLCI hybrid. We see that these shape restrictions can be informative: the lower bound of the robust confidence set over $\Delta^{SDD}(M)$ never falls substantially below the lower bound of the OLS confidence interval, even when M is large.

To evaluate the magnitudes of M , we consider the following calibration exercise: if violations of parallel trends were driven by confounding changes in education quality, what would a given value of M imply about the evolution of those confounds? Chetty, Friedman and Rockoff (2014) estimate that a 1 standard deviation increase in teacher value-added (VA) corresponds with a 0.4 percentage point increase in adult employment. Hence, a value of $M = 0.01$ would correspond with allowing the slope of the differential trend in education quality to change by one-fourtieth of a standard deviation of teacher VA across consecutive periods. Since the robust confidence sets for both men and women begin to include zero around this value of M , the strength with which we can rule out a null effect depends on our assessment of the economic plausibility of such non-linearities.


8.2 Estimating the incidence of a value-added tax cut

Benzarti and Carloni (2019, henceforth, BC) study the incidence of a decrease in the value-added tax (VAT) on restaurants in France. France reduced its VAT on sit-down restaurants from 19.6 to 5.5 percent in July of 2009. BC analyze the impact of this change using a dynamic difference-in-differences design that compares restaurants to a control group of other market services firms that were not affected by the VAT change. Their regression specification is

$$Y_{irt} = \sum_{s=2004}^{2012} \beta_s \times 1[t = s] \times D_{ir} + \phi_i + \lambda_t + \epsilon_{irt}, \quad (26)$$

where Y_{irt} is an outcome of interest for firm i in region r in year t ; D_{ir} is an indicator for whether firm i in region r is a restaurant; and ϕ_i and λ_t are firm and year fixed effects. The coefficient β_{2008} is normalized to 0. BC cluster standard errors at the regional level. BC's main finding is that the VAT reduction had a large effect on restaurant profits. Figure 9 shows the estimated event-study coefficients from specification (26) when the outcome is the log of (before-tax) restaurant profits.


Figure 9: Event-study coefficients $\{\beta_s\}$ for log profits, estimated using the event-study specification in (26).



The left panel of Figure 9 shows a sensitivity analysis for the treatment effect in 2009 using $\Delta = \Delta^{SD}(M)$. The confidence sets contain only positive values unless M exceeds 0.2, allowing us to reject a null effect on profits in 2009 if we are willing to restrict the slope of the differential trend to change by no more than 20 log points between periods. To provide additional context for the magnitudes of M , we consider benchmarks using changes in slope in the pre-period, as discussed in Section 7.3. A 95% interval for the largest change in slope using the pre-period data is $[0.1, 0.2]$. Thus, we can reject a null effect in 2009 unless we are willing to allow for changes in slope of the underlying trend that are larger than the upper bound for the largest change in slope in the pre-period.

Using context-specific knowledge, BC argue that their estimates likely understate the effect of the VAT cut on profits. Since the VAT cut occurred at the same time that a payroll subsidy for restaurants was terminated, BC write, “a conservative interpretation of our results is that we are estimating a lower bound on the effect of the VAT cut on profits” (pg. 40). This argument may be made precise by imposing that the bias of the post-period event-study coefficients is negative. The right panel of Figure 10 imposes the additional constraint that the sign of the bias be negative — that is, we set $\Delta = \Delta^{SDNB} := \Delta^{SD}(M) \cap \{\delta : \delta_{post} \leq 0\}$. With this added constraint, the robust confidence sets now rule out effects on profits smaller than 15 log points for all values of M .

Figure 10: Sensitivity analysis for $\theta = \tau_{2009}$ using $\Delta = \Delta^{SD}(M)$ and $\Delta = \Delta^{SDNB}(M)$.



9 Conclusion

This paper considers the problem of conducting inference in difference-in-differences and related designs that is robust to violations of the parallel trends assumption. We introduce a variety of possible restrictions on the class of possible differences in trends that formalize common intuitive arguments made in applied work. We then introduce inference procedures that are uniformly valid so long as the difference in trends satisfies these restrictions. We recommend that applied researchers use our proposed confidence sets to conduct formal sensitivity analyses, in which they report confidence sets for the causal effect of interest under a variety of possible restrictions on the underlying trends. Such sensitivity analyses make transparent what assumptions are needed in order to obtain informative inference and help researchers assess whether those assumptions are plausible in a given setting.

References

- Abadie, Alberto**, “Semiparametric Difference-in-Differences Estimators,” *The Review of Economic Studies*, 2005, 72 (1), 1–19.
- **and Guido W. Imbens**, “Large Sample Properties of Matching Estimators for Average Treatment Effects,” *Econometrica*, 2006, 74 (1), 235–267.

- Altonji, Joseph G., Todd E. Elder, and Christopher R. Taber**, “Selection on Observed and Unobserved Variables: Assessing the Effectiveness of Catholic Schools,” *Journal of Political Economy*, 2005, *113* (1), 151–184.
- Andrews, Donald W. K. and Panle Jia Barwick**, “Inference for Parameters Defined by Moment Inequalities: A Recommended Moment Selection Procedure,” *Econometrica*, 2012, *80* (6), 2805–2826.
- Andrews, Isaiah, Jonathan Roth, and Ariel Pakes**, “Inference for Linear Conditional Moment Inequalities,” Technical Report, NBER Working Paper No. 26374 2019.
- Armstrong, Timothy and Michal Kolesar**, “Optimal Inference in a Class of Regression Models,” *Econometrica*, 2018, *86*, 655–683.
- **and –**, “Sensitivity Analysis using Approximate Moment Condition Models,” Technical Report, arXiv preprint arXiv:1808.07387 2020.
- Ashenfelter, Orley**, “Estimating the Effect of Training Programs on Earnings,” *The Review of Economics and Statistics*, 1978, *60* (1), 47–57. Publisher: The MIT Press.
- Athey, Susan and Guido Imbens**, “Design-Based Analysis in Difference-In-Differences Settings with Staggered Adoption,” *arXiv:1808.05293 [cs, econ, math, stat]*, August 2018.
- Benzarti, Youssef and Dorian Carloni**, “Who Really Benefits from Consumption Tax Cuts? Evidence from a Large VAT Reform in France,” *American Economic Journal: Economic Policy*, February 2019, *11* (1), 38–63.
- Berger, Roger L. and Jason C. Hsu**, “Bioequivalence trials, intersection-union tests and equivalence confidence sets,” *Statistical Science*, November 1996, *11* (4), 283–319.
- Bhuller, Manudeep, Tarjei Havnes, Edwin Leuven, and Magne Mogstad**, “Broadband Internet: An Information Superhighway to Sex Crime?,” *The Review of Economics and Statistics*, 2013, *80*, 1237—1266.

- Bilinski, Alyssa and Laura A. Hatfield**, “Seeking Evidence of Absence: Reconsidering Tests of Model Assumptions,” *arXiv:1805.03273 [stat]*, May 2018.
- Bonhomme, Stephanie and Martin Weidner**, “Minimizing Sensitivity to Model Misspecification,” Technical Report 2018.
- Borusyak, Kirill and Xavier Jaravel**, “Revisiting Event Study Designs,” SSRN Scholarly Paper ID 2826228, Social Science Research Network, Rochester, NY August 2016.
- Bugni, Federico A., Ivan A. Canay, and Xiaoxia Shi**, “Inference for subvectors and other functions of partially identified parameters in moment inequality models,” *Quantitative Economics*, 2017, 8 (1), 1–38.
- Callaway, Brantly and Pedro H. C. Sant’Anna**, “Difference-in-Differences with Multiple Time Periods,” SSRN Scholarly Paper ID 3148250, Social Science Research Network, Rochester, NY March 2019.
- Chen, Xiaohong, Timothy M. Christensen, and Elie Tamer**, “Monte Carlo Confidence Sets for Identified Sets,” *Econometrica*, 2018, 86 (6), 1965–2018.
- Chernozhukov, Victor, Whitney K. Newey, and Andres Santos**, “Constrained Conditional Moment Restriction Models,” *arXiv:1509.06311 [math, stat]*, September 2015. arXiv: 1509.06311.
- Chetty, Raj, John N. Friedman, and Jonah E. Rockoff**, “Measuring the Impacts of Teachers II: Teacher Value-Added and Student Outcomes in Adulthood,” *American Economic Review*, September 2014, 104 (9), 2633–2679.
- Cho, JoonHwan and Thomas M. Russell**, “Simple Inference on Functionals of Set-Identified Parameters Defined by Linear Moments,” Technical Report, arXiv preprint arXiv:1810.03180v7 2019.
- Conley, Timothy, Christian Hansen, and Peter Rossi**, “Plausibly Exogenous,” *The Review of Economics and Statistics*, 2012, 94 (1), 260–272.
- Cox, Gregory and Xiaoxia Shi**, “A Simple Uniformly Valid Test for Inequalities,” *arXiv:1907.06317 [econ, math, stat]*, July 2019. arXiv: 1907.06317.

- de Chaisemartin, Clément and Xavier D’Haultfœuille**, “Fuzzy Differences-in-Differences,” *The Review of Economic Studies*, April 2018, *85* (2), 999–1028.
- **and** –, “Two-way fixed effects estimators with heterogeneous treatment effects,” *arXiv:1803.08807 [econ]*, March 2018. arXiv: 1803.08807.
- Dobkin, Carlos, Amy Finkelstein, Raymond Kluender, and Matthew J. Notowidigdo**, “The Economic Consequences of Hospital Admissions,” *American Economic Review*, 2018, *108* (2), 308–352.
- Donoho, David L.**, “Statistical Estimation and Optimal Recovery,” *The Annals of Statistics*, 1994, *22* (1), 238–270.
- Flynn, Zach**, “Inference Based on Continuous Linear Inequalities via Semi-Infinite Programming,” SSRN Scholarly Paper ID 3390788, Social Science Research Network, Rochester, NY May 2019.
- Frandsen, Brigham R.**, “The Effects of Collective Bargaining Rights on Public Employee Compensation: Evidence from Teachers, Firefighters, and Police,” *ILR Review*, 2016, *69* (1), 84–112.
- Freyaldenhoven, Simon, Christian Hansen, and Jesse Shapiro**, “Pre-event Trends in the Panel Event-study Design,” *American Economic Review*, 2019, *109* (9), 3307–3338.
- Gafarov, Bulat**, “Inference in high-dimensional set-identified affine models,” Technical Report, arXiv preprint arXiv:1904.00111 2019.
- Goodman-Bacon, Andrew**, “Difference-in-Differences with Variation in Treatment Timing,” Working Paper 25018, National Bureau of Economic Research September 2018.
- , “Public Insurance and Mortality: Evidence from Medicaid Implementation,” *Journal of Public Economics*, 2018, *126* (1), 216–262.
- Greenstone, Michael and Rema Hanna**, “Environmental Regulations, Air and Water Pollution, and Infant Mortality in India,” *American Economic Review*, October 2014, *104* (10), 3038–3072.

- Heckman, James, Hidehiko Ichimura, Jeffrey Smith, and Petra Todd**, “Characterizing Selection Bias Using Experimental Data,” *Econometrica*, 1998, *66* (5), 1017–1098.
- Hirano, Keisuke and Jack R. Porter**, “Impossibility Results for Nondifferentiable Functionals,” *Econometrica*, 2012, *80* (4), 1769–1790.
- Horowitz, Joel L. and Charles F. Manski**, “Identification and Robustness with Contaminated and Corrupted Data,” *Econometrica*, 1995, *63* (2), 281–302.
- Hudson, Sally, Peter Hull, and Jack Liebersohn**, “Interpreting Instrumented Difference-in-Differences,” Working Paper 2017.
- Imai, Kosuke and In Song Kim**, “On the Use of Two-way Fixed Effects Regression Models for Causal Inference with Panel Data,” 2019.
- Imbens, Guido**, “Sensitivity to Exogeneity Assumptions in Program Evaluation,” *American Economic Review*, 2003, *93* (2), 126–132.
- Joshi, VM**, “Admissibility of the usual confidence sets for the mean of a univariate or bivariate normal population,” *The Annals of Mathematical Statistics*, 1969, *40* (3), 1042–1067.
- Kadane, Joseph B.**, “Comparison of k-Class Estimators When the Disturbances Are Small,” *Econometrica*, 1971, *39* (5), 723–737.
- Kahn-Lang, Ariella and Kevin Lang**, “The Promise and Pitfalls of Differences-in-Differences: Reflections on 16 and Pregnant and Other Applications,” *Journal of Business and Economic Statistics*, 2019.
- Kaido, Hiroaki and Andres Santos**, “Asymptotically Efficient Estimation of Models Defined by Convex Moment Inequalities,” *Econometrica*, 2014, *82* (1), 387–413.
- , **Francesca Molinari, and Jörg Stoye**, “Confidence Intervals for Projections of Partially Identified Parameters,” *Econometrica*, 2019, *87*, 1397–1432.
- , – , **and –** , “Constraint Qualifications in Partial Identification,” *arXiv:1908.09103 [econ]*, August 2019. arXiv: 1908.09103.

- Keele, Luke J., Dylan S. Small, Jesse Y. Hsu, and Colin B. Fogarty**, “Patterns of Effects and Sensitivity Analysis for Differences-in-Differences,” Technical Report 2019.
- Kim, Wooyoung, Koohyun Kwon, Soonwoo Kwon, and Sokbae Lee**, “The identification power of smoothness assumptions in models with counterfactual outcomes,” *Quantitative Economics*, 2018, *9* (2), 617–642.
- Kline, Patrick and Andres Santos**, “Sensitivity to missing data assumptions: Theory and an evaluation of the U.S. wage structure,” *Quantitative Economics*, 2013, *4* (2), 231–267.
- Kolesar, Michal and Christoph Rothe**, “Inference in Regression Discontinuity Designs with a Discrete Running Variable,” *American Economic Review*, 2018, *108* (8), 2277–2304.
- Kropko, Jonathan and Robert Kubinec**, “Why the Two-Way Fixed Effects Model Is Difficult to Interpret, and What to Do About It,” SSRN Scholarly Paper ID 3062619, Social Science Research Network, Rochester, NY May 2018.
- Lee, Jason D., Dennis L. Sun, Yuekai Sun, and Jonathan E. Taylor**, “Exact Post-Selection Inference, with Application to the Lasso,” *The Annals of Statistics*, June 2016, *44* (3), 907–927.
- Lee, Jin Young and Gary Solon**, “The Fragility of Estimated Effects of Unilateral Divorce Laws on Divorce Rates,” *The B.E. Journal of Economic Analysis & Policy*, 2011, *11* (1).
- Lovenheim, Michael F. and Alexander Willen**, “The Long-Run Effects of Teacher Collective Bargaining,” *American Economic Journal: Economic Policy*, 2019, *11* (3), 292–324.
- Low, Mark G.**, “Bias-Variance Tradeoffs in Functional Estimation Problems,” *The Annals of Statistics*, 1995, *23* (3), 824–835.
- Malani, Anup and Julian Reif**, “Interpreting pre-trends as anticipation: Impact on estimated treatment effects from tort reform,” *Journal of Public Economics*, April 2015, *124*, 1–17.

- Manski, Charles F. and John V. Pepper**, “How Do Right-to-Carry Laws Affect Crime Rates? Coping with Ambiguity Using Bounded-Variation Assumptions,” *Review of Economics and Statistics*, 2018, *100* (2), 232–244.
- Masten, Matthew A. and Alexandre Poirier**, “Identification of Treatment Effects Under Conditional Partial Independence,” *Econometrica*, 2018, *86* (1), 317–351.
- and –, “Inference on Breakdown Frontiers,” *Quantitative Economics*, 2020, *11* (1), 41–111.
- Meer, Jonathan and Jeremy West**, “Effects of the Minimum Wage on Employment Dynamics,” *Journal of Human Resources*, 2016, *51* (2), 500–522.
- Moreira, Marcelo J.**, “A Conditional Likelihood Ratio Test for Structural Models,” *Econometrica*, 2003, *71* (4), 1027–1048. Publisher: [Wiley, Econometric Society].
- and **Geert Ridder**, “Efficiency Loss of Asymptotically Efficient Tests in an Instrumental Variables Regression,” SSRN Scholarly Paper ID 3348716, Social Science Research Network, Rochester, NY March 2019.
- Noack, Claudia and Christoph Rothe**, “Bias-aware inference in fuzzy regression discontinuity designs,” *arXiv preprint arXiv:1906.04631*, 2019.
- Oster, Emily**, “Unobservable Selection and Coefficient Stability: Theory and Evidence,” *Journal of Business & Economic Statistics*, 2019, *37* (2), 187–204.
- Rambachan, Ashesh and Jonathan Roth**, “Design-Based Uncertainty for Quasi-Experiments,” *arXiv:2008.00602 [econ, stat]*, August 2020. arXiv: 2008.00602.
- Romano, Joseph P. and Azeem M. Shaikh**, “Inference for identifiable parameters in partially identified econometric models,” *Journal of Statistical Planning and Inference*, September 2008, *138* (9), 2786–2807.
- , –, and **Michael Wolf**, “A Practical Two-Step Method for Testing Moment Inequalities,” *Econometrica*, 2014, *82* (5), 1979–2002.
- Rosenbaum, Paul**, “Sensitivity Analysis in Observational Studies,” in B. S. Everitt and D. C. Howell, eds., *Encyclopedia of Statistics in Behavioral Science*, 2005.

- Roth, Jonathan**, “Pre-test with Caution: Event-study Estimates After Testing for Parallel Trends,” *Working paper*, 2019.
- Sant’Anna, Pedro H. C. and Jun B. Zhao**, “Doubly Robust Difference-in-Differences Estimators,” *Journal of Econometrics*, 2020, *Forthcoming*.
- Schrijver, Alexander**, *Theory of Linear and Integer Programming*, Wiley-Interscience, 1986.
- Sun, Liyan and Sarah Abraham**, “Estimating Dynamic Treatment Effects in Event Studies with Heterogeneous Treatment Effects,” *Working Paper*, 2020.
- Słoczyński, Tymon**, “A General Weighted Average Representation of the Ordinary and Two-Stage Least Squares Estimands,” 2018.
- Wolfers, Justin**, “Did Unilateral Divorce Laws Raise Divorce Rates? A Reconciliation and New Results,” *American Economic Review*, 2006, *96*, 1802–1820.
- Ye, Ting, Luke Keele, Raiden Hasegawa, and Dylan S. Small**, “A Negative Correlation Strategy for Bracketing in Difference-in-Differences with Application to the Effect of Voter Identification Laws on Voter Turnout,” June 2020.

An Honest Approach to Parallel Trends

Online Appendix

Ashesh Rambachan Jonathan Roth

September 10, 2020

This online appendix contains proofs and additional results for the paper “An Honest Approach to Parallel Trends” by Ashesh Rambachan and Jonathan Roth. Section [A](#) collects together additional results that are referenced in the main text. Section [B](#) contains proofs and auxiliary lemmas for the results in the main text. The supplementary materials provide statements and proofs of uniform asymptotic results along with additional simulation results.

A Additional Results

A.1 Optimal local asymptotic power of FLCIs

As discussed in Remark [6](#), the FLCIs have local asymptotic power converging to the power envelope provided that Assumption [4](#) and Assumption [5](#) are satisfied. We now formally state this result. The proof is given in Section [B](#).

Proposition A.1. *Fix $\delta_A \in \Delta$, $\tau_A \in \mathbb{R}^{\bar{T}}$ and suppose Σ^* is positive definite. Let $\theta_A^{ub} = \sup_{\theta} \mathcal{S}(\Delta, \delta_A + \tau_A)$ be the upper bound of the identified set. Suppose that Assumption [5](#) holds and $\delta_{A,pre}$ satisfies Assumption [4](#). Then, for any $x > 0$ and $\alpha \in (0, 0.5]$,*

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta_A^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n}^{FLCI} \right) = \lim_{n \rightarrow \infty} \sup_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_{\alpha}(\Delta, \Sigma_n)} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta_A^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n} \right).$$

The analogous result holds replacing $\theta_A^{ub} + \frac{1}{\sqrt{n}}x$ with $\theta_A^{lb} - \frac{1}{\sqrt{n}}x$, for θ_A^{lb} the lower bound of the identified set.

Thus, $\mathcal{C}_{\alpha, n}^{FLCI}$ behaves similarly to $\mathcal{C}_{\alpha, n}^C$ as $n \rightarrow \infty$ when Assumption [4](#) and Assumption [5](#) hold.

A.2 Connection to linear independence constraint qualification (LICQ)

In this section, we draw connections between linear independence constraint qualification (LICQ) and Assumption 5, under which the power of the conditional test converges to the power envelope asymptotically. We show that LICQ implies Assumption 5. We follow the notation provided in [Kaido et al. \(2019\)](#).

Suppose $\Delta = \{\delta : A\delta \leq d\}$. Let $m(\tau_{post}; \beta) = A(\beta - M_{post}\tau_{post}) - d$, and let $T(\Delta, \beta) := \{\tau_{post} : m(\tau_{post}; \beta) \leq 0\}$ be the identified set for the full parameter vector τ_{post} . Define the set of support points in direction p to be $S(p, T) := \{\tau_{post} : p'\tau_{post} = \sup_{\tau_{post} \in T} p'\tau_{post}\}$.

Definition 1. The **linear constraint constraint qualification** (LICQ) is satisfied in the direction p if, for all support points in the direction p , the gradients of the binding constraints are linearly independent. That is, for all $\tau_{post} \in S(p, T)$, the set $\{D_{\tau_{post}} m_j(\tau_{post}, \beta) : m_j(\tau_{post}, \beta) = 0\}$ is linearly independent, where $D_{\tau_{post}}$ denotes the gradient with respect to τ_{post} .

LICQ in the directions l and $-l$ is equivalent to restrictions on the binding moments in the problems b^{min} and b^{max} .

Lemma A.1. *Suppose $\beta_A = \delta_A + M_{post}\tau_{A,post}$ for some $\delta_A \in \Delta = \{\delta : A\delta \leq d\}$ and $\tau_{A,post} \in \mathbb{R}^T$. Then the following are equivalent: (i) LICQ is satisfied in the direction l ; (ii) For any solution δ^{**} to the linear program*

$$b^{min}(\delta_{A,pre}) = \min_{\delta} l'\delta_{post} \text{ s.t. } A\delta \leq d, \delta_{pre} = \delta_{A,pre},$$

*the matrix $A_{(B(\delta^{**}), post)}$ with rows corresponding with the binding inequality constraints at δ^{**} has rank $|B(\delta^{**})|$. Analogous results hold replacing l with $-l$ in (i) and min with max in (ii).*

Proof. We first show (i) implies (ii). Let δ^{**} be a solution to the minimization problem for b^{min} . Let $\tau_{post}^{**} = \beta_{A,post} - \delta_{post}^{**}$. Observe that $l'\tau_{post}^{**} = l'\beta_{A,post} - b^{min}(\delta_{A,post})$. From (6), we then see that $l'\tau_{post}^{**} = \theta^{ub}$ and hence $\tau_{post}^{**} \in S(l, T)$. Now, note that by construction, $m(\beta_A, \tau_{post}^{**}) = A(\beta_A - M_{post}\tau_{post}^{**}) - d = A\delta^{**} - d$, so the binding constraints in $m(\beta_A, \tau_{post}^{**})$ correspond with the binding constraints in the minimization

for b^{min} . Finally, observe that $D_{\tau_{post}} m(\beta_A, \tau_{post}^{**}) = A_{(\cdot, post)}$. It then follows from (i) that the rows of $A_{(B(\delta^{**}), post)}$ are linearly independent, which gives the desired result.

Conversely, suppose $\tau_{post}^{**} \in S(l, \tau)$. By definition, there exists some $\delta^{**} \in \Delta$ such that $\delta^{**} = \beta_A - M_{post} \tau_{post}^{**}$, and $l' \tau_{post}^{**} = \theta^{ub}$. Thus, $\theta^{ub} = l' \beta_{A, post} - l' \delta^{**}$. It then follows from (6) that $l' \delta^{**} = b^{min}(\delta_{A, pre})$, so δ^{**} is a solution to the optimization b^{min} . (ii) then implies that $A_{(B(\delta^{**}), post)}$ has linearly independent rows. By the same argument as earlier in the proof, $A_{(B(\delta^{**}), post)}$ corresponds with the matrix of gradients for the binding constraints in $m(\beta_A, \tau_{post}^{**})$, from which we see that LICQ is satisfied. \square

Therefore, if LICQ holds in the directions l and $-l$, then Assumption 5 is satisfied. Indeed, if LICQ holds, then Lemma 5 implies that the rank condition in Assumption 5 holds for *any* solutions δ^* and δ^{**} to the problems b^{max} and b^{min} . By contrast, Assumption 5 only requires the rank condition to hold for *at least one* solution to b^{max} and b^{min} .

It is possible for a linear program to have multiple solutions, and for the rows of the binding constraints to be linearly independent (non-degenerate) for some solutions but not for others (e.g., see Example 1 on p. 146 of Sierksma (2001)). Assumption 5 is thus potentially weaker than LICQ if there are multiple solutions to the problem for b^{max} or b^{min} .

A.3 Details on the conditional-FLCI hybrid Confidence Sets

We provide further technical details on the construction of conditional-FLCI hybrid confidence sets discussed in Section 5. To construct the conditional-FLCI hybrid confidence sets, we first provide a lemma that implies that the affine estimator at which the optimal FLCI is centered can be written as an affine function of $A\hat{\beta}$, where recall that A is the matrix defining the polyhedral set, $\Delta = \{\delta : A\delta \leq d\}$.

Lemma A.2. *Suppose $\Delta = \{\delta : A\delta \leq d\} \neq \emptyset$, and (a, v) are such that $\bar{b}(a, v) < \infty$. Then, there exists \tilde{v} such that $v' = \tilde{v}' A$.*

Proof. Note that

$$\bar{b}(a, v) = \max_{\delta, \tau_{post}} |v'(\delta + M_{post} \tau_{post}) - l' \tau| \text{ s.t. } A\delta - d \leq 0.$$

We will show that if $\bar{b}(a, v)$ is finite, then for all $\tilde{\delta} \in \Delta$, $A\tilde{\delta} = 0$ implies $v'\tilde{\delta} = 0$. This implies that $\tilde{\delta}$ is in the rowspace of A , from which the result follows. To prove this,

suppose towards contradiction that $\tilde{\delta}$ is such that $A\tilde{\delta} = 0$ and $v'\tilde{\delta} \neq 0$. Since $A\tilde{\delta} = 0$, it follows that $\delta_c := (\delta_0 + c \cdot \tilde{\delta})$ is in Δ for any $\delta_0 \in \Delta$ and $c \in \mathbb{R}$. However, it then follows that for any fixed τ_{post} and δ_0 , the objective in the previous display can be made arbitrarily large at (δ_c, τ_{post}) by $c \rightarrow \infty$. \square

Consider the level $1 - \kappa$ optimal FLCI, $\mathcal{C}_{\kappa,n}^{FLCI} = a_n + v'_n \hat{\beta}_n \pm \chi_n$. By Lemma A.2, there exists some vector \tilde{v}_n such that the level $1 - \kappa$ optimal FLCI can be written as $a_n + \tilde{v}'_n A \hat{\beta}_n \pm \chi_n$. Since $\tilde{Y}_n(\theta) = A \hat{\beta}_n - \tilde{A}_{(\cdot,1)} \theta - d$, it follows that $\bar{\theta} \in \mathcal{C}_{\kappa,n}^{FLCI}$ if and only if

$$\begin{aligned} \tilde{v}'_n \tilde{Y}_n(\bar{\theta}) &\leq \chi_n - a_n - \tilde{v}'_n d + (1 - \tilde{v}'_n \tilde{A}_{(\cdot,1)}) \bar{\theta}, \\ -\tilde{v}'_n \tilde{Y}_n(\bar{\theta}) &\leq \chi_n + a_n + \tilde{v}'_n d - (1 - \tilde{v}'_n \tilde{A}_{(\cdot,1)}) \bar{\theta}. \end{aligned}$$

One can further show that $(1 - \tilde{v}'_n \tilde{A}_{(\cdot,1)}) \bar{\theta} = 0$, which simplifies the upper bounds above.³⁴ Defining the matrix $\tilde{V}_n = (\tilde{v}'_n, -\tilde{v}'_n)'$ and the vector $d_n(\bar{\theta})$ which stacks the upper-bounds of the inequalities in the previous display, we see that the optimal level $1 - \kappa$ FLCI contains the parameter value $\bar{\theta}$ if and only if $\tilde{V}_n \tilde{Y}_n(\bar{\theta}) \leq \bar{d}_n(\bar{\theta})$.

With this equivalent representation of the optimal FLCI, we now characterize the distribution of the test statistic $\hat{\eta}$ (17) conditional on the parameter θ falling within the optimal FLCI.

Lemma A.3. $\hat{\eta} \mid \left\{ \gamma_* \in \hat{V}_n, S_n = s, \bar{\theta} \in \mathcal{C}_{\kappa,n}^{FLCI} \right\} \sim \xi \mid \xi \in [v_{C-FLCI}^{lo}, v_{C-FLCI}^{up}]$, where $\xi \sim \mathcal{N}\left(\gamma'_* \tilde{\mu}, \gamma'_* \tilde{\Sigma}_n \gamma_*\right)$, $v_{C-FLCI}^{lo} := \max\{v^{lo}, v_{FLCI}^{lo}\}$, $v_{C-FLCI}^{up} := \min\{v^{up}, v_{FLCI}^{up}\}$; v^{lo} and v^{up} are as defined in Section 4.2, $v_{FLCI}^{lo} := \max_{\{j: (\tilde{V}_n c_{n,\gamma_*})_j < 0\}} \frac{\bar{d}_n(\theta)_j - (\tilde{V}_n S_n)_j}{(\tilde{V}_n c_{n,\gamma_*})_j}$, $v_{FLCI}^{up} := \min_{\{j: (\tilde{V}_n c_{n,\gamma_*})_j > 0\}} \frac{\bar{d}_n(\theta)_j - (\tilde{V}_n S_n)_j}{(\tilde{V}_n c_{n,\gamma_*})_j}$, $c_{n,\gamma_*} = \frac{\tilde{\Sigma}_n \gamma_*}{\gamma'_* \tilde{\Sigma}_n \gamma_*}$ and $S_n = (I - \frac{\tilde{\Sigma}_n \gamma_*}{\gamma'_* \tilde{\Sigma}_n \gamma_*} \gamma'_*) \tilde{Y}_n(\bar{\theta})$, and γ_* is the vector of Lagrange multipliers for the primal problem (18).

Proof. The proof follows an analogous argument to Lemma 9 in ARP. Recall that conditional on $\gamma_* \in \hat{V}_n$, $\hat{\eta} = \gamma'_* \tilde{Y}_n$. Recall also that $\theta \in \mathcal{C}_{\kappa,n}^{FLCI}$ if and only if $\tilde{V}_n \tilde{Y}_n \leq \bar{d}_n$. Observe that the set of values of \tilde{Y}_n such that

$$\gamma'_* \tilde{Y}_n = \left(\max \gamma'_* \tilde{Y}_n \text{ s.t. } \gamma \geq 0, \gamma' \tilde{A}_{(\cdot,-1)} = 0, \gamma' \tilde{\sigma}_n = 1 \right) \text{ and } \tilde{V}_n \tilde{Y}_n \leq \bar{d}_n$$

³⁴Applying the definitions of \tilde{A} and \tilde{v}_n , we obtain that $\tilde{v}_n \tilde{A}_{(\cdot,1)} = v_{n,post} \Gamma^{-1} e_1$. However, we show in the proof to Lemma B.19 that $v_{n,post} = l'$, so $\tilde{v}'_n \tilde{A}_{(\cdot,1)} = l' \Gamma^{-1} e_1$. The result then follows from the fact that $e'_1 \Gamma = l'$ by construction.

is convex. This follows from the fact that if the expression above holds for both \tilde{Y}_n and \tilde{Y}_n^* , then $\gamma'_* \tilde{Y}_n \geq \gamma' \tilde{Y}_n$ and $\gamma'_* \tilde{Y}_n^* \geq \gamma' \tilde{Y}_n^*$ for all γ feasible in the maximization. It then follows that $\gamma'_*(\alpha \tilde{Y}_n^* + (1 - \alpha) \tilde{Y}_n) \geq \gamma'(\alpha \tilde{Y}_n^* + (1 - \alpha) \tilde{Y}_n)$ for any $\alpha \in (0, 1)$. Thus, $(\alpha \tilde{Y}_n^* + (1 - \alpha) \tilde{Y}_n)$ is also equal to the maximum. It is likewise clear that the second constraint holds for a convex combination of \tilde{Y}_n and \tilde{Y}_n^* .

Thus, once we condition on S_n , the set of values of $\gamma'_* \tilde{Y}_n$ such that $\gamma_* \in \hat{V}_n$ and $\tilde{V}_n \tilde{Y}_n \leq \bar{d}_n$ is an interval. It thus suffices to find the endpoints. Without loss of generality, we focus on the lower bound. For ease of notation, let $F := \{\gamma \geq 0, \gamma' \tilde{A}_{(\cdot, -1)} = 0, \gamma' \tilde{\sigma}_n = 1\}$ denote the feasible region for the maximization. Then we are interested in

$$\min_{\{\tilde{Y}_n : S_n = s\}} \left\{ \gamma'_* \tilde{Y}_n : \gamma'_* \tilde{Y}_n = \max_{\gamma \in F} \gamma' \tilde{Y}_n, \tilde{V}_n \tilde{Y}_n \leq \bar{d}_n \right\}.$$

Recalling that $S_n = (I - c_{n, \gamma_*} \gamma'_*) \tilde{Y}_n$ for $c_{n, \gamma_*} = \frac{\tilde{\Sigma}_n \gamma_*}{\gamma'_* \tilde{\Sigma}_n \gamma_*}$, the expression becomes

$$\min_{\{\tilde{Y}_n : S_n = s\}} \left\{ \gamma'_* \tilde{Y}_n : \gamma'_* \tilde{Y}_n = \max_{\gamma \in F} \gamma' (s + c_{n, \gamma_*} \gamma'_* \tilde{Y}_n), \tilde{V}_n \tilde{Y}_n \leq \bar{d}_n \right\},$$

which is equivalent to

$$\min \left\{ \left\{ x : x = \max_{\gamma \in F} \gamma' (s + c_{n, \gamma_*} x) \right\} \cap \left\{ \gamma'_* \tilde{Y}_n : \tilde{Y}_n \text{ s.t. } S_n = s, \tilde{V}_n \tilde{Y}_n \leq \bar{d}_n \right\} \right\}.$$

However, the first set in the minimization above is the interval $[v^{lo}, v^{up}]$, and the polyhedral lemma in [Lee et al. \(2016\)](#) (Lemma 5.1) implies that second set is the interval $[v_{FLCI}^{lo}, v_{FLCI}^{up}]$. Thus, the expression above is $\max\{v^{lo}, v_{FLCI}^{lo}\}$, as desired. The argument for the lower bound of the interval is analogous. Finally, the independence of $\gamma'_* \tilde{Y}_n$ and S_n implies that the distribution of $\gamma'_* \tilde{Y}_n$ conditional on $\{\gamma_* \in \hat{V}_n, S_n = s, \bar{\theta} \in \mathcal{C}_{n, \kappa}^{FLCI}\}$ is truncated normal. \square

We now show that the conditional-FLCI hybrid confidence sets have asymptotic properties similar to those of the conditional test. The proofs of these results are provided in [Section B](#).

Proposition A.2. *The conditional-FLCI hybrid test is consistent. For any $\delta_A \in$*

$\Delta, \tau_A \in \mathbb{R}^{\bar{T}}, \theta_{out} \notin \mathcal{S}(\Delta, \delta_A + \tau_A), \alpha \in (0, .5],$ and $\kappa \in (0, \alpha),$

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} (\theta_{out} \notin \mathcal{C}_{\kappa, \alpha, n}^{C-FLCI}) = 1.$$

Proposition A.3. Fix $\delta_A \in \Delta, \tau_A,$ and Σ^* positive definite. Suppose Assumption 5 holds. Suppose $\alpha \in (0, .5], \kappa \in (0, \alpha),$ and let $\tilde{\alpha} = \frac{\alpha - \kappa}{1 - \kappa}.$ Then,

$$\liminf_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left((\theta_A^{ub} + \frac{1}{\sqrt{n}}x) \notin \mathcal{C}_{\kappa, \alpha, n}^{C-FLCI} \right) \geq \lim_{n \rightarrow \infty} \sup_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_{\tilde{\alpha}}(\Delta, \Sigma_n)} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left((\theta_A^{ub} + \frac{1}{\sqrt{n}}x) \notin \mathcal{C}_{\alpha, n} \right).$$

The analogous result holds replacing $\theta_A^{ub} + \frac{1}{\sqrt{n}}x$ with $\theta_A^{lb} - \frac{1}{\sqrt{n}}x,$ for θ_A^{lb} the lower bound of the identified set (although the constant c^* may differ).

B Proofs of Main Finite Sample Normal Results

B.1 Proofs of Main Finite Sample Normal Results

Throughout the proofs, we will use the following notation. Let $\tilde{Y}(\hat{\beta}_n, A, d, \bar{\theta}) := A\hat{\beta}_n - d - \tilde{A}_{(\cdot, 1)}\bar{\theta}.$ Define $\psi_\alpha^C(\hat{\beta}_n; A, d, \bar{\theta}, \Sigma) := \psi_\alpha^C(\tilde{Y}(\hat{\beta}_n; A, d, \bar{\theta}), A\Sigma A')$ to be an indicator for whether the conditional test constructed using $(\hat{\beta}_n, A, d, \bar{\theta}, \Sigma)$ rejects.

Proof of Proposition 3.2

Proof. First, suppose Assumption 4 holds. Without loss of generality, we show $\mathbb{P}((\theta^{ub} + x) \in \mathcal{C}_{\alpha, n}^{FLCI}) \rightarrow 0$ for any $x > 0.$ By Lemma B.20 there exists (\bar{a}, \bar{v}) such that $\bar{b}(\bar{a}, \bar{v}) = \frac{1}{2}LID(\Delta, \delta_{pre}) =: \bar{b}_{min}$ and $\mathbb{E}_{(\delta_A, \tau_A, \Sigma_n)} \left[\bar{a} + \bar{v}'\hat{\beta}_n \right] = \frac{1}{2}(\theta^{ub} + \theta^{lb}) =: \theta^{mid}.$ Let $\bar{\mathcal{C}}_n := \bar{a} + \bar{v}'\hat{\beta}_n \pm \chi_n(\bar{a}, \bar{v})$ denote the fixed length confidence interval based on $(\bar{a}, \bar{v}).$

By construction, $\bar{\chi}_n := \chi_n(\bar{a}, \bar{v})$ is the $1 - \alpha$ quantile of the $|\mathcal{N}(\bar{b}_{min}, \sigma_{\bar{v}, n}^2)|$ distribution. Since $\sigma_{\bar{v}, n}^2 = \frac{1}{n}\sigma_{\bar{v}, 1}^2 \rightarrow 0,$ the $|\mathcal{N}(\bar{b}_{min}, \sigma_{\bar{v}, n}^2)|$ distribution collapses to a point mass at $\bar{b}_{min},$ and thus $\bar{\chi}_n \rightarrow \bar{b}_{min}.$ By construction the length of the shortest FLCI $\chi_n := \chi_n(a_n, v_n)$ must be less than or equal to $\bar{\chi}_n,$ and so $\limsup_{n \rightarrow \infty} \chi_n \leq \bar{b}_{min}.$ Let $b_n := \bar{b}(a_n, v_n)$ be the worst-case bias of the optimal FLCI. Since $\alpha \in (0, .5],$ Lemma B.21 implies that $\chi_n \geq b_n.$ Additionally, Lemma B.19 implies that $b_n \geq \frac{1}{2}LID(\Delta, \delta_{pre}) = \bar{b}_{min},$ and thus $\chi_n \geq \bar{b}_{min}.$ Hence, $\chi_n \rightarrow \bar{b}_{min}$ implies $b_n \rightarrow \bar{b}_{min}.$ Additionally, note that for $\alpha \in (0, 0.5],$ $\chi_n(a, v)$ is increasing in both $\bar{b}(a, v)$ and $\sigma_{v, n}.$

Since $\bar{b}_{min} \leq b_n$ and $\chi_n \leq \bar{\chi}_n$, it must be that $\sigma_{v_n,n} \leq \sigma_{\bar{v},n}$, from which it follows that $\sigma_{v_n,n} \rightarrow 0$.

Now, we claim that $\mu_n := \mathbb{E}_{(\delta_A, \tau_A, \Sigma_n)} \left[a_n + v'_n \hat{\beta}_n \right]$ converges to $\theta^{mid} := \frac{1}{2}(\theta^{ub} + \theta^{lb})$. To show this, note that $\mu_n = a_n + v'_n \beta_A$ for $\beta_A = \delta_A + \tau_A$. Since $\theta^{ub}, \theta^{lb} \in \mathcal{S}(\Delta, \beta_A)$, by the definition of the identified set there exist $\delta^{ub}, \delta^{lb} \in \Delta$ and τ^{ub}, τ^{lb} such that $\beta_A = \delta^{ub} + \tau^{ub} = \delta^{lb} + \tau^{lb}$, $\theta^{ub} = l'_{\tau^{ub}}$ and $\theta^{lb} = l'_{\tau^{lb}}$. Thus, $\theta^{ub} - \mathbb{E}_{(\delta^{ub}, \tau^{ub}, \Sigma_n)} \left[a_n + v'_n \hat{\beta}_n \right] = \theta^{ub} - \mu_n$ and $\mathbb{E}_{(\delta^{lb}, \tau^{lb}, \Sigma_n)} \left[a_n + v'_n \hat{\beta}_n \right] - \theta^{lb} = \mu_n - \theta^{lb}$. This implies that $b_n \geq \max\{\theta^{ub} - \mu_n, \mu_n - \theta^{lb}\} = \bar{b}_{min} + |\mu_n - \theta^{mid}|$, where the equality uses the fact that $\theta^{ub} - \theta^{lb} = LID(\Delta, \delta_{A,pre}) = 2\bar{b}_{min}$. Since we've shown that $b_n \rightarrow \bar{b}_{min}$, it follows that $\mu_n \rightarrow \theta^{mid}$, as desired.

Next, note that if $\hat{\beta}_n \sim \mathcal{N}(\delta_A + \tau_A, \Sigma_n)$, then $a_n + v'_n \hat{\beta}_n \sim \mathcal{N}(\mu_n, \sigma_{v_n,n}^2)$. Observe that $\bar{\theta} \in \mathcal{C}_{\alpha,n}^{FLCI}$ if and only if $a_n + v'_n \hat{\beta}_n \in [\bar{\theta} - \chi_n, \bar{\theta} + \chi_n]$. Thus,

$$\mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} (\bar{\theta} \in \mathcal{C}_{\alpha,n}^{FLCI}) = \Phi \left(\frac{\bar{\theta} + \chi_n - \mu_n}{\sigma_{v_n,n}} \right) - \Phi \left(\frac{\bar{\theta} - \chi_n - \mu_n}{\sigma_{v_n,n}} \right).$$

Now, recalling that $\theta^{ub} = \theta^{mid} + \bar{b}_{min}$ by construction, we have $\mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} ((\theta^{ub} + x) \in \mathcal{C}_{\alpha,n}^{FLCI})$ equals

$$\Phi \left(\frac{\theta^{mid} + \bar{b}_{min} + x + \chi_n - \mu_n}{\sigma_{v_n,n}} \right) - \Phi \left(\frac{\theta^{mid} + \bar{b}_{min} + x - \chi_n - \mu_n}{\sigma_{v_n,n}} \right). \quad (27)$$

Note that the term inside the second normal CDF in the previous display equals

$$-\frac{\chi_n - b_n}{\sigma_{v_n,n}} + \frac{x + \theta^{mid} - \mu_n + \bar{b}_{min} - b_n}{\sigma_{v_n,n}}.$$

However, the first summand above is bounded between $-z_{1-\alpha/2}$ and $-z_{1-\alpha}$ by Lemma B.21. Additionally, we've shown that the $\theta^{mid} - \mu_n \rightarrow 0$ and $\bar{b}_{min} - b_n \rightarrow 0$, so the numerator of the second summand converges to $x > 0$. Since the denominator $\sigma_{v_n,n} \rightarrow 0$, the expression in the previous display diverges to ∞ , and hence the second normal CDF term in (27) converges to 1, which implies that $\mathbb{P}((\theta^{ub} + x) \in \mathcal{C}_{\alpha,n}^{FLCI}) \rightarrow 0$, as needed.

Conversely, suppose Assumption 4 fails. Let $L_A := LID(\Delta, \delta_{A,pre})$ and $\bar{L} := \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre})$. By Lemma B.19, $b_n := \bar{b}(a_n, v_n) \geq \frac{1}{2}\bar{L} =: \bar{b}_{min}$. As argued earlier in the proof, since $\alpha \in (0, .5]$, $\chi_n \geq b_n = \frac{1}{2}\bar{L}$. If $\bar{L} = \infty$, then $\mathcal{C}_{\alpha,n}^{FLCI}$ is the real line, and thus never rejects, so $\mathcal{C}_{\alpha,n}^{FLCI}$ is trivially inconsistent under the assumption

that $\mathcal{S}_\theta(\Delta, \delta_A + \tau_A) \neq \mathbb{R}$. For the remainder of the proof, we assume $L_A < \bar{L} < \infty$. From Lemma 2.1, $\mathcal{S}(\delta_A + \tau_A, \Delta) = [\theta^{lb}, \theta^{ub}]$, where $\theta^{ub} - \theta^{lb} = LID(\Delta, \delta_{A,pre}) = L_A$. Let $\epsilon = \frac{1}{4}(\bar{L} - L_A)$, and set $\theta_1^{out} := \theta^{ub} + \epsilon$ and $\theta_2^{out} := \theta^{lb} - \epsilon$. Let $\theta^{mid} = \frac{1}{2}(\theta^{ub} + \theta^{lb})$ be the midpoint of the identified set. By construction, $\theta_1^{out} - \theta^{mid} = \theta^{mid} - \theta_2^{out} = \frac{1}{2}L_A + \epsilon < \frac{1}{2}\bar{L}$. Since $\mathcal{C}_{\alpha,n}^{FLCI}$ is an interval with half-length at least $\frac{1}{2}\bar{L}$, it follows that if $\theta^{mid} \in \mathcal{C}_{\alpha,n}^{FLCI}$ then at least one of $\theta_1^{out}, \theta_2^{out}$ is also in $\mathcal{C}_{\alpha,n}^{FLCI}$. Hence, $\mathbb{P}(\theta_1^{out} \in \mathcal{C}_{\alpha,n}^{FLCI}) + \mathbb{P}(\theta_2^{out} \in \mathcal{C}_{\alpha,n}^{FLCI}) \geq \mathbb{P}(\theta^{mid} \in \mathcal{C}_{\alpha,n}^{FLCI}) \geq 1 - \alpha$, where the final bound follows since $\mathcal{C}_{\alpha,n}^{FLCI}$ controls size. It follows that $\limsup_{n \rightarrow \infty} \mathbb{P}(\theta_j^{out} \in \mathcal{C}_{\alpha,n}^{FLCI}) \geq \frac{1}{2}(1 - \alpha) > 0$ for at least one $j \in \{1, 2\}$. \square

Proof of Proposition 4.1

Proof. Lemma 2.1 showed that the identified set is an interval, $\mathcal{S}(\delta_A + \tau_A, \Delta) = [\theta^{lb}, \theta^{ub}]$, and so if $\theta^{out} \notin \mathcal{S}(\delta_A + \tau_A, \Delta)$, then we must have either $\theta^{out} = \theta^{ub} + x$ or $\theta^{out} = \theta^{lb} - x$ for some $x > 0$. Without loss of generality, consider the case $\theta^{out} = \theta^{ub} + x$, so

$$\liminf_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)}(\theta^{out} \notin \mathcal{C}_{\alpha,n}^C) = \liminf_{n \rightarrow \infty} \mathbb{E}_{(\delta_A, \tau_A, \Sigma_n)} \left[\psi_\alpha^C(\hat{\beta}_n; A, d, \theta^{ub} + x, \Sigma_n) \right].$$

Lemma B.2 along with $\Sigma_n = \frac{1}{n}\Sigma^*$ imply $\psi_\alpha^C(\hat{\beta}_n; A, d, \theta^{ub} + x, \Sigma_n) = \psi_\alpha^C(\sqrt{n}\hat{\beta}_n; A, \sqrt{nd}, \sqrt{n}\theta^{ub} + \sqrt{nx}, \Sigma^*)$. Thus,

$$\liminf_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)}(\theta^{out} \notin \mathcal{C}_{\alpha,n}^C) = \liminf_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^C(\hat{\beta}_n; A, \sqrt{nd}, \sqrt{n}\theta^{ub} + \sqrt{nx}, \Sigma^*) \right],$$

where we further used that $\hat{\beta}_n \sim \mathcal{N}(\beta_A, \Sigma_n)$ implies $\sqrt{n}\hat{\beta}_n \sim \mathcal{N}(\sqrt{n}\beta_A, \Sigma^*)$. Lemma B.1 implies that $\sqrt{n}\theta^{ub} = \theta_n^{ub}$, for $\theta_n^{ub} = \sup \mathcal{S}(\sqrt{n}\delta_A + \sqrt{n}\tau_A, \Delta_n)$ and $\Delta_n = \{\delta : A\delta \leq \sqrt{nd}\}$. It follows from Lemma B.18 that

$$\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^C(\hat{\beta}_n; A, \sqrt{nd}, \sqrt{n}\theta^{ub} + \sqrt{nx}, \Sigma^*) \right] \geq \rho_{LB}(x, \Sigma^*),$$

for ρ_{LB} a function with $\lim_{x \rightarrow \infty} \rho_{LB}(x, \Sigma^*) = 1$. It is immediate from the previous two displays that $\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)}(\theta^{out} \notin \mathcal{C}_{\alpha,n}^C) = 1$ as desired. \square

Proof of Proposition 4.2

Proof. We show that each of the limits in the proposition equals $\Phi(c^*x - z_{1-\alpha})$. Following the same steps as proof of Proposition 4.1, the first limit of interest can be written as

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n}^C \right) = \lim_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^C(\hat{\beta}_n; A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) \right].$$

The term on the right-hand side converges to $\Phi(c^*x - z_{1-\alpha})$ by Lemma B.8.

We next turn to the second limit. Consider testing $H_0 : \delta \in \Delta = \{\delta : A\delta \leq d\}, \theta = \bar{\theta}$ against $H_1 : (\delta, \tau) = (\delta_A, \tau_A)$. Observe that the null is equivalent to $H_0 : \beta \in \mathcal{B}_0(\bar{\theta}) := \{\beta : \exists \tau_{post} \text{ s.t. } l'\tau_{post} = \bar{\theta}, A\beta - d - AM_{post}\tau_{post} \leq 0\}$ and the alternative is equivalent to $H_1 : \beta = \delta_A + \tau_A =: \beta_A$. It is clear from the definition of \mathcal{B}_0 that it is convex. From Lemma B.5, the most powerful test that controls size is a one-sided t-test (Neyman-Pearson test) that rejects for large values of $(\beta_A - \tilde{\beta}_n)' \Sigma_n^{-1} \hat{\beta}_n$, where $\tilde{\beta} := \arg \min_{\beta \in \mathcal{B}_0} \|\beta_A - \beta\|_{\Sigma_n}$. Define $\psi_\alpha^{MP}(\hat{\beta}_n; A, d, \bar{\theta}, \Sigma_n, \delta_A, \tau_A)$ to be an indicator for whether the Neyman-Pearson test rejects H_0 in favor H_1 given a draw $\hat{\beta}_n$ that is assumed to be normally distributed with covariance Σ_n . The second limit can thus be written as

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_\alpha(\Delta, \frac{1}{n}\Sigma^*)} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n} \right) = \\ & \lim_{n \rightarrow \infty} \mathbb{E}_{(\delta_A, \tau_A, \Sigma_n)} \left[\psi_\alpha^{MP}(\hat{\beta}_n; A, d, \theta^{ub} + \frac{1}{\sqrt{n}}x, \Sigma_n, \delta_A, \tau_A) \right] \end{aligned}$$

From Lemma B.6 and the fact that $\Sigma_n = \frac{1}{n}\Sigma^*$, $\psi_\alpha^{MP}(\hat{\beta}_n; A, d, \theta^{ub} + \frac{1}{\sqrt{n}}x, \Sigma_n, \delta_A, \tau_A) = \psi_\alpha^{MP}(\sqrt{n}\hat{\beta}_n; A, \sqrt{n}d, \sqrt{n}\theta^{ub} + x, \Sigma^*, \sqrt{n}\delta_A, \sqrt{n}\tau_A)$. It follows that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sup_{\mathcal{C}_{\alpha, n} \in \mathcal{I}_\alpha(\Delta, \Sigma_n)} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left(\left(\theta^{ub} + \frac{1}{\sqrt{n}}x \right) \notin \mathcal{C}_{\alpha, n} \right) \stackrel{(1)}{=} \\ & \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^{MP}(\hat{\beta}_n; A, \sqrt{n}d, \sqrt{n}\theta^{ub} + x, \Sigma^*, \sqrt{n}\delta_A, \sqrt{n}\tau_A) \right] \stackrel{(2)}{=} \\ & \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^{MP}(\hat{\beta}_n; A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*, \sqrt{n}\delta_A, \sqrt{n}\tau_A) \right] \stackrel{(3)}{\rightarrow} \Phi(c^*x - z_{1-\alpha}), \end{aligned}$$

where (1) used that if $\hat{\beta}_n \sim \mathcal{N}(\beta_A, \frac{1}{n}\Sigma^*)$, then $\sqrt{n}\hat{\beta}_n \sim \mathcal{N}(\sqrt{n}\beta_A, \Sigma^*)$, (2) used that $\theta_n^{ub} = \sqrt{n}\theta^{ub}$ and (3) follows by Lemma B.12. \square

Proof of Proposition A.1

Proof. Following the same argument as in the proof to Proposition 3.2, we can show that $\mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left((\theta^{ub} + \frac{x}{\sqrt{n}}) \in \mathcal{C}_{\alpha, n}^{FLCI} \right)$ equals

$$\Phi \left(\frac{\theta^{mid} + \bar{b}_{min} + \frac{x}{\sqrt{n}} + \chi_n - \mu_n}{\sigma_{v_n, n}} \right) - \Phi \left(\frac{\theta^{mid} + \bar{b}_{min} + \frac{x}{\sqrt{n}} - \chi_n - \mu_n}{\sigma_{v_n, n}} \right),$$

where $\bar{b}_{min} = \frac{1}{2}LID(\Delta, \delta_{A, pre})$, $\mu_n = a_n + v'_n(\delta_A + \tau_A)$, and θ^{mid} is the midpoint of $\mathcal{S}(\Delta, \delta_A + \tau_A)$.

The term inside the second normal CDF in the previous display equals

$$-\frac{\chi_n - b_n}{\sigma_{v_n, n}} + \frac{x}{\sqrt{n}\sigma_{v_n, n}} + \frac{\theta^{mid} - \mu_n + \bar{b}_{min} - b_n}{\sigma_{v_n, n}}. \quad (28)$$

We first show the first term in (28) converges to $-z_{1-\alpha}$. Since χ_n is the $1 - \alpha$ quantile of the $|\mathcal{N}(b_n, \sigma_{v_n, n}^2)|$ distribution, $\Phi \left(\frac{\chi_n - b_n}{\sigma_{v_n, n}} \right) - \Phi \left(\frac{-\chi_n - b_n}{\sigma_{v_n, n}} \right) = 1 - \alpha$. Lemma B.25 implies that $\bar{b}_{min} = \frac{1}{2} \sup_{\tilde{\delta}_{pre}} LID(\Delta, \tilde{\delta}_{pre}) > 0$. We argued in the proof to Proposition 3.2 that $b_n \geq \bar{b}_{min} > 0$, $\chi_n \geq 0$, and $\sigma_{v_n, n} \rightarrow 0$, from which we see that the term inside the second normal CDF in the previous display diverges to $-\infty$, and thus the second normal CDF term converges to 0. It follows that the first CDF term in the previous display converges to $1 - \alpha$, and hence $\frac{\chi_n - b_n}{\sigma_{v_n, n}} \rightarrow z_{1-\alpha}$.

Next, we show the second term in (28) converges to c^*x , for the constant c^* defined in Proposition 4.2. Lemma B.23 implies that $\lim_{n \rightarrow \infty} \frac{x}{\sqrt{n}\sigma_{v_n, n}} = \lim_{n \rightarrow \infty} \frac{x}{\sqrt{n}\sigma_{\bar{v}, n}} = \frac{x}{\sigma_{\bar{v}, 1}}$, where \bar{v} is the unique value such that there exists (\bar{a}, \bar{v}) with $\bar{b}(\bar{a}, \bar{v}) = \bar{b}_{min}$. Moreover, Lemma B.22 implies that $1/\sigma_{\bar{v}, 1} = c^*$, from which we see that the limit of the second term is c^*x , as desired.

Now, we show the third term in (28) converges to 0. We argued in the proof to Proposition 3.2 that $|\mu_n - \theta^{mid}| \leq b_n - \bar{b}_{min}$. It thus suffices to show that $\frac{b_n - \bar{b}_{min}}{\sigma_{v_n, n}} \rightarrow 0$. Lemma B.22 implies that there is a unique pair (\bar{a}, \bar{v}) such that $\bar{b}(\bar{a}, \bar{v}) = \bar{b}_{min}$. Let $\bar{\chi}_n = \chi_n(\bar{a}, \bar{v})$ and $\chi_n = \chi_n(a_n, v_n)$. Note that $\chi_n \leq \bar{\chi}_n$ by construction, and $b_n \geq \bar{b}_{min}$ by Lemma B.19. Hence, using the bounds from Lemma B.21, we have that $b_n + \sigma_{v_n, n}z_{1-\alpha} \leq \chi_n \leq \bar{\chi}_n = \sigma_{\bar{v}, n}cV_\alpha \left(\frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right)$, which, along with the inequality

$b_n \geq \bar{b}_{min}$, implies that

$$\begin{aligned} 0 \leq \frac{b_n - \bar{b}_{min}}{\sigma_{v_n, n}} &\leq \frac{\sigma_{\bar{v}, n}}{\sigma_{v_n, n}} cv_\alpha \left(\frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right) - \left(z_{1-\alpha} + \frac{\bar{b}_{min}}{\sigma_{v_n, n}} \right) \\ &= \left[cv_\alpha \left(\frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right) - \left(z_{1-\alpha} + \frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right) \right] + \\ &\quad \left[\left(\frac{\sigma_{\bar{v}, n}}{\sigma_{v_n, n}} - 1 \right) cv_\alpha \left(\frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right) - \left(\frac{\bar{b}_{min}}{\sigma_{v_n, n}} - \frac{\bar{b}_{min}}{\sigma_{\bar{v}, n}} \right) \right]. \end{aligned}$$

The first bracketed expression in the upper bound above converges to 0 by Lemma B.24. Applying the upper bound from Lemma B.21 to the cv_α term in the second bracketed expression, we obtain that the second bracketed expression is bounded above by $\left(\frac{\sigma_{\bar{v}, n}}{\sigma_{v_n, n}} - 1 \right) z_{1-\alpha/2}$, which converges to 0 by Lemma B.23.

Combining the results above, we see that the expression in (28) converges to $c^*x - z_{1-\alpha}$. It follows that $\limsup_{n \rightarrow \infty} \mathbb{P} \left((\theta^{ub} + \frac{x}{\sqrt{n}}) \in \mathcal{C}_{\alpha, n}^{FLCI} \right) \leq 1 - \Phi(c^*x - z_{1-\alpha})$, and hence $\liminf_{n \rightarrow \infty} \mathbb{P} \left((\theta^{ub} + \frac{x}{\sqrt{n}}) \notin \mathcal{C}_{\alpha, n}^{FLCI} \right) \geq \Phi(c^*x - z_{1-\alpha})$. Proposition 4.2 gives that $\Phi(c^*x - z_{1-\alpha})$ is the optimal local asymptotic power over procedures that control size, from which the result follows. \square

Proof of Proposition A.2

Proof. The proof follows from the same argument as for Proposition 4.1, replacing Lemma B.2 with Lemma B.4, and Lemma B.16 with Lemma B.17. \square

Proof of Proposition A.3

Proof. By an invariance to scale argument analogous to that in the proof of Proposition 4.2, $\liminf_{n \rightarrow \infty} \mathbb{P}_{(\delta_A, \tau_A, \Sigma_n)} \left((\theta_A^{ub} + \frac{1}{\sqrt{n}}x) \notin \mathcal{C}_{\kappa, \alpha, n}^{C-FLCI} \right)$ equals

$$\liminf_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_{\kappa, \alpha}^{C-FLCI}(\hat{\beta}_n, A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) \right].$$

From Proposition 4.2, it thus suffices to show

$$\begin{aligned} \liminf_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_{\kappa, \alpha}^{C-FLCI}(\hat{\beta}_n, A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) \right] &\geq \\ \liminf_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_{\tilde{\alpha}}^C(\hat{\beta}_n, A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) \right]. \end{aligned}$$

Note that the second stage of the test $\psi_{\kappa,\alpha}^{C-FLCI}$ is nearly identical to ψ_{α}^C except it uses $v_{C-FLCI}^{lo} := \max\{v^{lo}, v_{FLCI}^{lo}\}$ and $v_{C-FLCI}^{up} := \min\{v^{up}, v_{FLCI}^{up}\}$ instead of v^{lo} and v^{up} . Since $F_{\xi:\xi \in [v^{lo}, v^{up}]}(\hat{\eta})$ is decreasing in v^{lo} and v^{up} , it suffices to show that $v_{FLCI}^{lo} \xrightarrow{P_n} -\infty$, where P_n denotes the sequence of distributions under which $(\delta, \tau, \Sigma) = (\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)$.

Let $\Delta_n = \{\delta : A\delta \leq \sqrt{nd}\}$. Let $v_n = v_n(\Delta_n, \Sigma^*)$ and $\check{v}_n = v_n(\Delta, \Sigma_n)$. Define a_n and \check{a}_n , and $\check{\chi}_n$ and χ_n analogously. By Lemma B.3, $v_n = \check{v}_n$, $a_n = \sqrt{n}\check{a}_n$, $\chi_n = \sqrt{n}\check{\chi}_n$, and $\bar{b}(a, v; \Delta_n) = \sqrt{n}\bar{b}(\check{a}, \check{v}, \Delta)$. We argued in the proof to Lemma B.23 that $\check{v}_n \rightarrow \bar{v}$. Further, we showed in the proof to Lemma B.22 that $\bar{v} = -\bar{\gamma}A$, where $\bar{\gamma}_{-B} = 0$ and $\bar{\gamma}_B$ is the unique vector such that $\bar{\gamma}'_B \tilde{A}_{(B,1)} = 1, \bar{\gamma}'_B \tilde{A}_{(B,-1)} = 0$. Likewise, we argued in the proof to Lemma B.23 that $\check{a}_n \rightarrow \bar{a}$, for \bar{a} the unique value such that $\bar{b}(\bar{a}, \bar{v}; \Delta) = b_{min}(\Delta)$. We also showed in the proof to Proposition 3.2 that $\check{\chi}_n \rightarrow b_{min}(\Delta)$.

Let \tilde{v}_n be a vector such that $\tilde{v}'_n A = v'$ (which exists by Lemma A.2). Observe that

$$\tilde{v}'_n \tilde{\Sigma} \tilde{\gamma} = \tilde{v}'_n A \Sigma^* A' \tilde{\gamma} \rightarrow \bar{v}' \Sigma^* A' \tilde{\gamma} = -\bar{\gamma}' A \Sigma^* A' \tilde{\gamma} = -\bar{\gamma}' \tilde{\Sigma} \tilde{\gamma},$$

where we use $\tilde{v}'_n A = v_n$, $v_n \rightarrow \bar{v} = -\bar{\gamma}' A$ as shown above, and the identity $\tilde{\Sigma} = A \Sigma^* A'$.

Now, Lemma B.27 implies that there is a constant $c > 0$ such that, with probability approaching one under P_n , $c \cdot \bar{\gamma}$ is an optimal vertex of the dual problem for $\psi_{\kappa,\alpha}^{C-FLCI}(\hat{\beta}_n, A, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*)$. Observe from Lemma A.3 that if γ_* is an optimal vertex, and $\frac{v'_n \tilde{\Sigma} \gamma_*}{\gamma'_* \tilde{\Sigma} \gamma_*} < 0$, then the value of v_{FLCI}^{lo} used in $\psi_{\kappa,\alpha}^{C-FLCI}(\hat{\beta}_n, A, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*)$ is

$$v_{FLCI}^{lo} = \frac{d_{n,1} - \tilde{v}'_n \left(I - \frac{\tilde{\Sigma} \gamma_*}{\gamma'_* \tilde{\Sigma} \gamma_*} \gamma'_* \right) \tilde{Y}_n}{\tilde{v}'_n \tilde{\Sigma} \gamma_* / \gamma'_* \tilde{\Sigma} \gamma_*} = \frac{-(\tilde{v}'_n \tilde{Y}_n - d_{n,1})}{v'_n \tilde{\Sigma} \gamma_* / \gamma'_* \tilde{\Sigma} \gamma_*} + \gamma'_* \tilde{Y}_n, \quad (29)$$

where $\tilde{Y}_n = A \hat{\beta}_n - \sqrt{nd} - \tilde{A}_{(\cdot,1)}(\theta_n^{ub} + x)$. Since $c\bar{\gamma}$ is optimal in the dual with probability approaching one under P_n , and

$$\frac{v'_n \tilde{\Sigma} c \bar{\gamma}}{c^2 \bar{\gamma}' \tilde{\Sigma} \bar{\gamma}} \rightarrow -\frac{1}{c} < 0 \quad (30)$$

by the argument above, we have that with probability approaching 1 under P_n ,

$$v_{FLCI}^{lo} = \frac{-(\tilde{v}'_n \tilde{Y}_n - d_{n,1})}{v'_n \tilde{\Sigma} c \tilde{\gamma} / c^2 \tilde{\gamma}' \tilde{\Sigma} \tilde{\gamma}} + c \tilde{\gamma}' \tilde{Y}_n. \quad (31)$$

Now, we showed in the proof to Lemma B.8 that $\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\check{Y}_{n,B}] = -\tilde{A}_{(B,1)}x$ regardless of n , where $\check{Y}_n = \tilde{Y}_n - \tilde{A}_{(\cdot,-1)}(\sqrt{n}\tilde{\tau}_1^{ub})$ for a vector $\tilde{\tau}_1^{ub}$. Since $\tilde{\gamma}_{-B} = 0$ and $\tilde{\gamma}'_B \tilde{A}_{(B,-1)} = 0$, it follows that $\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\tilde{\gamma}' \tilde{Y}_n] = -\tilde{\gamma}'_B \tilde{A}_{(B,1)}x$ regardless of n . Thus,

$$c \tilde{\gamma}' \tilde{Y}_n \stackrel{P_n}{\approx} \mathcal{N} \left(-c \tilde{\gamma}'_B \tilde{A}_{(B,1)}x, c^2 \tilde{\gamma}' \tilde{\Sigma} \tilde{\gamma} \right). \quad (32)$$

Now, note that by construction, $\tilde{v}'_n \tilde{Y}_n - d_{n,1} = a_n + v'_n \hat{\beta} - (\theta_n^{ub} + x) - \chi_n$. Further, we have that $\hat{\beta}_n \stackrel{P_n}{\approx} \mathcal{N}(\sqrt{n}\beta_A, \Sigma^*)$, where $\beta_A = \delta_A + \tau_A$. It follows that under P_n , $\tilde{v}'_n \tilde{Y}_n - d_{n,1} = a_n + v'_n \sqrt{n}\beta_A - (\theta_n^{ub} + x) - \chi_n + v'_n \xi$, where $\xi \sim \mathcal{N}(0, \Sigma^*)$. Applying the equalities $v_n = \check{v}_n$, $a = \sqrt{n}\check{a}_n$, $\chi_n = \sqrt{n}\check{\chi}$ derived above, along with the fact that $\theta_n^{ub} = \sqrt{n}\theta_1^{ub}$ by Lemma B.1, we see that under P_n ,

$$\tilde{v}'_n \tilde{Y}_n - \tilde{d}_1 = \sqrt{n} (\check{a}_n + \check{v}'_n \beta_A - \theta_1^{ub} - \check{\chi}_n) - x + \check{v}'_n \xi, \quad (33)$$

Since $\check{v}'_n \rightarrow \bar{v}$, $\check{v}'_n \xi \rightarrow_d \bar{v}' \xi$ by Slutsky's lemma.

Additionally, the results above imply that $\check{a}_n + \check{v}'_n \beta_A - \theta_1^{ub} - \check{\chi}_n \rightarrow \bar{a} + \bar{v}' \beta_A - \theta_1^{ub} - \bar{b}_{min}(\Delta)$. We claim that this limit is strictly negative. Since Assumption 5 holds, Lemma B.25 implies that $LID(\Delta, \delta_A) > 0$. Hence, for $\epsilon > 0$ sufficiently small, we have that $\theta_1^{ub} - \epsilon \in \mathcal{S}(\Delta, \beta_A)$. If the limit above were weakly positive, then we would have $\bar{a} + \bar{v}' \beta_A - (\theta_1^{ub} - \epsilon) - \bar{b}_{min}(\Delta) > 0$. However, this implies that $\bar{b}(\bar{a}, \bar{v}) > \bar{b}_{min}(\Delta)$, which is a contradiction. The limit must thus be strictly negative, as desired. We then see from (33) that

$$\tilde{v}'_n \tilde{Y}_n (\theta_n^{ub} + x) - d_{n,1} \xrightarrow{P_n} -\infty. \quad (34)$$

Displays (30), (31), (32), and (34) together give that $v_{FLCI}^{lo} \xrightarrow{P_n} -\infty$, as desired. \square

B.2 Auxiliary Lemmas for Finite Sample Normal Results

Lemma B.1. For any $n > 0$, let $\Delta_n = \{\delta : A\delta \leq \sqrt{nd}\}$. Fix $\delta_A \in \Delta_1$ and τ_A . Then, $\mathcal{S}(\sqrt{n}\delta_A + \sqrt{n}\tau_A, \Delta_n) = \sqrt{n}\mathcal{S}(\delta_A + \tau_A, \Delta_1)$. This implies $\theta_n^{ub} = \sqrt{n}\theta_1^{ub}$, where $\theta_n^{ub} := \sup_{\theta} \mathcal{S}(\sqrt{n}\delta_A + \sqrt{n}\tau_A, \Delta_n)$, and $\theta_n^{lb} = \sqrt{n}\theta_1^{lb}$, for $\theta_n^{lb} := \inf_{\theta} \mathcal{S}(\sqrt{n}\delta_A + \sqrt{n}\tau_A, \Delta_n)$

Proof. Let $\mathcal{S}_n = \mathcal{S}(\sqrt{n}\delta_A + \sqrt{n}\tau_A, \Delta_n)$ and $\beta_A = \delta_A + \tau_A$. By definition, $\bar{\theta}_n \in \mathcal{S}_n$ iff there exists a vector $\tau_{post} \in \mathbb{R}^T$ such that $l'\tau_{post} = \bar{\theta}_n$ and $A(\sqrt{n}\beta_A - M_{post}\tau_{post} - \sqrt{nd}) \leq 0$. Using the change of basis described in Section 4.1, it follows that $\bar{\theta}_n \in \mathcal{S}_n$ iff there exists $\tilde{\tau}_n \in \mathbb{R}^{T-1}$ such that

$$A\sqrt{n}\beta_A - \sqrt{nd} - \tilde{A}_{(\cdot,1)}\bar{\theta}_n - \tilde{A}_{(\cdot,-1)}\tilde{\tau}_n \leq 0. \quad (35)$$

Thus, $\bar{\theta}_1 \in \mathcal{S}_1$ iff there exists $\tilde{\tau}_1$ such that

$$A\beta_A - d - \tilde{A}_{(\cdot,1)}\bar{\theta}_1 - \tilde{A}_{(\cdot,-1)}\tilde{\tau}_1 \leq 0. \quad (36)$$

If there exists a $\tilde{\tau}_1$ such that (36) holds for $\bar{\theta}_1$, then multiplying both sides of (36) by \sqrt{n} implies that (35) holds with $\bar{\theta}_n = \sqrt{n}\bar{\theta}_1$ and $\tilde{\tau}_n = \sqrt{n}\tilde{\tau}_1$. Likewise, if there exists a $\tilde{\tau}_n$ such that (35) holds for $\bar{\theta}_n$, then multiplying both sides of (35) by $\frac{1}{\sqrt{n}}$ implies that (36) holds with $\bar{\theta}_1 = \frac{1}{\sqrt{n}}\bar{\theta}_n$ and $\tilde{\tau}_1 = \frac{1}{\sqrt{n}}\tilde{\tau}_n$. The desired result follows immediately. \square

Lemma B.2. For any $n > 0$ and $(\hat{\beta}; A, d, \bar{\theta}, \Sigma)$, $\psi_{\alpha}^C(\hat{\beta}; A, d, \bar{\theta}, \Sigma) = \psi_{\alpha}^C(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma)$.

Proof. Using the change of basis described in Section 4.1, the test statistic used to calculate $\psi_{\alpha}^C(\hat{\beta}; A, d, \bar{\theta}, \Sigma)$ is

$$\min_{\eta, \tilde{\tau}} \eta \quad \text{s.t.} \quad A\hat{\beta} - d - \tilde{A}_{(\cdot,1)}\bar{\theta} - \tilde{A}_{(\cdot,-1)}\tilde{\tau} \leq \eta\tilde{\sigma},$$

where $\tilde{\sigma}$ is the vector containing the square roots of the diagonal elements of $\tilde{\Sigma} = A\Sigma A'$. Since multiplying the constraints by \sqrt{n} does not change the feasible set, this optimization is equivalent to

$$\min_{\eta, \tilde{\tau}} \eta \quad \text{s.t.} \quad A\sqrt{n}\hat{\beta} - \sqrt{nd} - \tilde{A}_{(\cdot,1)}\sqrt{n}\bar{\theta} - \tilde{A}_{(\cdot,-1)}\sqrt{n}\tilde{\tau} \leq \eta\sqrt{n}\tilde{\sigma}.$$

However, since $\tilde{\tau}$ enters only in the constraint, and $\{\sqrt{n}\tilde{\tau} : \tilde{\tau} \in \mathbb{R}^{T-1}\} = \{\tilde{\tau} \in \mathbb{R}^{T-1}\}$, this linear program is equivalent to

$$\min_{\eta, \tilde{\tau}} \eta \text{ s.t. } A\sqrt{n}\hat{\beta} - \sqrt{nd} - \tilde{A}_{(\cdot,1)}\sqrt{n}\bar{\theta} - \tilde{A}_{(\cdot,-1)}\tilde{\tau} \leq \eta\sqrt{n}\tilde{\sigma},$$

which is the test statistic used to calculate $\psi_\alpha^C(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma)$. Thus, the test statistics used for the two problems are the same. Additionally, the feasible set for the dual for the unscaled problem is $F_1 = \{\gamma : \gamma' \tilde{A}_{(\cdot,-1)} = 0, \gamma' \tilde{\sigma} = 1, \gamma \geq 0\}$, whereas for the problem scaled by \sqrt{n} it is $F_n = \{\gamma : \gamma' \tilde{A}_{(\cdot,-1)} = 0, \gamma' \sqrt{n}\tilde{\sigma} = 1, \gamma \geq 0\} = \frac{1}{\sqrt{n}}F_1$. It follows that $V_n = \frac{1}{\sqrt{n}}V_1$, for V_1 and V_n respectively the vertices of F_1 and F_n . Moreover, it is immediate that if γ_1 is an optimal vertex of the unscaled problem, then $\gamma_n = \frac{1}{\sqrt{n}}\gamma_1$ will be an optimal vertex of the problem scaled by \sqrt{n} .

Recall that the critical value for the conditional test depends on $\gamma'_* \tilde{\Sigma} \gamma_*$, where γ_* is an optimal vertex, and the values v^{lo} and v^{up} which are functions of γ_* , $\tilde{\Sigma}$, and a sufficient statistic S . However, for $\gamma_n = \frac{1}{\sqrt{n}}\gamma_1$, we have that $\gamma'_n(n\tilde{\Sigma})\gamma_n = \frac{1}{\sqrt{n}}\gamma'_1(n\tilde{\Sigma})\frac{1}{\sqrt{n}}\gamma_1 = \gamma'_1\tilde{\Sigma}\gamma_1$, and so the variances are the same. Let $\tilde{Y}_1 = A\hat{\beta} - d - \tilde{A}_{(\cdot,1)}\bar{\theta}$ and $\tilde{Y}_n = A\sqrt{n}\hat{\beta} - \sqrt{nd} - \tilde{A}_{(\cdot,1)}\sqrt{n}\bar{\theta} = \sqrt{n}\tilde{Y}_1$. The sufficient statistic used to construct v^{lo} and v^{up} in the first problem is $S_1 = (I - \frac{\tilde{\Sigma}\gamma_1\gamma'_1}{\gamma'_1\tilde{\Sigma}\gamma_1})\tilde{Y}_1$, whereas for the second problem it is $S_n = (I - \frac{n\tilde{\Sigma}\gamma_n\gamma'_n}{\gamma'_n n\tilde{\Sigma}\gamma_n})\tilde{Y}_n$. The identities $\tilde{Y}_n = \sqrt{n}\tilde{Y}_1$ and $\gamma_1 = \sqrt{n}\gamma_n$ immediately imply that $S_n = \sqrt{n}S_1$. The values v^{lo} and v^{up} for the first problem are then the minimum and maximum of $C_1 = \{c : c = \max_{\tilde{\gamma}_1 \in V_1} \tilde{\gamma}'_1(S_1 + \frac{\tilde{\Sigma}\gamma_1}{\gamma'_1\tilde{\Sigma}\gamma_1}c)\}$. Likewise, the values v^{lo} and v^{up} for the second problem are the the minimum and maximum of $C_n = \{c : c = \max_{\tilde{\gamma}_n \in V_n} \tilde{\gamma}'_n(S_n + \frac{n\tilde{\Sigma}\gamma_n}{\gamma'_n n\tilde{\Sigma}\gamma_n}c)\}$. However, since $V_n = \sqrt{n}V_1$, $S_n = \sqrt{n}S_1$, and $\gamma_n = \frac{1}{\sqrt{n}}\gamma_1$, we have that for any c ,

$$\max_{\tilde{\gamma}_1 \in V_1} \tilde{\gamma}'_1(S_1 + \frac{\tilde{\Sigma}\gamma_1}{\gamma'_1\tilde{\Sigma}\gamma_1}c) = \max_{\tilde{\gamma}_n \in V_n} \sqrt{n}\tilde{\gamma}'_n(\frac{1}{\sqrt{n}}S_n + \frac{\tilde{\Sigma}\sqrt{n}\gamma_n}{\gamma'_n n\tilde{\Sigma}\gamma_n}c) = \max_{\tilde{\gamma}_n \in V_n} \tilde{\gamma}'_n(S_n + \frac{n\tilde{\Sigma}\gamma_n}{\gamma'_n n\tilde{\Sigma}\gamma_n}c),$$

from which it is immediate that $C_1 = C_n$, and hence the values of v^{lo} and v^{up} are the same across the two problems as well. Since the test statistics and critical values of the two problems are the same, they are equivalent. \square

Lemma B.3. *Let $v_n(\Delta, \Sigma_n)$ be the vector v_n used in the optimal FLCI as defined in Section 3, making the dependence on (Δ, Σ_n) explicit. Define, $\chi_n(\Delta, \Sigma_n)$, $a_n(\Delta, \Sigma_n)$, and $\bar{b}_n(\Delta, \Sigma_n)$ analogously. Let $\Sigma_n = \frac{1}{n}\Sigma^*$ and $\Delta_n = \sqrt{n}\Delta$. Then*

1. $v_n(\Delta_n, \Sigma^*) = v_n(\Delta, \frac{1}{n}\Sigma_n)$
2. $a_n(\Delta_n, \Sigma^*) = \sqrt{n}a_n(\Delta, \frac{1}{n}\Sigma_n)$

$$3. \chi_n(\Delta_n, \Sigma^*) = \sqrt{n}\chi_n(\Delta, \frac{1}{n}\Sigma_n)$$

$$4. \bar{b}(a_n(\Delta_n, \Sigma^*), v_n(\Delta_n, \Sigma^*); \Delta_n) = \sqrt{n}\bar{b}(a_n(\Delta, \Sigma_n), v_n(\Delta, \Sigma_n); \Delta).$$

Proof. Recall the proof to Lemma B.19 that $\bar{b}(a, v; \Delta_n)$ is finite only if $v_{post} = l$, in which case $\bar{b}(a, v; \Delta) = \max_{\delta \in \Delta} |a + v'\delta|$. Likewise, $\bar{b}(\sqrt{na}, v; \Delta_n)$ is finite only if $v_{post} = l$, in which case

$$\bar{b}(\sqrt{na}, v; \Delta) = \max_{\delta_n \in \Delta_n} |\sqrt{na} + v'\delta_n| = \max_{\delta \in \Delta} |\sqrt{na} + v'\sqrt{n}\delta| = \sqrt{n}\bar{b}(a, v, \Delta).$$

Next, observe that using the invariance above and $\Sigma_n = \frac{1}{n}\Sigma^*$,

$$\begin{aligned} \chi(\sqrt{na}, v; \Sigma^*, \Delta_n) &= \sqrt{v'\Sigma^*v} \cdot cv_\alpha \left(\frac{\bar{b}(\sqrt{na}, v; \Delta_n)}{\sqrt{v'\Sigma^*v}} \right) \\ &= \sqrt{n} \cdot \sqrt{v'\Sigma_n v} \cdot cv_\alpha \left(\frac{\bar{b}(a, v; \Delta)}{\sqrt{v'\Sigma_n v}} \right) = \sqrt{n}\chi(a, v; \Sigma^*, \Delta). \end{aligned}$$

It is then immediate that if $(a^*, v^*) = \arg \min_{(a,v)} \chi(a, v; \Delta, \Sigma_n)$, then $(\sqrt{na}^*, v^*) = \arg \min_{(a,v)} \chi(a, v; \Delta_n, \Sigma^*)$, from which the first two results follow. The second two results then follow from the two invariances derived above. \square

Lemma B.4. *For any $n > 0$ and $(\hat{\beta}; A, d, \bar{\theta}, \Sigma)$,*

$$\psi_{\kappa, \alpha}^{C-FLCI}(\hat{\beta}; A, d, \bar{\theta}, \Sigma) = \psi_{\kappa, \alpha}^{C-FLCI}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma).$$

Proof. From Lemma B.3, if $\mathcal{C}_\kappa^{FLCI}(\hat{\beta}; A, d, \bar{\theta}, \Sigma) = a_1 + v_1'\hat{\beta} \pm \chi_1$, then $\mathcal{C}_\kappa^{FLCI}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, \sqrt{n}\Sigma) = \sqrt{n} \left(a_1 + v_1'\hat{\beta} \pm \chi_1 \right)$. Thus, $\bar{\theta} \in \mathcal{C}_\kappa^{FLCI}(\hat{\beta}; A, d, \bar{\theta}, \Sigma)$ iff $\sqrt{n}\bar{\theta} \in \mathcal{C}_\kappa^{FLCI}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, \sqrt{n}\Sigma)$, so the first stage tests are equivalent. The second stage test is almost identical to ψ_α^C , which is invariant to scale by Lemma B.2, except it replaces v^{lo} with $\max\{v^{lo}, v_{FLCI}^{lo}\}$ and v^{up} with $\min\{v^{up}, v_{FLCI}^{up}\}$. It thus suffices to show that v_{FLCI}^{lo} and v_{FLCI}^{up} are invariant to scale. We show in the proof to Lemma B.2 that $S_n = \sqrt{n}S_1$, and if $\gamma_n \in \hat{V}_n$ then $\gamma_1 = \sqrt{n}\gamma_n \in \hat{V}_n$; where objects subscripted by 1 indicate values based on $(\hat{\beta}; A, d, \bar{\theta}, \Sigma)$ and values subscripted by n indicate those based on $(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, \sqrt{n}\Sigma)$. Additionally, Lemma B.3 implies that $\tilde{V}_1 = \tilde{V}_n$, and $\bar{d}_n = \sqrt{n}\bar{d}_n$. The desired invariance is then immediate from the formulas in Lemma A.3. \square

Lemma B.5. *Suppose $\hat{\beta} \sim \mathcal{N}(\beta, \Sigma)$ for Σ known. Let B_0 be a closed, convex set. Then the most-powerful size α test of $H_0 : \beta \in B_0$ against the point alternative*

$H_A : \beta = \beta_A$ is equivalent to the most powerful test of $H_0 : \beta = \tilde{\beta}$ against $H_A : \beta = \beta_A$, where $\tilde{\beta} = \arg \min_{\beta} \|\beta - \beta_A\|_{\Sigma}$ and $\|\cdot\|_{\Sigma}$ is the Mahalanobis norm in Σ , $\|x\|_{\Sigma} = \sqrt{x'\Sigma^{-1}x}$. The most powerful test rejects for values of $(\beta_A - \tilde{\beta})'\Sigma^{-1}\hat{\beta}$ greater than $(\beta_A - \tilde{\beta})'\Sigma^{-1}\tilde{\beta} + z_{1-\alpha}\|\beta_A - \tilde{\beta}\|_{\Sigma}$, and has power against the alternative of $\Phi(\|\beta_A - \tilde{\beta}\|_{\Sigma} - z_{1-\alpha})$, for $z_{1-\alpha}$ the $1 - \alpha$ quantile of the standard normal.

Proof. Define $\langle \cdot, \cdot \rangle_{\Sigma}$ by $\langle x, y \rangle_{\Sigma} = x'\Sigma^{-1}y$, and observe that $\langle \cdot, \cdot \rangle_{\Sigma}$ is an inner product. The result then follows immediately from the discussion in Section 2.4.3 of [Ingster and Suslina \(2003\)](#), replacing all instances of the usual euclidean inner product with $\langle \cdot, \cdot \rangle_{\Sigma}$. \square

Lemma B.6. *Suppose $\Delta = \{\delta : A\delta \leq d\}$. As in the proof to Proposition 4.2, let $\psi_{\alpha}^{MP}(\hat{\beta}; A, d, \bar{\theta}, \Sigma, \delta_A, \tau_A)$ be an indicator for whether the most powerful (Neyman-Pearson) test between the null hypothesis $H_0 : \delta \in \Delta, \theta = \bar{\theta}$ and the alternative $H_A : (\delta, \tau) = (\delta_A, \tau_A)$ rejects the null, given the realization $\hat{\beta}$ which is assumed to come from a normal distribution with variance Σ . Then for any $n > 0$,*

$$\psi_{\alpha}^{MP}(\hat{\beta}; A, d, \bar{\theta}, \Sigma, \delta_A, \tau_A) = \psi_{\alpha}^{MP}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma, \sqrt{n}\delta_A, \sqrt{n}\tau_A)$$

Proof. As argued in the proof to Proposition 4.2, the null $H_0 : \delta \in \Delta, \theta = \bar{\theta}$ is equivalent to the null $H_0 : \beta \in \mathcal{B}_0(\bar{\theta}, d) := \{\beta : \exists \tau_{post} \text{ s.t. } l'\tau_{post} = \bar{\theta}, A\beta - d - AM_{post}\tau_{post} \leq 0\}$. Likewise, the alternative that $(\delta, \tau) = (\delta_A, \tau_A)$ is equivalent to $H_A : \beta = \delta_A + \tau_A =: \beta_A$. It is clear from the definition that \mathcal{B}_0 is convex. Thus, by Lemma B.5, the most powerful test of H_0 against H_A when the covariance of $\hat{\beta}$ is Σ is a t-test that rejects for large values of $(\beta_A - \tilde{\beta}_1)'\Sigma^{-1}\hat{\beta}$, where $\tilde{\beta}_1 = \arg \min_{\beta \in \mathcal{B}_0(\bar{\theta}, d)} \|\beta_A - \beta\|_{\Sigma}$. Its critical value is $(\beta_A - \tilde{\beta}_1)'\Sigma^{-1}\tilde{\beta}_1 + z_{1-\alpha}\|\beta_A - \tilde{\beta}_1\|_{\Sigma}$, for $z_{1-\alpha}$ the $1 - \alpha$ quantile of the standard normal distribution.

Similarly, the null hypothesis $\delta \in \{\delta : A\delta \leq \sqrt{nd}\}, \theta = \sqrt{n}\bar{\theta}$ is equivalent to $H_0 : \beta \in \mathcal{B}_0(\sqrt{n}\bar{\theta}, \sqrt{nd}) = \{\beta : \exists \tau_{post} \text{ s.t. } l'\tau_{post} = \sqrt{n}\bar{\theta}, A\beta - \sqrt{nd} - AM_{post}\tau_{post} \leq 0\}$. Likewise, the alternative that $(\delta, \tau) = (\sqrt{n}\delta_A, \sqrt{n}\tau_A)$ is equivalent to $H_A : \beta = \sqrt{n}\delta_A + \sqrt{n}\tau_A = \sqrt{n}\beta_A$. It is clear from the definition that $\mathcal{B}_0(\sqrt{n}\bar{\theta}, \sqrt{nd})$ is convex. Thus, by Lemma B.5, the most powerful test of H_0 against H_A when the covariance of $\hat{\beta}$ is $n\Sigma$ is a t-test that rejects for large values of $(\sqrt{n}\beta_A - \tilde{\beta}_2)'\Sigma^{-1}\hat{\beta}$, where $\tilde{\beta}_2 = \arg \min_{\beta \in \mathcal{B}_0(\sqrt{n}\bar{\theta}, \sqrt{nd})} \|\sqrt{n}\beta_A - \beta\|_{(n\Sigma)}$. Its critical value is $(\sqrt{n}\beta_A - \tilde{\beta}_2)'\Sigma^{-1}\tilde{\beta}_2 + z_{1-\alpha}\|\beta_A - \tilde{\beta}_2\|_{(n\Sigma)}$.

Now, define

$$\eta(\hat{\beta}, A, d, \bar{\theta}, \Sigma) := \min_{\eta, \tilde{\tau}} \eta \text{ s.t. } A\hat{\beta} - d - \tilde{A}_{(\cdot, 1)}\bar{\theta} - \tilde{A}_{(\cdot, -1)}\tilde{\tau} \leq \eta\tilde{\sigma},$$

where $\tilde{\sigma}$ is the square root of the diagonal elements of $A\Sigma A'$. It follows immediately from the definition of \mathcal{B}_0 and the function η that we can write

$$\begin{aligned} \mathcal{B}_0(\bar{\theta}, d) &= \{\beta : \eta(\beta, A, d, \bar{\theta}, \Sigma) \leq 0\} \\ \mathcal{B}_0(\sqrt{n}\bar{\theta}, \sqrt{nd}) &= \{\beta : \eta(\beta, A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma) \leq 0\} \end{aligned}$$

As argued in the proof to Lemma B.2 above, for any $n > 0$, $\eta(\beta, A, d, \bar{\theta}, \Sigma) = \eta(\sqrt{n}\beta, A, \sqrt{nd}, \sqrt{n}\bar{\theta}, n\Sigma)$, from which it follows that $\sqrt{n}\mathcal{B}_0(\bar{\theta}, d) = \mathcal{B}_0(\sqrt{n}\bar{\theta}, \sqrt{nd})$. Thus,

$$\begin{aligned} \tilde{\beta}_2 &= \sqrt{n} \arg \min_{\beta \in \mathcal{B}_0(\bar{\theta}, d)} \|\sqrt{n}\beta_A - \sqrt{n}\beta\|_{(n\Sigma)} \\ &= \sqrt{n} \arg \min_{\beta \in \mathcal{B}_0(\bar{\theta}, d)} \|\beta_A - \beta\|_{\Sigma} = \sqrt{n}\tilde{\beta}_1, \end{aligned}$$

where the second equality uses the fact that $\|\sqrt{n}x\|_{(n\Sigma)} = \|x\|_{\Sigma}$. Thus, the test statistic used for $\psi_{\alpha}^{MP}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, \sqrt{n}\delta_A, \sqrt{n}\tau_A)$ is

$$(\sqrt{n}\beta_A - \tilde{\beta}_2)'(n\Sigma)^{-1}(\sqrt{n}\hat{\beta}) = (\sqrt{n}\beta_A - \sqrt{n}\tilde{\beta}_1)'(n\Sigma)^{-1}(\sqrt{n}\hat{\beta}) = (\beta_A - \tilde{\beta}_1)'\Sigma^{-1}\hat{\beta},$$

which is the test statistic used for $\psi_{\alpha}^{MP}(\hat{\beta}; A, d, \bar{\theta}, \delta_A, \tau_A)$.

Likewise, the critical value used for $\psi_{\alpha}^{MP}(\sqrt{n}\hat{\beta}; A, \sqrt{nd}, \sqrt{n}\bar{\theta}, \sqrt{n}\delta_A, \sqrt{n}\tau_A)$ is

$$\begin{aligned} &(\sqrt{n}\beta_A - \tilde{\beta}_2)'(n\Sigma)^{-1}\tilde{\beta}_2 + z_{1-\alpha}\|\sqrt{n}\beta_A - \tilde{\beta}_2\|_{(n\Sigma)} = \\ &(\sqrt{n}\beta_A - \sqrt{n}\tilde{\beta}_1)'(n\Sigma)^{-1}\sqrt{n}\tilde{\beta}_1 + z_{1-\alpha}\|\sqrt{n}\beta_A - \sqrt{n}\tilde{\beta}_1\|_{(n\Sigma)} = \\ &(\beta_A - \tilde{\beta}_1)'\Sigma^{-1}\tilde{\beta}_1 + z_{1-\alpha}\|\beta_A - \tilde{\beta}_1\|_{\Sigma}, \end{aligned}$$

which is the critical value used for $\psi_{\alpha}^{MP}(\hat{\beta}; A, d, \bar{\theta}, \delta_A, \tau_A)$. We have thus shown that the test statistics and critical values for the two tests align, which gives the desired result. □

Lemma B.7. *Suppose Assumption 5 holds. Let $\theta^{ub} := \sup_{\theta} \mathcal{S}(\Delta, \delta_A + \tau_A)$ and $\beta_A =$*

$\delta_A + \tau_A$. Then there exists a vector $\tilde{\tau}^{ub} \in \mathbb{R}^{\bar{T}-1}$ such that for $B = B(\delta^{**})$ as defined in Assumption 5,

$$\begin{aligned} A_{(B,\cdot)}\beta_A - d_B - \tilde{A}_{(B,1)}\theta^{ub} - \tilde{A}_{(B,-1)}\tilde{\tau}^{ub} &= 0 \\ A_{(-B,\cdot)}\beta_A - d_{-B} - \tilde{A}_{(-B,1)}\theta^{ub} - \tilde{A}_{(-B,-1)}\tilde{\tau}^{ub} &= -\epsilon < 0, \end{aligned}$$

for ϵ a vector with strictly positive entries. Additionally, the matrix $\tilde{A}_{(B,-1)}$ has rank equal to $|B| - 1$, and $\{\gamma_B : \gamma'_B \tilde{A}_{(B,-1)} = 0\} = \{c\bar{\gamma}_B : c \in \mathbb{R}\}$ for a non-zero vector $\bar{\gamma}_B \geq 0$.

Proof. From (6), we have that $\theta^{ub} = l'\beta_{A,post} - l'\delta_{post}^{**}$, for δ^{**} a solution to

$$\min_{\delta} l'\delta_{post} \text{ s.t. } A\delta \leq d, \delta_{pre} = \delta_{A,pre}. \quad (37)$$

Let $B = B(\delta^{**})$ index the binding inequalities of the optimization above at δ^{**} , so that

$$A_{(B,\cdot)}\delta^{**} - d_B = 0 \quad (38)$$

$$A_{(-B,\cdot)}\delta^{**} - d_{-B} = -\epsilon < 0. \quad (39)$$

By Assumption 5, $A_{(B,post)}$ has rank $|B|$.

Now, let $\tau^{**} = (\delta_{A,post} + \tau_{A,post}) - \delta_{post}^{**}$. Since by construction $\theta^{ub} \in \mathcal{S}_{\theta}(\Delta, \beta_A)$, we have

$$\begin{pmatrix} \delta_{pre}^{**} \\ \delta_{post}^{**} + \tau^{**} \end{pmatrix} = \begin{pmatrix} \delta_{A,pre} \\ \delta_{A,post} + \tau_{A,post} \end{pmatrix} = \beta_A.$$

It follows that

$$\begin{aligned} A\delta^{**} &= A\beta_A - AM_{post}\tau^{**} \\ &= A\beta_A - AM_{post}\Gamma^{-1}\Gamma\tau^{**} \\ &= A\beta_A - \tilde{A}_{(\cdot,1)}(l'\tau^{**}) - \tilde{A}_{(\cdot,-1)}\Gamma_{(-1,\cdot)}\tau^{**} \\ &= A\beta_A - \tilde{A}_{(\cdot,1)}(\theta^{ub}) - \tilde{A}_{(\cdot,-1)}\tilde{\tau}^{ub}, \end{aligned}$$

where the third equality uses the definition of \tilde{A} and the fact that the first row of Γ is l' ; and the fourth equality uses the fact that $\theta^{ub} = l'((\delta_{A,post} + \tau_{A,post}) - \delta_{post}^{**}) = l'\tau^{**}$

and defines $\tilde{\tau}^{ub} := -\Gamma_{(-1,\cdot)}\tau^{**}$. The first result then follows immediately from the previous display along with (38) and (39).

To show the second set of results, note that $\tilde{A}_{(B,\cdot)} = A_{(B,post)}\Gamma^{-1}$. Since $A_{(B,post)}$ has rank $|B|$ by assumption and Γ^{-1} is full rank, $\tilde{A}_{(B,\cdot)}$ also has rank $|B|$. This implies that $\tilde{A}_{(B,-1)}$ has rank of either $|B| - 1$ or $|B|$. To show that the rank must be $|B| - 1$, note that the optimization (37) can be re-written as

$$\min_{\delta_{post}} l' \delta_{post} \quad \text{s.t.} \quad A_{(\cdot,post)} \delta_{post} \leq d - A_{(\cdot,pre)} \delta_{A,pre}. \quad (40)$$

Since the optimization is assumed to have a finite solution, it is equivalent to its dual formulation,

$$\max_{\gamma} \gamma'(A_{(\cdot,pre)} \delta_{A,pre} - d) \quad \text{s.t.} \quad -\gamma' A_{(\cdot,post)} = l', \gamma \geq 0.$$

Let $\bar{\gamma}$ be a solution to the dual problem. Since $\bar{\gamma}$ is feasible in the dual, $-\bar{\gamma}' A_{(\cdot,post)} = l'$ and $\bar{\gamma} \geq 0$. Additionally, by the complementary slackness conditions, it must be that $\bar{\gamma}_{-B} = 0$. Hence, we have $-\bar{\gamma}'_B A_{(B,post)} = l'$. Multiplying on the right by Γ^{-1} , we obtain $-\bar{\gamma}'_B \tilde{A}_{(B,\cdot)} = l' \Gamma^{-1}$. Recall that by construction the first row of Γ is l' , so $l' = e'_1 \Gamma$, and thus $-\bar{\gamma}'_B \tilde{A}_{(B,\cdot)} = l' \Gamma^{-1} = e'_1$. This shows, however, that $\bar{\gamma}_B$ is in the nullspace of $\tilde{A}'_{(B,-1)}$ but not in the null space of $\tilde{A}'_{(B,\cdot)}$. It follows that the rank of $\tilde{A}_{(B,-1)}$ is strictly less than that of $\tilde{A}_{(B,\cdot)}$, and thus must be equal to $|B| - 1$. Since $\tilde{A}_{(B,-1)}$ has $|B|$ rows and rank $|B| - 1$, by the rank nullity theorem the set $\{\gamma_B : \gamma'_B \tilde{A}_{(B,-1)} = 0\}$ must be one dimensional. We've shown that $\bar{\gamma}'_B \tilde{A}_{(B,-1)} = 0$, and $\bar{\gamma}_B \neq 0$ since $\bar{\gamma}'_B A_{(B,post)} = -l' \neq 0$, which implies that $\{\gamma_B : \gamma'_B \tilde{A}_{(B,-1)} = 0\} = \{c \bar{\gamma}_B : c \in \mathbb{R}\}$, as needed. \square

Lemma B.8. *Let $\Delta = \{\delta : A\delta \leq d\}$, and fix $\delta_A \in \Delta$, τ_A , and Σ^* positive definite. If Assumption 5 holds, then for any $x > 0$,*

$$\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_{\alpha}^C(\hat{\beta}_n; A, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*) \right] \rightarrow 1 - \Phi(z_{1-\alpha} - c^* x),$$

where $\theta_n^{ub} := \sup \mathcal{S}(\Delta_n, \sqrt{n}\delta_A + \sqrt{n}\tau_A)$, $\Delta_n = \{\delta : A\delta \leq \sqrt{nd}\}$, and c^* is a positive constant (not depending on x or α). In particular, $c^* = -\bar{\gamma}'_B \tilde{A}_{(B,1)} / \sigma_B$, where $\sigma_B = \sqrt{\bar{\gamma}'_B A_{(B,\cdot)} \Sigma^* A'_{(B,\cdot)} \bar{\gamma}_B}$ and $\bar{\gamma}_B$ is the unique vector such that $\bar{\gamma}'_B \tilde{A}_{(B,-1)} = 0$, $\bar{\gamma}_B \geq 0$, $\|\bar{\gamma}_B\| = 1$.

Proof. From Lemma B.7, there exists a vector $\tilde{\tau}_1^{ub}$ and a set of indices B such that

$$A_{(B,\cdot)}\beta_A - d_B - \tilde{A}_{(B,1)}\theta_1^{ub} - \tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub} = 0 \quad (41)$$

$$A_{(-B,\cdot)}\beta_A - d_{-B} - \tilde{A}_{(-B,1)}\theta_1^{ub} - \tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub} = -\epsilon < 0, \quad (42)$$

and the set $\{\gamma_B \in \mathbb{R}^{|B|} : \gamma'_B \tilde{A}_{(B,-1)} = 0\} = \{c\bar{\gamma}_B : c \in \mathbb{R}\}$ for some non-zero vector $\bar{\gamma}_B \geq 0$, which without loss of generality we can normalize so that $\|\bar{\gamma}_B\| = 1$. Let $\tilde{\sigma}$ be the vector containing the square roots of the diagonal elements of $A\Sigma^*A'$. It follows that the set $\{\gamma_B \in \mathbb{R}^{|B|} : \gamma'_B \tilde{A}_{(B,-1)} = 0, \gamma'_B \tilde{\sigma}_B = 1, \gamma_B \geq 0\}$ is a singleton. In particular, its lone element is $\gamma_B^* := (\tilde{\sigma}'_B \bar{\gamma}_B)^{-1} \bar{\gamma}_B$. Note that $(\tilde{\sigma}'_B \bar{\gamma}_B)^{-1}$ is well-defined since $\gamma_B \geq 0$ and has at least one strictly positive element, and $\tilde{\sigma} > 0$ since by assumption A has no all-zero rows and Σ^* is positive definite.

Now, consider $\psi_\alpha^C(\hat{\beta}_n, A_{(B,\cdot)}, \sqrt{nd_B}, \theta_n^{ub} + x, \Sigma^*)$, the conditional test that uses only the moments in B . The test statistic for the conditional test that uses only the moments in B is

$$\begin{aligned} \eta(\hat{\beta}_n, A_{(B,\cdot)}, \sqrt{nd_B}, \theta_n^{ub} + x, \Sigma^*) &= \min_{\eta, \tilde{\tau}} \eta \\ \text{s.t. } &A_{(B,\cdot)}\hat{\beta}_n - \sqrt{nd_B} - \tilde{A}_{(B,1)}(\theta_n^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau} \leq \eta\tilde{\sigma}_B. \end{aligned}$$

The equivalent dual problem is

$$\max_{\gamma_B} \gamma'_B \tilde{Y}_{B,n} \text{ s.t. } \gamma'_B \tilde{A}_{(B,-1)} = 0, \gamma'_B \tilde{\sigma}_B = 1, \gamma_B \geq 0,$$

where $\tilde{Y}_{B,n} = A_{(B,\cdot)}\hat{\beta}_n - \sqrt{nd_B} - \tilde{A}_{(B,1)}(\theta_n^{ub} + x)$. We have shown, however, that there is a single value, γ_B^* , that satisfies the constraints of the dual problem, and so the solution to the problem in the previous display is $\gamma_B^* \tilde{Y}_{B,n}$. Additionally, since the set of dual vertices is a singleton, the conditioning event that γ_B^* is optimal is trivial, so $v^{lo} = -\infty$ and $v^{up} = \infty$. It follows that the conditional test using only the moments B is a one-sided t-test that rejects for large values of $\gamma_B^* \tilde{Y}_{B,n}$. Specifically, the critical value is $z_{1-\alpha} \sigma_B^*$, for $\sigma_B^* = \sqrt{\gamma_B^{*'} A_{(B,\cdot)} \Sigma^* A'_{(B,\cdot)} \gamma_B^*}$ the standard deviation of the test statistic $\gamma_B^* \tilde{Y}_{B,n}$. We claim that $\sigma_B^* > 0$. To see why this is the case, observe that Assumption 5 implies that $A_{(B,\cdot)}$ has full row rank, and by construction $\gamma_B^* \neq 0$, so $\gamma_B^{*'} A_{(B,\cdot)} \neq 0$. That $\sigma_B^* > 0$ then follows from the fact that Σ^* is positive definite.

Additionally, observe that

$$\begin{aligned}
\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\gamma_B^{*'} \tilde{Y}_{B,n} \right] &= \gamma_B^{*'} \left[A_{(B,\cdot)} \sqrt{n} \beta_A - \sqrt{n} d_B - \tilde{A}_{(B,1)} (\theta_n^{ub} + x) \right] \\
&= \gamma_B^{*'} \sqrt{n} \left[A_{(B,\cdot)} \beta_A - d_B - \tilde{A}_{(B,1)} \theta_1^{ub} \right] - \gamma_B^{*'} \tilde{A}_{(B,1)} x \\
&= \gamma_B^{*'} [\sqrt{n} \tilde{A}_{(B,-1)} \tilde{\tau}_1^{ub}] - \gamma_B^{*'} \tilde{A}_{(B,1)} x = -\gamma_B^{*'} \tilde{A}_{(B,1)} x
\end{aligned}$$

where the second equality uses $\theta_n^{ub} = \sqrt{n} \theta_1^{ub}$ from Lemma B.1, the third equality uses (41) to substitute for the term in brackets, and the final equality follows from the fact that $\gamma_B^{*'} \tilde{A}_{(B,-1)} = 0$ by construction. Thus, regardless of n , the conditional test using only the moments in B rejects with probability

$$\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^C(\hat{\beta}_n, A_{(B,\cdot)}, \sqrt{n} d_B, \theta_n^{ub} + x, \Sigma^*) \right] = 1 - \Phi(z_{1-\alpha} - (-\gamma_B^{*'} \tilde{A}_{(B,1)} / \sigma_B^*) \cdot x). \quad (43)$$

Note also that we showed in the proof of Lemma B.7 that $-\tilde{\gamma}'_B \tilde{A}_{(B,\cdot)} = e'_1$, which implies that $-\tilde{\gamma}'_B \tilde{A}_{(B,1)} = 1$, and hence $c^* := -\tilde{\gamma}'_B \tilde{A}_{(B,1)} / \sigma_B^* > 0$ since γ_B^* is a positive multiple of $\tilde{\gamma}_B$. Moreover, observe that if we define $\sigma_B = \sqrt{\tilde{\gamma}'_B A_{(B,\cdot)} \Sigma^* A'_{(B,\cdot)} \tilde{\gamma}_B}$, then $\tilde{\gamma}_B^* / \sigma_B^* = \tilde{\gamma}_B / \sigma_B$, so $c^* = -\tilde{\gamma}'_B \tilde{A}_{(B,1)} / \sigma_B$.

Recall that $\psi_\alpha^C(\hat{\beta}_n; A, \sqrt{n} d, \theta_n^{ub} + x, \Sigma^*) = \psi_\alpha^C(\tilde{Y}_n, A \Sigma^* A')$ for $\tilde{Y}_n = A \hat{\beta}_n - \sqrt{n} d - \tilde{A}_{(\cdot,1)} (\theta_n^{ub} + x)$. Since the conditional test optimizes over $\tilde{\tau}$, and $\tilde{\tau}$ appears in this optimization only in the term $\tilde{A}_{(\cdot,-1)} \tilde{\tau}$, the result of the conditional test using \tilde{Y}_n is equivalent to the result of the conditional test that replaces \tilde{Y}_n with $\check{Y}_n = \tilde{Y}_n - \tilde{A}_{(\cdot,-1)} (\sqrt{n} \tilde{\tau}_1^{ub})$ (see Lemma 16 in ARP for a formal justification). That is, $\psi_\alpha^C(\tilde{Y}_n, A \Sigma^* A') = \psi_\alpha^C(\check{Y}_n, A \Sigma^* A')$. The expectation of the elements of \check{Y}_n corresponding with the rows B is

$$\begin{aligned}
\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\check{Y}_{n,B} \right] &= A_{(B,\cdot)} \sqrt{n} \beta_A - \sqrt{n} d_B - \tilde{A}_{(B,1)} (\theta_n^{ub} + x) - \tilde{A}_{(B,-1)} (\sqrt{n} \tilde{\tau}_1^{ub}) \\
&= \sqrt{n} \left(A_{(B,\cdot)} \beta_A - d_B - \tilde{A}_{(B,1)} \theta_1^{ub} - \tilde{A}_{(B,-1)} \tilde{\tau}_1^{ub} \right) - \tilde{A}_{(B,1)} x \\
&= -\tilde{A}_{(B,1)} x,
\end{aligned}$$

where the second line uses the fact that $\theta_n^{ub} = \sqrt{n} \theta_1^{ub}$ from Lemma B.1, and the third uses (41). Similarly, the expectation of the elements of \check{Y}_n corresponding to the rows other than B is

$$\begin{aligned}
\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\check{Y}_{n,-B} \right] &= A_{(-B, \cdot)} \sqrt{n} \beta_A - \sqrt{n} d_{-B} - \tilde{A}_{(-B, 1)} (\theta_n^{ub} + x) - \tilde{A}_{(-B, 1)} (\sqrt{n} \tilde{\tau}_1^{ub}) \\
&= \sqrt{n} \left(A_{(-B, \cdot)} \beta_A - d_{-B} - \tilde{A}_{(-B, 1)} \theta_1^{ub} - \tilde{A}_{(-B, 1)} \tilde{\tau}_1^{ub} \right) - \tilde{A}_{(-B, 1)} x \\
&= \sqrt{n} (-\epsilon) - \tilde{A}_{(-B, 1)} x,
\end{aligned}$$

where the last line uses (42). Since $-\epsilon < 0$, all of the elements of $\mathbb{E} \left[\check{Y}_{n,-B} \right]$ converge to $-\infty$ as $n \rightarrow \infty$, whereas $\mathbb{E} \left[\check{Y}_{n,B} \right]$ does not depend on n . It follows from Proposition 3 in ARP that the conditional test based on the full set of moments is equal to the conditional test that only uses the moments B with probability approaching one,

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left(\psi_\alpha^C(\hat{\beta}_n; A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) = \psi_\alpha^C(\hat{\beta}_n, A_{(B, \cdot)}, \sqrt{n}d_B, \theta_n^{ub} + x, \Sigma^*) \right) = 1.$$

This, combined with (43), gives the desired result. \square

Lemma B.9. *Let \mathcal{B} be a closed, convex subset of \mathbb{R}^K , and $\beta_A \notin \mathcal{B}$. Let $\tilde{\beta} = \arg \min_{\beta \in \mathcal{B}} \|\beta - \beta_A\|_\Sigma$, where $\|x\|_\Sigma^2 = x' \Sigma^{-1} x$ for some positive definite matrix Σ . Then for any $\beta \in \mathcal{B}$, $(\tilde{\beta} - \beta_A)' \Sigma^{-1} (\beta - \tilde{\beta}) \geq 0$.*

Proof. Consider any $\beta \in \mathcal{B}$. Define $\beta_\theta = \theta(\beta - \tilde{\beta}) + \tilde{\beta}$, and note that since \mathcal{B} is convex $\beta_\theta \in \mathcal{B}$ for any $\theta \in [0, 1]$. Further,

$$\|\beta_\theta - \beta_A\|_\Sigma^2 = \theta^2 \|\beta - \tilde{\beta}\|_\Sigma^2 + 2\theta(\tilde{\beta} - \beta_A)' \Sigma^{-1} (\beta - \tilde{\beta}) + \|\tilde{\beta} - \beta_A\|_\Sigma^2.$$

Differentiating with respect to θ , we have

$$\frac{\partial}{\partial \theta} \|\beta_\theta - \beta_A\|_\Sigma^2 = 2\theta \|\beta - \tilde{\beta}\|_\Sigma^2 + 2(\tilde{\beta} - \beta_A)' \Sigma^{-1} (\beta - \tilde{\beta}),$$

from which we see that the derivative evaluated at $\theta = 0$ is $2(\tilde{\beta} - \beta_A)' \Sigma^{-1} (\beta - \tilde{\beta})$. Since $\tilde{\beta}$ minimizes the norm, it follows that we must have $2(\tilde{\beta} - \beta_A)' \Sigma^{-1} (\beta - \tilde{\beta}) \geq 0$, else we could achieve a lower value of the norm at β_θ by choosing θ sufficiently small. \square

Lemma B.10. *Let $\mathcal{B} = \{\beta \in \mathbb{R}^K : v' \beta \leq d\}$ for some $v \in \mathbb{R}^K \setminus \{0\}$ and $d \in \mathbb{R}$. Let $\tilde{\beta} = \arg \min_{\beta \in \mathcal{B}} \|\beta - \beta_A\|_\Sigma$ for some $\beta_A \notin \mathcal{B}$, where $\|x\|_\Sigma^2 = x' \Sigma^{-1} x$ and Σ is positive definite. Then $(\beta_A - \tilde{\beta})' \Sigma^{-1} = c \cdot v'$ for the positive constant $c = \frac{v' \beta_A - d}{\sqrt{v' \Sigma v}}$.*

Proof. Note that we can form a basis $v, \tilde{v}_2, \dots, \tilde{v}_K$ such that $v'\tilde{v}_j = 0$ for $j = 2, \dots, K$. It follows by construction that for any $j = 2, \dots, K$ and any $t \in \mathbb{R}$, $\tilde{\beta} + t \cdot \tilde{v}_j \in \mathcal{B}$. Hence, from Lemma B.9, $-(\beta_A - \tilde{\beta})'\Sigma^{-1}(t\tilde{v}_j) \geq 0$. Since we can choose t both positive and negative, it follows that $(\beta_A - \tilde{\beta})'\Sigma^{-1}\tilde{v}_j = 0$ for all j . Since $(\beta_A - \tilde{\beta})'\Sigma^{-1}$ is orthogonal to $\{\tilde{v}_2, \dots, \tilde{v}_K\}$, and $\{v, \tilde{v}_2, \dots, \tilde{v}_K\}$ form a basis, we have that $(\beta_A - \tilde{\beta})'\Sigma^{-1} = c \cdot v'$, for some $c \in \mathbb{R}$. Multiplying both sides of the equation on the right by Σv , we obtain that $(\beta_A - \tilde{\beta})'v = c \cdot v'\Sigma v$. However, since $\tilde{\beta}$ is the closest point to β_A in Mahalanobis distance, it must be on the boundary of \mathcal{B} , and so $v'\tilde{\beta} = d$. It follows that $c = (v'\beta_A - d)/(v'\Sigma v)$, which is clearly positive since $\beta_A \notin \mathcal{B}$ and thus $v'\beta_A > d$. \square

Lemma B.11. *Let $\mathcal{B} = \{\beta \in \mathbb{R}^K : v'\beta \leq d\}$ for some $v \in \mathbb{R}^K \setminus \{0\}$ and $d \in \mathbb{R}$. Suppose $\hat{\beta} \sim \mathcal{N}(\beta, \Sigma)$ for Σ positive definite known, and consider the problem of testing $H_0 : \beta \in \mathcal{B}$ against $H_A : \beta = \beta_A$ for some $\beta_A \notin \mathcal{B}$. Then the most powerful size- α test of H_0 against H_A is a one-sided t -test that rejects for large values of $v'\hat{\beta}$, and has power equal to $\Phi((v'\beta_A - d)/\sqrt{v'\Sigma v} - z_{1-\alpha})$.*

Proof. From Lemma B.5, the most powerful test rejects for large values of $(\beta_A - \tilde{\beta})'\Sigma^{-1}\hat{\beta}$, where $\tilde{\beta} = \arg \min_{\beta \in \mathcal{B}} \|\beta - \beta_A\|_\Sigma$, and has power $\Phi(\|\beta_A - \tilde{\beta}\|_\Sigma - z_{1-\alpha})$. By Lemma B.10, $(\beta_A - \tilde{\beta})'\Sigma^{-1} = cv'$, for $c = (v'\beta_A - d)/(v'\Sigma v)$. It follows that

$$\begin{aligned} \|\beta_A - \tilde{\beta}\|_\Sigma^2 &= (\beta_A - \tilde{\beta})\Sigma^{-1}(\beta_A - \tilde{\beta}) \\ &= cv'(\beta_A - \tilde{\beta}) \\ &= c(v'\beta_A - d) = (v'\beta_A - d)^2/(v'\Sigma v), \end{aligned}$$

where we use the fact that $v'\tilde{\beta} = d$, since $\tilde{\beta}$ must be on the boundary of \mathcal{B} , as argued in the proof to Lemma B.10. The result then follows immediately. \square

Lemma B.12. *Let $\Delta = \{\delta : A\delta \leq d\}$, and fix $\delta_A \in \Delta$, τ_A , and Σ^* positive definite. If Assumption 5 holds, then for any $x > 0$,*

$$\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left[\psi_\alpha^{MP}(\hat{\beta}_n; A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*) \right] \rightarrow 1 - \Phi(z_{1-\alpha} - c^*x),$$

where ψ_α^{MP} is as defined in the proof to Proposition 4.2, $\theta_n^{ub} := \sup \mathcal{S}(\Delta_n, \sqrt{n}\delta_A + \sqrt{n}\tau_A)$, $\Delta_n = \{\delta : A\delta \leq \sqrt{n}d\}$, and c^* is the same positive constant as in Lemma B.8.

Proof. As argued in the proof to Lemma B.6, the null hypothesis $H_0 : \theta = \bar{\theta}, \delta \in$

$\{A\delta \leq d\}$ is equivalent to the null $H_0 : \beta \in \mathcal{B}_0(\bar{\theta}, d) = \{\beta : \exists \tau_{post} \text{ s.t. } l'_{\tau_{post}} = \bar{\theta}, A\beta - d - AM_{post}\tau_{post} \leq 0\}$, which we showed in Lemma B.6 to be equivalent to $\mathcal{B}_0(\bar{\theta}, d) = \{\beta : \eta(\beta, A, d, \bar{\theta}, \Sigma^*) \leq 0\}$ for the function η as defined in (37). Thus, the null hypothesis for the test associated with $\psi_{\alpha}^{MP}(\hat{\beta}_n; A, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*)$ can be written as $H_0 : \beta_n \in \mathcal{B}_{n,0} := \{\beta : \eta(\beta, A, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*) \leq 0\}$. Under the alternative for this test, $\beta_n = \sqrt{n}\beta_A$, so by Lemma B.5 the most powerful test uses the test statistic $(\sqrt{n}\beta_A - \tilde{\beta}_n)' \Sigma^{*-1} \tilde{\beta}_n$, where $\tilde{\beta}_n = \arg \min_{\beta \in \mathcal{B}_{n,0}} \|\beta - \sqrt{n}\beta_A\|_{\Sigma^*}$.

Now, from Lemma B.7, there exists a vector $\tilde{\tau}_1^{ub}$ and a set of indices B such that

$$A_{(B,\cdot)}\beta_A - d_B - \tilde{A}_{(B,1)}\theta_1^{ub} - \tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub} = 0 \quad (44)$$

$$A_{(-B,\cdot)}\beta_A - d_{-B} - \tilde{A}_{(-B,1)}\theta_1^{ub} - \tilde{A}_{(-B,-1)}\tilde{\tau}_1^{ub} = -\epsilon < 0, \quad (45)$$

where $\{\gamma_B \in \mathbb{R}^{|B|} : \gamma_B' \tilde{A}_{(B,-1)} = 0\} = \{c\bar{\gamma}_B : c \in \mathbb{R}\}$ for some non-zero vector $\bar{\gamma}_B \geq 0$. Define $\mathcal{B}_n^B := \{\beta : \eta(\beta, A_{(B,\cdot)}, \sqrt{nd}_B, \theta_n^{ub} + x, \Sigma^*) \leq 0\}$, the analog to $\mathcal{B}_{n,0}$ that restricts attention only to the set of moments B . By an argument analogous to that in the proof to Lemma B.8 (replacing \hat{Y} with μ), we can show that $\eta(\beta, A_{(B,\cdot)}, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*) = \gamma_B^{*'} \mu_{B,n}(\beta)$, where $\mu_{B,n}(\beta) = A_{(B,\cdot)}\beta - \sqrt{nd}_B - \tilde{A}_{(B,1)}(\theta_n^{ub} + x)$ and $\gamma_B^* = (\bar{\gamma}_B' \tilde{\sigma})^{-1} \bar{\gamma}_B$. Note also that (44) implies that $\tilde{A}_{(B,1)}\theta_1^{ub} = A_{(B,\cdot)}\beta_A - d_B - \tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub}$. Substituting into the expression for $\mu_{B,n}(\beta)$ and using the fact that $\theta_n^{ub} = \sqrt{n}\theta_1^{ub}$ by Lemma B.1, we obtain $\mu_{B,n}(\beta) = A_{(B,\cdot)}(\beta - \sqrt{n}\beta_A) - \tilde{A}_{(B,1)}x + \sqrt{n}\tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub}$. Since $\gamma_B^{*'} \tilde{A}_{(B,-1)} = 0$ by construction, this implies that $\gamma_B^{*'} \mu_{B,n}(\beta) = \gamma_B^{*'} (A_{(B,\cdot)}(\beta - \sqrt{n}\beta_A) - \tilde{A}_{(B,1)}x)$. Hence,

$$\begin{aligned} \mathcal{B}_n^B &= \{\beta : \eta(\beta, A_{(B,\cdot)}, \sqrt{nd}, \theta_n^{ub} + x, \Sigma^*) \leq 0\} \\ &= \{\beta : \gamma_B^{*'} (A_{(B,\cdot)}(\beta - \sqrt{n}\beta_A) - \tilde{A}_{(B,1)}x) \leq 0\} \\ &= \{\beta : \gamma_B^{*'} (A_{(B,\cdot)}((\beta - (\sqrt{n} - 1)\beta_A) - \beta_A) - \tilde{A}_{(B,1)}x) \leq 0\} \\ &= \{\beta : (\beta - (\sqrt{n} - 1)\beta_A) \in \mathcal{B}_1^B\} = (\sqrt{n} - 1)\beta_A + \mathcal{B}_1^B. \end{aligned}$$

Now, define $\beta_n^* = \arg \min_{\beta \in \mathcal{B}_n^B} \|\beta - \sqrt{n}\beta_A\|_{\Sigma^*}$. The results above imply that

$$\begin{aligned}
\beta_n^* &= \arg \min_{\beta \in \mathcal{B}_n^B} \|\beta - \sqrt{n}\beta_A\|_{\Sigma^*} \\
&= \arg \min_{\beta \in (\sqrt{n}-1)\beta_A + \mathcal{B}_1^B} \|\beta - \sqrt{n}\beta_A\|_{\Sigma^*} \\
&= (\sqrt{n}-1)\beta_A + \arg \min_{\beta \in \mathcal{B}_1^B} \underbrace{\|\beta + (\sqrt{n}-1)\beta_A - \sqrt{n}\beta_A\|_{\Sigma^*}}_{=(\beta - \beta_A)} \\
&= (\sqrt{n}-1)\beta_A + \beta_1^*.
\end{aligned}$$

Observe that $\mathcal{B}_n^B \supseteq \mathcal{B}_{n,0}$ since \mathcal{B}_n^B is the set of values β that are consistent with a subset of the moments used in $\mathcal{B}_{n,0}$ (formally, $\eta(\beta, A_{(B,\cdot)}, \sqrt{n}d_B\theta_n^{ub} + x, \Sigma^*) \leq \eta(\beta, A, \sqrt{n}d\theta_n^{ub} + x, \Sigma^*)$ since the RHS minimizes the same objective function subject to additional constraints). Thus, $\beta_n^* = \tilde{\beta}_n$ iff $\beta_n^* \in \mathcal{B}_{n,0}$. From the definition of $\mathcal{B}_{n,0}$, this occurs iff there exists a value $\tilde{\tau}_n$ such that

$$\begin{aligned}
A_{(B,\cdot)}\beta_n^* - \sqrt{n}d_B - \tilde{A}_{(B,1)}(\theta_n^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau}_n &\leq 0 \\
A_{(-B,\cdot)}\beta_n^* - \sqrt{n}d_{-B} - \tilde{A}_{(-B,1)}\sqrt{n}(\theta_n^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau}_n &\leq 0
\end{aligned}$$

Now, since $\beta_1^* \in \mathcal{B}_1^B$, there exists a value $\tilde{\tau}_1^*$ such that

$$A_{(B,\cdot)}\beta_1^* - d_B - \tilde{A}_{(B,1)}(\theta_1^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau}_1^* \leq 0.$$

It follows that

$$\begin{aligned}
&A_{(B,\cdot)}\beta_n^* - \sqrt{n}d_B - \tilde{A}_{(B,1)}(\theta_n^{ub} + x) - \tilde{A}_{(B,-1)}(\tilde{\tau}_1^* + (\sqrt{n}-1)\tilde{\tau}_1^{ub}) \\
&= A_{(B,\cdot)}\beta_1^* - d_B - \tilde{A}_{(B,1)}(\theta_1^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau}_1^* + (\sqrt{n}-1) \left[A_{(B,\cdot)}\beta_A - d_B - \tilde{A}_{(B,1)}\theta_1^{ub} - \tilde{A}_{(B,-1)}\tilde{\tau}_1^{ub} \right] \\
&= A_{(B,\cdot)}\beta_1^* - d_B - \tilde{A}_{(B,1)}(\theta_1^{ub} + x) - \tilde{A}_{(B,-1)}\tilde{\tau}_1^* \leq 0,
\end{aligned}$$

where the first equality uses the fact that $\theta_n^{ub} = \sqrt{n}\theta_1^{ub}$ by Lemma B.1 and $\beta_n^* = \beta_1^* + (\sqrt{n}-1)\beta_A$ as shown above, and the second equality uses (44).

Similarly, we have

$$\begin{aligned}
& A_{(-B,\cdot)}\beta_n^* - \sqrt{n}d_{-B} - \tilde{A}_{(-B,1)}\sqrt{n}(\theta_n^{ub} + x) - \tilde{A}_{(B,-1)}(\tilde{\tau}_1^* + (\sqrt{n} - 1)\tilde{\tau}_1^{ub}) = \\
& = A_{(-B,\cdot)}\beta_1^* - d_{-B} - \tilde{A}_{(-B,1)}(\theta_1^{ub} + x) - \tilde{A}_{(-B,1)}\tilde{\tau}_1^* + \\
& (\sqrt{n} - 1) \left[A_{(-B,\cdot)}\beta_A - d_{-B} - \tilde{A}_{(-B,1)}\theta_1^{ub} - \tilde{A}_{(-B,1)}\tilde{\tau}_1^{ub} \right] \\
& = \left[A_{(-B,\cdot)}\beta_1^* - d_{-B} - \tilde{A}_{(-B,1)}(\theta_1^{ub} + x) - \tilde{A}_{(-B,1)}\tilde{\tau}_1^* \right] - (\sqrt{n} - 1)\epsilon,
\end{aligned}$$

for ϵ a vector with strictly positive elements, where the first equality again uses that $\theta_n^{ub} = \sqrt{n}\theta_1^{ub}$ and $\beta_n^* = \beta_1^* + (\sqrt{n} - 1)\beta_A$, and the second equality uses (45). Since the term in brackets in the final expression in the previous display does not depend on n and all elements of the final term go to $-\infty$, for n sufficiently large the expression in the previous display will be less than or equal to 0. Thus, for n sufficiently large, $\beta_n^* = \tilde{\beta}_n$, and hence the MP test of $H_0 : \beta \in \mathcal{B}_{n,0}$ against $H_A : \beta = \sqrt{n}\beta_A$ is equivalent to the most powerful test of $H_0 : \beta \in \mathcal{B}_n^B$ against $H_A : \beta = \sqrt{n}\beta_A$.

We showed earlier in the proof that $\mathcal{B}_n^B = \{\beta : v'\beta \leq \bar{d}_n\}$, for $v = \gamma_B^* A_{(B,\cdot)}$ and $\bar{d}_n = \gamma_B^* \tilde{A}_{(B,1)}x + v'\sqrt{n}\beta_A$. From Lemma B.11, the MP test of $H_0 : \beta \in \mathcal{B}_n^B$ against $H_A : \beta = \beta_A$ has power equal to $\Phi((v'\sqrt{n}\beta_A - \bar{d}_n)/(v'\Sigma^*v) - z_{1-\alpha})$. Plugging in the definitions of v and \bar{d} and cancelling like terms, we obtain that the power of the test is $\Phi(-\gamma_B^* \tilde{A}_{(B,1)}x/\sigma_B^* - z_{1-\alpha})$, for $\sigma_B^* = \sqrt{\gamma_B^* A_{(B,\cdot)}\Sigma^* A'_{(B,\cdot)}\gamma_B^*}$, which coincides with the expression for the limiting power of the conditional test in Lemma B.8, as needed. \square

Lemma B.13. *Let $\eta(\beta, A, d, \bar{\theta}, \Sigma)$ be as defined in (37). Fix Σ^* positive definite. For any δ_A, τ_A , and d , let $\beta_A(\delta_A, \tau_A) = \delta_A + \tau_A$ and $\theta^{ub}(\delta_A, \tau_A, d) = \sup \mathcal{S}(\Delta, \delta_A + \tau_A)$ for $\Delta = \{\delta : A\delta \leq d\}$. Let $\eta^*(x; \delta_A, \tau_A, d) := \eta(\beta_A(\delta_A, \tau_A), A, d, \theta^{ub}(\delta_A, \tau_A, d) + x, \Sigma^*)$. Then there exists a scalar $c(\Sigma^*, A) > 0$ such that $\eta^*(x; \delta_A, \tau_A, d) \geq c(\Sigma^*, A) \cdot x$ for all δ_A, τ_A , and d .*

Proof. Observe that $\eta(\beta_A, A, d, \bar{\theta}, x)$ is equivalent to the linear program

$$\min_{\eta, \tau} \eta \text{ s.t. } A\beta_A - d - A\tau \leq \eta\tilde{\sigma}, \quad l'\tau = \bar{\theta}.$$

The dual formulation for this problem is

$$\max_{\gamma} \begin{pmatrix} \gamma_A \\ \gamma_{\theta} \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \bar{\theta} \end{pmatrix} \text{ s.t. } \gamma'_A A + \gamma_{\theta} l' = 0, \quad \gamma'_A \tilde{\sigma} = 1, \quad \gamma_A \geq 0,$$

where γ_A is a vector with length equal to the number of rows of A , and γ_θ is a scalar. Note that the feasible set for the dual depends on A and Σ^* but not on d, δ_A , or τ_A . Let V_D denote the set of vertices of the dual, which is finite, and recall that maximizing over the feasible set is equivalent to maximizing over the set of vertices.

Now, we first claim that $\eta(\beta_A, A, d, \theta^{ub}, \Sigma^*) = 0$. Note that since θ^{ub} is in the identified set, it must be that $\eta(\beta_A, A, d, \theta^{ub}, \Sigma^*) \leq 0$. Towards contradiction, suppose that $\eta(\beta_A, A, d, \theta^{ub}, \Sigma^*) = -\epsilon_1 < 0$. Then for all $\gamma = \begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix} \in V_D$,

$$\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} \end{pmatrix} \leq -\epsilon_1.$$

Since V_D is finite, $\bar{\gamma}_\theta := \max_{\gamma \in V_D} \gamma_\theta$ is finite. But then for $\epsilon_2 > 0$,

$$\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} + \epsilon_2 \end{pmatrix} \leq -\epsilon_1 + \bar{\gamma}_\theta \epsilon_2.$$

By choosing ϵ_2 sufficiently small, we can make the upper bound in the previous display less than or equal to 0. However, this implies that $\eta(\beta_A, A, d, \theta^{ub} + \epsilon_2, \Sigma^*) \leq 0$. But this in turn implies $\theta^{ub} + \epsilon_2$ is in the identified set, which contradicts θ^{ub} being maximal. Therefore, $\eta(\beta_A, A, d, \theta^{ub}, \Sigma^*) = 0$.

Additionally, we claim that for $\bar{\theta} = \theta^{ub}$, there must be an optimal dual vertex with $\gamma_\theta > 0$. Towards contradiction, suppose not. Then there exists $\epsilon_3 > 0$ such that for all $\gamma = \begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix} \in V_{D,+} := \{\gamma \in V_D : \gamma_\theta > 0\}$, $\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} \end{pmatrix} < -\epsilon_3$. Letting $\epsilon_4 = \epsilon_3 / \max_{\gamma \in V_{D,+}} \gamma_\theta$, it follows that for all $\gamma \in V_{D,+}$, $\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} + \epsilon_4 \end{pmatrix} < 0$.

Additionally, for $\gamma = \begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix} \in V_D \setminus V_{D,+}$, $\gamma_\theta \leq 0$, and so

$$\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} + \epsilon_4 \end{pmatrix} \leq \begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} \end{pmatrix} \leq 0.$$

Thus, $\begin{pmatrix} \gamma_A \\ \gamma_\theta \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} + \epsilon_4 \end{pmatrix} \leq 0$ for all $\gamma \in V_D$, and so $\eta(\beta_A, A, d, \theta^{ub} + \epsilon_4, \Sigma^*) \leq 0$.

However, this implies that $\theta^{ub} + \epsilon_4$ is in the identified set, which contradicts θ^{ub} being maximal. Thus, there must be at least one $\gamma^* \in V_{D,+}$ such that

$$\begin{pmatrix} \gamma_A^* \\ \gamma_\theta^* \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} \end{pmatrix} = 0.$$

Since γ^* remains feasible in the dual with $\bar{\theta} = \theta^{ub} + x$, it follows that $\eta(\beta_A, A, d, \theta^{ub} + x, \Sigma^*)$ is lower bounded by

$$\begin{pmatrix} \gamma_A^* \\ \gamma_\theta^* \end{pmatrix}' \begin{pmatrix} A\beta_A - d \\ \theta^{ub} + x \end{pmatrix} = \gamma_\theta^* \cdot x.$$

Note that the choice of $\gamma^* \in V_{D,+}$ depended on d, δ_A , and τ_A . However, as noted earlier in the proof, the set $V_{D,+}$ depends on A and Σ^* but does not on d, δ_A, τ_A . Since $V_{D,+}$ is finite and $\gamma_\theta > 0$ for all $\gamma \in V_{D,+}$, there is a value $c > 0$ such that $\gamma_\theta \geq c$ for all $\gamma \in V_{D,+}$. Hence, $\eta^*(x; \delta_A, \tau_A, d) \geq c \cdot x$ for all δ_A, τ_A, d , as needed. \square

Lemma B.14. *Let $\alpha \in (0, 1)$ and $c > z_{1-\alpha}$. Then there exists a unique constant $\zeta(c) > 0$ such that*

$$\frac{\Phi(c) - \Phi(c - \zeta(c))}{1 - \Phi(c - \zeta(c))} = 1 - \alpha.$$

Additionally, for any values $z_{lo} < z_{up}$, with z_{lo} and z_{up} potentially infinite-valued, and $\eta > \max\{c, z_{lo} + \zeta(c)\}$,

$$F_{\xi | \xi \in [z_{lo}, \infty)}(\eta) > 1 - \alpha,$$

where $F_{\xi | \xi \in [z_{lo}, z_{up})}(\cdot)$ is the CDF of $\xi \sim \mathcal{N}(0, 1)$ truncated to $[z_{lo}, z_{up})$.

Proof. First, we show that $F_{\xi | \xi \in [z_{lo}, z_{up})}(t)$ is increasing in t and decreasing in z_{lo} and z_{up} , and these comparative statics are strict for $t \in (z_{lo}, z_{up})$. To see this, note that

$$F_{\xi | \xi \in [z_{lo}, z_{up})}(t) = \begin{cases} 0 & \text{for } t \leq z_{lo} \\ \frac{\Phi(t) - \Phi(z_{lo})}{\Phi(z_{up}) - \Phi(z_{lo})} & \text{for } t \in (z_{lo}, z_{up}) \\ 1 & \text{for } t \geq z_{up} \end{cases}$$

It is immediate that $F_{\xi | \xi \in [z_{lo}, z_{up})}(t)$ is increasing in t and decreasing in z_{up} , and strictly so when $t \in (z_{lo}, z_{up})$. Additionally, we have

$$\frac{\partial}{\partial z_{lo}} \frac{\Phi(t) - \Phi(z_{lo})}{\Phi(z_{up}) - \Phi(z_{lo})} = \frac{-\phi(z_{lo}) (\Phi(z_{up}) - \Phi(t))}{(\Phi(z_{up}) - \Phi(z_{lo}))^2},$$

which is clearly negative for $t \in (z_{lo}, z_{up})$, which gives the desired result for z_{lo} .

Next, consider the function

$$f(\zeta) = \frac{\Phi(c) - \Phi(c - \zeta)}{1 - \Phi(c - \zeta)}.$$

Observe that $f(0) = 0$ and $\lim_{\zeta \rightarrow \infty} f(\zeta) = \Phi(c) > 1 - \alpha$. Additionally, the derivative in the previous paragraph (with $z_{up} = \infty$) implies that $\frac{d}{d\zeta} f(\zeta) > 0$ for $\zeta > 0$. It follows that there is a unique value $\zeta(c) > 0$ such that

$$f(c, \zeta(c)) = \frac{\Phi(c) - \Phi(c - \zeta(c))}{1 - \Phi(c - \zeta(c))} = 1 - \alpha,$$

which gives the first result.

Next, we claim that for $z_{lo} \in (-\infty, \infty)$ and $\zeta > 0$, $F_{\xi | \xi \in [z_{lo}, \infty)}(z_{lo} + \zeta)$ is increasing in z_{lo} . To see why this is the case, note that

$$F_{\xi | \xi \in [z_{lo}, \infty)}(z_{lo} + \zeta) = \frac{\Phi(z_{lo} + \zeta) - \Phi(z_{lo})}{1 - \Phi(z_{lo})}.$$

Differentiating with respect to z_{lo} , we obtain

$$\frac{\phi(z_{lo} + \zeta)(1 - \Phi(z_{lo})) - \phi(z_{lo})(1 - \Phi(z_{lo} + \zeta))}{[1 - \Phi(z_{lo})]^2},$$

which is greater than zero iff

$$\frac{\phi(z_{lo} + \zeta)}{1 - \Phi(z_{lo} + \zeta)} > \frac{\phi(z_{lo})}{1 - \Phi(z_{lo})},$$

which holds since the normal hazard function is strictly increasing.

Now, suppose that $\eta > \max\{c, z_{lo} + \zeta(c)\}$. Then

$$\begin{aligned} F_{\xi | \xi \in [z_{lo}, z_{up})}(\eta) &\geq F_{\xi | \xi \in [z_{lo}, \infty)}(\eta) \\ &> F_{\xi | \xi \in [z_{lo}, \infty)}(\max\{c, z_{lo} + \zeta(c)\}) \\ &= \max\{F_{\xi | \xi \in [z_{lo}, \infty)}(c), F_{\xi | \xi \in [z_{lo}, \infty)}(z_{lo} + \zeta(c))\}, \end{aligned} \quad (46)$$

where the first inequality uses the fact that $F_{\xi | \xi \in [z_{lo}, z_{up})}(t)$ is decreasing in z_{up} and the second inequality uses the fact that $F_{\xi | \xi \in [z_{lo}, z_{up})}(t)$ is strictly increasing in t when $t \in (z_{lo}, \infty)$, and that $\max\{c, z_{lo} + \zeta(c)\} > z_{lo}$ since $\zeta(c) > 0$. The final equality again uses the fact that $F_{\xi | \xi \in [z_{lo}, z_{up})}(t)$ is increasing in t .

However, if $z_{l_0} \leq c - \zeta(c)$, then

$$F_{\xi | \xi \in [z_{l_0}, \infty)}(c) \geq F_{\xi | \xi \in [c - \zeta(c), \infty)}(c) = 1 - \alpha,$$

since we've shown that the expression on the left hand side is decreasing in z_{l_0} . On the other hand, if $z_{l_0} \geq c - \zeta(c)$, then

$$F_{\xi | \xi \in [z_{l_0}, \infty)}(z_{l_0} + \zeta(c)) \geq F_{\xi | \xi \in [c - \zeta(c), \infty)}(c) = 1 - \alpha,$$

since we've shown that $F_{\xi | \xi \in [z_{l_0}, \infty)}(z_{l_0} + \zeta)$ is increasing in z_{l_0} . We have thus shown that the max on the right-hand side of (46) is at least $1 - \alpha$, which gives the desired result. \square

Lemma B.15. *For any $t \in \mathbb{R}$, $\int_{-\infty}^t \Phi(x) dx$ is finite. In particular, $\int_{-\infty}^t \Phi(x) dx = t\Phi(t) + \phi(t)$.*

Proof. We have

$$\begin{aligned} \int_{-\infty}^t \Phi(x) dx &= \int_{-\infty}^{\infty} 1[x \leq t] \int_{-\infty}^{\infty} 1[s \leq x] \phi(s) ds dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} 1[s \leq x \leq t] \phi(s) ds dx \\ &= \int_{-\infty}^{\infty} (t - s) 1[s \leq t] \phi(s) ds \\ &= t \int_{-\infty}^{\infty} 1[s \leq t] \phi(s) ds - \int_{-\infty}^{\infty} s 1[s \leq t] \phi(s) ds \\ &= t\Phi(t) - \Phi(t) \mathbb{E}[\xi | \xi \leq t, \xi \sim \mathcal{N}(0, 1)] = t\Phi(t) - \Phi(t) \frac{-\phi(t)}{\Phi(t)}, \end{aligned}$$

where the last line uses the formula for the mean of a truncated normal distribution. Note that we exchange the order of integration via Fubini's theorem, which is valid since the integrand is weakly positive everywhere and thus equal to its absolute value, and we've shown that the integral after switching the order is finite. \square

Lemma B.16. *Suppose $\tilde{Y}(x) \sim \mathcal{N}(\tilde{\mu}(x), \tilde{\Sigma})$ for some $\tilde{\mu}(x)$ such that $\max_{\gamma \in V(\tilde{\Sigma})} \gamma' \tilde{\mu}(x) \geq x > 0$, where $V(\tilde{\Sigma})$ is the set of vertices of the dual feasible set, $F = \{\gamma : \gamma \geq 0, \gamma' \tilde{A}_{(\cdot, -1)} = 0, \gamma' \tilde{\sigma} = 1\}$, and $\tilde{\sigma}$ contains the square root of the diagonal elements of $\tilde{\Sigma}$. Then there exists a function $\rho(x, \tilde{\Sigma})$, not depending on $\tilde{\mu}(x)$, such that $\mathbb{E}[\psi_{\alpha}^C(\tilde{Y}(x), \tilde{\Sigma})] \geq \rho(x, \tilde{\Sigma})$ and $\lim_{n \rightarrow \infty} \rho(x, \tilde{\Sigma}) = 1$.*

Proof. Recall that ψ_α^C is based on the solution to the dual problem, $\hat{\eta} = \max_{\gamma \in V(\tilde{\Sigma})} \gamma' \tilde{Y}$. Specifically, $\psi_\alpha^C(\tilde{Y}, \tilde{\Sigma}) = 1$ iff

$$F_{\xi | \xi \in [v_{\gamma_*}^{lo}, v_{\gamma_*}^{up}]}(\hat{\eta}; \gamma_*' \tilde{\Sigma} \gamma_*) > 1 - \alpha,$$

where γ_* is an optimal solution to the dual ($\gamma_* \in \hat{V}$), and v^{lo}, v^{up} are functions of γ_* , $\tilde{\Sigma}$, and a sufficient statistic $S_{\gamma_*}(\tilde{Y})$ that by construction is independent of $\gamma_*' \tilde{Y}$. (In this proof only, we make the dependence of v^{lo} and v^{up} on γ_* explicit.) If $\gamma_*' \tilde{\Sigma} \gamma_* \neq 0$, then using the standard formula for the CDF of a truncated normal distribution, we have that the conditional test rejects iff

$$\frac{\Phi(\hat{\eta}/\sigma_{\gamma_*}) - \Phi(z_{\gamma_*}^{lo})}{\Phi(z_{\gamma_*}^{up}) - \Phi(z_{\gamma_*}^{lo})} > 1 - \alpha, \quad (47)$$

where $\sigma_{\gamma_*} = \sqrt{\gamma_*' \tilde{\Sigma} \gamma_*}$ and $z_{\gamma_*}^{lo} = \hat{v}_{\gamma_*}^{lo} / \sigma_{\gamma_*}$, $z_{\gamma_*}^{up} = \hat{v}_{\gamma_*}^{up} / \sigma_{\gamma_*}$. By Lemma B.14, for any $c > z_{1-\alpha}$, (47) holds whenever $\hat{\eta}/\sigma_{\gamma_*} > \max\{c, z_{\gamma_*}^{lo} + \zeta(c)\}$, where $\zeta(c)$ is the unique value that solves

$$\frac{\Phi(c) - \Phi(c - \zeta(c))}{1 - \Phi(c - \zeta(c))} = 1 - \alpha.$$

Thus, when $\sigma_{\gamma_*} \neq 0$, $\psi_\alpha^C = 1$ whenever $\hat{\eta}/\sigma_{\gamma_*} > \max\{c, z_{\gamma_*}^{lo} + \zeta(c)\}$, or equivalently, whenever $\hat{\eta} > \sigma_{\gamma_*} c$ and $\hat{\eta}/\sigma_{\gamma_*} - z_{\gamma_*}^{lo} > \zeta(c)$. Additionally, if $\sigma_{\gamma_*} = 0$, then $\psi_\alpha^C = 1$ whenever $\hat{\eta} > 0$.

Let $\bar{\sigma} = \max_{\gamma \in V(\tilde{\Sigma})} \sigma_\gamma$, which is finite since $V(\tilde{\Sigma})$ is finite. Then the preceding discussion implies that for any $c > \max\{z_{1-\alpha}, 0\}$, $\psi_\alpha^C = 1$ whenever

1) $\hat{\eta} > \bar{\sigma} c$, AND

2) $\exists \gamma_* \in \hat{V}$ such that either i) $\sigma_{\gamma_*} = 0$, OR ii) $\sigma_{\gamma_*} > 0$ and $\gamma_*' \tilde{Y} / \sigma_{\gamma_*} - z_{\gamma_*}^{lo} > \zeta(c)$,

where for the second part of condition 2) we use the fact that $\hat{\eta} = \gamma_*' \tilde{Y}$ when $\gamma_* \in \hat{V}$. Hence, $\psi_\alpha^C = 0$ only if either

A) $\hat{\eta} \leq \bar{\sigma} c$, OR

B) $\exists \gamma_* \in \hat{V}$ such that $\sigma_{\gamma_*} > 0$ and $\gamma_*' \tilde{Y} / \sigma_{\gamma_*} - z_{\gamma_*}^{lo} \leq \zeta(c)$,

Now, by assumption there exists some $\bar{\gamma} \in V(\tilde{\Sigma})$ such that $\bar{\gamma}' \tilde{\mu}(x) \geq x$. Since $\bar{\gamma}$ is feasible in the dual problem for $\hat{\eta}$, we see that $\hat{\eta}$ is lower bounded by $\bar{\gamma}' \tilde{Y}$, which is

distributed $\mathcal{N}(\bar{\gamma}'\tilde{\mu}(x), \sigma_{\bar{\gamma}}^2)$. Thus, the probability that condition A) holds is bounded above by the probability that $\bar{\gamma}'\tilde{Y} \leq \bar{\sigma}c$. If $\sigma_{\bar{\gamma}} = 0$, then the probability condition A) holds is 0 so long as $c < \frac{x}{\bar{\sigma}}$. If $\sigma_{\bar{\gamma}} > 0$, then $\mathbb{P}(\bar{\gamma}'\tilde{Y} \leq \bar{\sigma}c) = \Phi\left(\frac{\bar{\sigma}c - \bar{\gamma}'\tilde{\mu}(x)}{\sigma_{\bar{\gamma}}}\right)$. If $\sigma_{\bar{\gamma}} > 0$, then the set $V^+(\tilde{\Sigma}) := \{\gamma \in V(\tilde{\Sigma}); \sigma_{\gamma} > 0\}$ is non-empty. In this case, let $\underline{\sigma} = \min_{\gamma \in V^+} \sigma_{\gamma}$ and note that $\underline{\sigma} > 0$ since V^+ is finite. Then

$$\Phi\left(\frac{\bar{\sigma}c - \bar{\gamma}'\tilde{\mu}(x)}{\sigma_{\bar{\gamma}}}\right) \leq \Phi\left(\frac{\bar{\sigma}c - x}{\sigma_{\bar{\gamma}}}\right) \leq \Phi\left(\frac{\bar{\sigma}}{\underline{\sigma}}c - \frac{x}{\bar{\sigma}}\right),$$

where we use the fact that $\Phi(\cdot)$ is increasing, $c \geq 0$ and $\bar{\gamma}'\tilde{\mu}(x) \geq x > 0$. Thus, if $c < \frac{x}{\bar{\sigma}}$, we have that condition A) holds with probability bounded above by $\Phi\left(\frac{\bar{\sigma}}{\underline{\sigma}}c - \frac{x}{\bar{\sigma}}\right)$.

Now, the probability that condition B) holds is equal to

$$\begin{aligned} & \mathbb{P}\left(\exists \gamma_* \in \hat{V} \text{ s.t. } \sigma_{\gamma_*} > 0 \text{ and } \gamma_*'\tilde{Y}/\sigma_{\gamma_*} - z_{\gamma_*}^{lo} \leq \zeta(c)\right) = \\ & \mathbb{P}\left(\exists \gamma_* \in \hat{V} \text{ s.t. } \sigma_{\gamma_*} > 0 \text{ and } \left|\gamma_*'\tilde{Y}/\sigma_{\gamma_*} - z_{\gamma_*}^{lo}\right| \leq \zeta(c)\right) \leq \\ & \mathbb{P}\left(\exists \gamma_+ \in V^+ \text{ s.t. } \left|\gamma_+' \tilde{Y}/\sigma_{\gamma_+} - z_{\gamma_+}^{lo}\right| \leq \zeta(c)\right) \leq \\ & \sum_{\gamma_+ \in V^+} \mathbb{P}\left(\left|\gamma_+' \tilde{Y}/\sigma_{\gamma_+} - z_{\gamma_+}^{lo}\right| \leq \zeta(c)\right). \end{aligned}$$

The equality above uses the fact that $\gamma_* \in \hat{V}$ implies that $\gamma_*'\tilde{Y}/\sigma_{\gamma_*} - z_{\gamma_*}^{lo} \geq 0$ since $\hat{\eta} \geq v^{lo}$ by construction; and the remaining inequalities follow from standard properties of probability. Next, observe that for $\gamma_+ \in V^+(\tilde{\Sigma})$, $\gamma_+' \tilde{Y}/\sigma_{\gamma_+}$ is normally distributed with variance 1. Additionally, the random variable $z_{\gamma_+}^{lo}$ is by construction independent of $\gamma_+' \tilde{Y}/\sigma_{\gamma_+}$. However, for any variable ξ that is normally distributed with variance 1 and any variable Z independent of ξ ,

$$\begin{aligned} \mathbb{P}_{(\xi, Z)}(|\xi - Z| \leq \zeta) &= \mathbb{E}_Z \left[\mathbb{P}_{\xi|Z}(\xi \in [z - \zeta, z + \zeta] | Z = z) \right] \\ &\leq \max_{v \in \mathbb{R}} \mathbb{P}_{\xi}(\xi \in [v - \zeta, v + \zeta]) = \Phi(\zeta) - \Phi(-\zeta), \end{aligned}$$

where the first equality follows from iterated expectations, the inequality uses the fact that the distribution of ξ is independent of Z , and the final equality uses the fact that the normal distribution is single-peaked at its mean, so the maximal probability that a normal distribution with variance 1 falls in an interval of length 2ζ is $\Phi(\zeta) - \Phi(-\zeta)$. Additionally, observe that $\Phi(\zeta) - \Phi(-\zeta) = \int_{-\zeta}^{\zeta} \phi(t) dt \leq 2\phi(0)\zeta$. It follows that for

any constant $c > \max\{z_{1-\alpha}, 0\}$, the probability condition B) holds is bounded above by $\kappa\zeta(c)$, where we define the constant $\kappa = 2|V^+|\phi(0)$.

Since $\psi_\alpha^C = 0$ only if either condition A) or condition B) holds, the probability that $\psi_\alpha^C = 0$ is bounded above by $\Phi\left(\frac{\bar{\sigma}}{\sigma}c - \frac{x}{\bar{\sigma}}\right) + \kappa\zeta(c)$,

for any $c \in [\max\{z_{1-\alpha}, 0\}, \frac{x}{\bar{\sigma}}]$. Let $c(x) = c_0 \cdot x$ for $c_0 = \frac{1}{2} \frac{\sigma}{\bar{\sigma}^2}$. Note that $c(x) > \max\{z_{1-\alpha}, 0\}$ for $x > \max\{z_{1-\alpha}/c_0, 0\} =: x_{min}$. Note also that $c_0 = \frac{1}{2} \frac{\sigma}{\bar{\sigma}} < \frac{1}{\bar{\sigma}}$, so $c(x) < \frac{1}{\bar{\sigma}}x$. For $x > x_{min}$, we then have that the probability $\psi_\alpha^C = 0$ is bounded above by $\Phi\left(-\frac{1}{2\bar{\sigma}}x\right) + \kappa\zeta(c_0x)$.

Define $\rho(x, \tilde{\Sigma}) = 1 - \Phi\left(-\frac{1}{2\bar{\sigma}}x\right) - \kappa\zeta(c_0x)$ for $x > x_{min}$ and $\rho(x, \tilde{\Sigma}) = 0$ otherwise.

By construction, $\mathbb{E}\left[\psi_\alpha^C(\tilde{Y}, \tilde{\Sigma})\right] \geq \rho(x, \tilde{\Sigma})$. Note that as $x \rightarrow \infty$, $\Phi\left(-\frac{1}{2\bar{\sigma}}x\right) \rightarrow 0$. To complete the proof that $\rho \rightarrow 1$, we show that $\kappa\zeta(c) \rightarrow 0$ as $c \rightarrow \infty$. To show this, observe that for any $\epsilon > 0$, by L'Hospitale's rule,

$$\begin{aligned} \lim_{c \rightarrow \infty} \frac{\Phi(c) - \Phi(c - \epsilon)}{1 - \Phi(c - \epsilon)} &= \lim_{c \rightarrow \infty} \frac{\phi(c) - \phi(c - \epsilon)}{-\phi(c - \epsilon)} \\ &= 1 - \lim_{c \rightarrow \infty} \frac{\phi(c)}{\phi(c - \epsilon)} \\ &= 1 - \lim_{c \rightarrow \infty} \exp\left(-\frac{1}{2}(2c\epsilon - \epsilon^2)\right) = 1. \end{aligned}$$

Additionally, as shown in the proof to Lemma B.14, $\frac{\Phi(c) - \Phi(c - \zeta)}{1 - \Phi(c - \zeta)}$ is increasing in ζ .

It is then immediate that $\limsup_{c \rightarrow \infty} \zeta(c) < \epsilon$ for all $\epsilon > 0$, and hence $\lim_{c \rightarrow \infty} \zeta(c) = 0$. \square

Lemma B.17. *Suppose $\tilde{Y}(x) \sim \mathcal{N}(\tilde{\mu}, \tilde{\Sigma})$ for some $\tilde{\mu}(x)$ such that $\max_{\gamma \in V(\tilde{\Sigma})} \gamma' \tilde{\mu}(x) \geq x > 0$, where $V(\tilde{\Sigma})$ is the set of vertices of the dual feasible set, $F = \{\gamma : \gamma \geq 0, \gamma' \tilde{A}_{(\cdot, -1)} = 0, \gamma' \tilde{\sigma} = 1\}$, and $\tilde{\sigma}$ contains the square root of the diagonal elements of $\tilde{\Sigma}$. Then there exists a function $\rho(x, \tilde{\Sigma})$, not depending on $\tilde{\mu}(x)$, such that $\mathbb{E}\left[\psi_{\kappa, \alpha}^{C-FLCI}(\tilde{Y}(x), \tilde{\Sigma})\right] \geq \rho(x, \tilde{\Sigma})$ and $\lim_{n \rightarrow \infty} \rho(x, \tilde{\Sigma}) = 1$.*

Proof. The proof is nearly identical to that of Lemma B.16. In particular, by analogous argument we can show that the test $\psi_{\kappa, \alpha}^{C-FLCI} = 0$ only if A) $\hat{\eta} \leq \bar{\sigma}c_{\bar{\alpha}}$, or B) $\exists \gamma_* \in \hat{V}$ such that $\sigma_{\gamma_*} > 0$ and $0 \leq \gamma_*' \tilde{Y} / \sigma_{\gamma_*} - z_{C-FLCI, \gamma_*}^{lo} \leq \zeta(c_{\bar{\alpha}})$, where

$z_{C-FLCI, \gamma_*}^{lo} = v_{C-FLCI, \gamma_*}^{lo} / \sigma_{\gamma_*}$, and $c_{\tilde{\alpha}}$ solves $\frac{\Phi(c_{\tilde{\alpha}}) - \Phi(c_{\tilde{\alpha}} - \zeta(c_{\tilde{\alpha}}))}{1 - \Phi(c_{\tilde{\alpha}} - \zeta(c_{\tilde{\alpha}}))} = 1 - \tilde{\alpha}$. Noting that $z_{C-FLCI, \gamma_*}^{lo}$ is independent of $\gamma'_* \tilde{Y}$, we can then obtain upper bounds on the probability that conditions A) or B) hold by an analogous argument to that in the proof to Lemma B.16. \square

Lemma B.18. *Let $\Delta = \{\delta : A\delta \leq d\}$, and let $\theta^{ub}(\Delta, \beta) := \sup \mathcal{S}(\Delta, \beta)$. Then there exists a function $\rho_{LB}(\cdot, \cdot)$ such that for any $\delta \in \Delta$, τ , Σ^* , and $x > 0$,*

$$\mathbb{E}_{(\delta, \tau, \Sigma^*)} \left[\psi_{\alpha}^C(\hat{\beta}_n; A, d, \theta^{ub}(\Delta, \delta + \tau) + x, \Sigma^*) \right] \geq \rho_{LB}(x, \Sigma^*),$$

and for any Σ^* fixed, $\rho_{LB}(x, \Sigma^*) \rightarrow 1$ as $x \rightarrow \infty$. Analogously, there exists a function $\tilde{\rho}_{LB}(\cdot, \cdot)$ such that for any $\delta \in \Delta$, τ , Σ^* , and $x > 0$,

$$\mathbb{E}_{(\delta, \tau, \Sigma^*)} \left[\psi_{\kappa, \alpha}^{C-FLCI}(\hat{\beta}_n; A, d, \theta^{ub}(\Delta, \delta + \tau) + x, \Sigma^*) \right] \geq \tilde{\rho}_{LB}(x, \Sigma^*),$$

and for any Σ^* fixed, $\tilde{\rho}_{LB}(x, \Sigma^*) \rightarrow 1$ as $x \rightarrow \infty$.

Proof. Lemma B.13 implies that there exists a scalar $c(\Sigma^*, A) > 0$ such that

$$c(\Sigma^*, A) \cdot x \leq \eta^*(x; \delta, \tau, \Sigma^*) := \min_{\eta, \tilde{\tau}} \eta \text{ s.t. } \underbrace{A\beta - d - \tilde{A}_{(\cdot, 1)}(\theta^{ub} + x)}_{:= \tilde{\mu}} - \tilde{A}_{(\cdot, -1)}\tilde{\tau} \leq \eta\tilde{\sigma},$$

where $\beta = \delta + \tau$. Reformulating the minimization above in terms of its dual, we have that $c(\Sigma^*, A) \cdot x \leq \max_{\gamma \in V(\tilde{\Sigma}^*)} \gamma' \tilde{\mu}$, where $V(\tilde{\Sigma}^*)$ is the set of vertices of $F = \{\gamma' \tilde{A}_{(\cdot, -1)} = 0, \gamma' \tilde{\sigma} = 0, \gamma \geq 0\}$. Next, recall that by definition, $\psi_{\alpha}^C(\hat{\beta}; A, d, \bar{\theta}, \Sigma^*) = \psi_{\alpha}^C(\tilde{Y}(\hat{\beta}, A, d, \bar{\theta}), A\Sigma^*A')$, where $\tilde{Y}(\hat{\beta}, A, d, \bar{\theta}) = A\hat{\beta} - d - \tilde{A}_{(\cdot, -1)}\bar{\theta}$. Observe that $\mathbb{E}_{(\delta, \tau, \Sigma^*)} \left[\tilde{Y}(\hat{\beta}, A, d, \bar{\theta}) \right] = \tilde{\mu}$. Lemma B.16 then implies that there exists a function $\rho(\cdot, \cdot)$ such that

$$\mathbb{E}_{(\delta, \tau, \Sigma^*)} \left[\psi_{\alpha}^C(\hat{\beta}_n; A, d, \theta^{ub}(\Delta, \delta + \tau) + x, \Sigma^*) \right] \geq \rho(c(\Sigma^*, A) \cdot x, A\Sigma^*A'),$$

and $\rho(\tilde{x}, A\Sigma^*A') \rightarrow 1$ as $\tilde{x} \rightarrow \infty$. The first desired result then follows by defining $\rho_{LB}(x, \Sigma^*) := \rho(c(\Sigma^*, A) \cdot x, A\Sigma^*A')$. The second desired result follows from an analogous argument, appealing to Lemma B.17 instead of Lemma B.16. \square

Lemma B.19. *For any (a, v) , $\bar{b}(a, v) \geq \frac{1}{2} \sup_{\delta_{pre} \in \Delta_{pre}} LID(\Delta, \delta_{pre})$.*

Proof. Since $\beta = \delta + \tau$, we can write the bias of the affine estimator $a + v'\hat{\beta}$ as $b = a + v'\delta + (v_{post} - l)'\tau_{post}$. Since τ_{post} is unrestricted in the maximization in (11), we see that the worst-case bias will be infinite if $v_{post} \neq l$ and the lemma holds trivially. We can thus restrict attention to affine estimators with $v_{post} = l$, in which case the worst-case bias reduces to

$$\bar{b}(a, v) = \sup_{\delta \in \Delta} |a + v'\delta| = \sup_{\delta \in \Delta} |a + v'_{pre}\delta_{pre} + l'\delta_{post}|. \quad (48)$$

Now, pick any $\delta_{pre}^* \in \Delta_{pre}$. First, suppose that the minimum ($\min_{\delta} l'\delta_{post}$, s.t. $\delta \in \Delta, \delta_{pre} = \delta_{pre}^*$) and the equivalent maximum ($\max_{\delta} l'\delta_{post}$, s.t. $\delta \in \Delta, \delta_{pre} = \delta_{pre}^*$) are finite. Let δ^{min} and δ^{max} be the associated solutions. By construction, $\delta_{pre}^{max} = \delta_{pre}^{min} = \delta_{pre}^*$. For any v_{pre} , we apply the triangle inequality to show that

$$\begin{aligned} |a + v'_{pre}\delta_{pre}^{max} + l'\delta_{post}^{max}| + |a + v'_{pre}\delta_{pre}^{min} + l'\delta_{post}^{min}| &\geq |(a + v'_{pre}\delta_{pre}^{max} + l'\delta_{post}^{max}) - (a + v'_{pre}\delta_{pre}^{min} + l'\delta_{post}^{min})| \\ &= |l'\delta_{post}^{max} - l'\delta_{post}^{min}| = LID(\Delta, \delta_{pre}^*). \end{aligned}$$

Note that for any $x_1, x_2 \geq 0$, $\max\{x_1, x_2\} \geq \frac{1}{2}(x_1 + x_2)$. It then follows from the previous display that

$$\max\{|a + v'_{pre}\delta_{pre}^{max} + l'\delta_{post}^{max}|, |a + v'_{pre}\delta_{pre}^{min} + l'\delta_{post}^{min}|\} \geq \frac{1}{2}LID(\Delta, \delta_{pre}^*).$$

Since δ_{pre}^{max} and δ_{pre}^{min} are feasible in the maximization (48), we see that $\bar{b} \geq \frac{1}{2}LID(\Delta, \delta_{pre}^*)$, as needed. To complete the proof, now suppose without loss of generality that $(\max_{\delta} l'\delta_{post}$, s.t. $\delta \in \Delta, \delta_{pre} = \delta_{pre}^*$) = ∞ . Then, we can replay the argument above replacing δ^{max} with a sequence of values $\{\delta_j\}$ such that $l'\delta_j$ diverges, which gives that \bar{b} is infinite and the result follows. \square

Lemma B.20. *Suppose Δ is convex. Suppose there exists $\delta \in \Delta$ such that $LID(\Delta, \delta_{pre}) = \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre}) < \infty$. Then there exists (a, v) such that $\bar{b}(a, v) = \frac{1}{2} \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre})$. Additionally, for any τ and Σ_n , $\mathbb{E}_{(\delta, \tau, \Sigma_n)} [a + v'\hat{\beta}_n] = \frac{1}{2}(\theta^{ub} + \theta^{lb})$, where θ^{ub} and θ^{lb} are the upper and lower bounds of the identified set $\mathcal{S}(\Delta, \delta + \tau)$.*

Proof. Let $b^{max}(\delta_{pre}^*) := (\max_{\delta} l'\delta_{post}$, s.t. $\delta \in \Delta, \delta_{pre} = \delta_{pre}^*)$, where we define $b^{max} = -\infty$ if $\delta_{pre}^* \notin \Delta_{pre}$. Likewise, define $b^{min}(\delta_{pre}^*) := (\min_{\delta} l'\delta_{post}$, s.t. $\delta \in \Delta, \delta_{pre} = \delta_{pre}^*)$, where we define $b^{min} = \infty$ if $\delta_{pre}^* \notin \Delta_{pre}$. Note that Δ convex implies that b^{max} is concave and b^{min} is convex. Thus, $-LID(\delta_{pre}^*) = b^{min}(\delta_{pre}^*) - b^{max}(\delta_{pre}^*)$ is convex

(where we define $LID(\delta_{pre}^*) = -\infty$ if $\delta_{pre}^* \notin \Delta_{pre}$). The domain of $-LID(\delta_{pre}^*)$ (i.e. the set of values for which it is finite) is Δ_{pre} , since it is infinite for $\delta_{pre}^* \notin \Delta_{pre}$ by construction, and by assumption, $LID(\delta_{pre}^*)$ is finite for all $\delta_{pre}^* \in \Delta_{pre}$. Since Δ is assumed to be convex, it is easy to verify that Δ_{pre} is a non-empty convex set, and thus has non-empty relative interior, so the relative interior of the domain of $-LID$ is non-empty.³⁵ It follows from Theorem 8.2 in [Mau Nam \(2019\)](#) that $\partial(-LID) = \partial(-b^{max}) + \partial(b^{min})$ where for a convex function f , ∂f is the subdifferential $\partial f(\bar{x}) := \{v : f(\bar{x}) + v'(x - \bar{x}) \leq f(x), \forall x\}$ and $\partial(-b^{max}) + \partial(b^{min})$ is the Minkowski sum of the two subdifferentials.

Additionally, if $LID(\delta_{pre}) = \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\tilde{\delta}_{pre})$, then $-LID(\delta_{pre}) = \inf_{\tilde{\delta}_{pre} \in \Delta_{pre}} -LID(\tilde{\delta}_{pre})$. Thus, standard results in convex analysis (see, e.g., Theorem 16.2 in [Mau Nam \(2019\)](#)) give that $0 \in \partial(-LID)(\delta_{pre}) + N(\Delta; \delta_{pre})$, where $N(\Delta; \delta_{pre}) = \{v_{pre} : v'_{pre}(\tilde{\delta}_{pre} - \delta_{pre}) \leq 0, \forall \tilde{\delta}_{pre} \in \Delta_{pre}\}$ is the normal cone to Δ_{pre} at δ_{pre} . Hence, there exist vectors $\bar{v}_{min}, \bar{v}_{max}$ such that for all $\tilde{\delta}_{pre} \in \Delta_{pre}$,

$$b^{min}(\delta_{pre}) + \bar{v}'_{min}(\tilde{\delta}_{pre} - \delta_{pre}) \leq b^{min}(\tilde{\delta}_{pre}) \quad (49)$$

$$-b^{max}(\delta_{pre}) + \bar{v}'_{max}(\tilde{\delta}_{pre} - \delta_{pre}) \leq -b^{max}(\tilde{\delta}_{pre}) \quad (50)$$

$$-(\bar{v}_{min} + \bar{v}_{max})'(\tilde{\delta}_{pre} - \delta_{pre}) \leq 0. \quad (51)$$

The inequalities (50) and (51) together imply that for all $\tilde{\delta}_{pre} \in \Delta_{pre}$,

$$b^{max}(\delta_{pre}) + \bar{v}'_{min}(\tilde{\delta}_{pre} - \delta_{pre}) \geq b^{max}(\tilde{\delta}_{pre}). \quad (52)$$

Now, let v be the vector such that $v_{post} = l$ and $v_{pre} = -\bar{v}_{min}$. Observe that

$$\begin{aligned} \max_{\tilde{\delta} \in \Delta} a + v'_{pre} \tilde{\delta}_{pre} + l' \tilde{\delta}_{post} &= \max_{\tilde{\delta}_{pre} \in \Delta_{pre}} \left(a + v'_{pre} \tilde{\delta}_{pre} + \max_{\tilde{\delta} \in \Delta, \tilde{\delta}_{pre} = \tilde{\delta}_{pre}} l' \tilde{\delta}_{post} \right) \\ &= \max_{\tilde{\delta}_{pre} \in \Delta_{pre}} a + v'_{pre} \tilde{\delta}_{pre} + b^{max}(\tilde{\delta}_{pre}) \\ &\leq a + v'_{pre} \delta_{pre} + b^{max}(\delta_{pre}), \end{aligned} \quad (53)$$

where the first equality nests the maximization, the second equality uses the definition of b^{max} , and the inequality follows from (52). An analogous argument using (49) yields

³⁵The relative interior of a set is the interior of the set relative to its affine hull. See, e.g., [Mau Nam \(2019\)](#), Chapter 5.

that

$$\begin{aligned} \min_{\tilde{\delta} \in \Delta} a + v'_{pre} \tilde{\delta}_{pre} + l' \tilde{\delta}_{post} &= \min_{\tilde{\delta}_{pre} \in \Delta_{pre}} a + v'_{pre} \tilde{\delta}_{pre} + b^{min}(\tilde{\delta}_{pre}) \\ &\geq a + v'_{pre} \delta_{pre} + b^{min}(\delta_{pre}). \end{aligned} \quad (54)$$

Now, it is apparent from the formulation of the identified set in (48) that

$$\bar{b}(a, v) = \max \left\{ \left| \max_{\tilde{\delta} \in \Delta} a + v'_{pre} \tilde{\delta}_{pre} + l' \tilde{\delta}_{post} \right|, \left| \min_{\tilde{\delta} \in \Delta} a + v'_{pre} \tilde{\delta}_{pre} + l' \tilde{\delta}_{post} \right| \right\}, \quad (55)$$

which is bounded above by $\max \{ |a + v'_{pre} \delta_{pre} + b^{max}(\delta_{pre})|, |a + v'_{pre} \delta_{pre} + b^{min}(\delta_{pre})| \}$ from the results above. Setting $a = -v'_{pre} \delta_{pre} - \frac{1}{2}(b^{max}(\delta_{pre}) + b^{min}(\delta_{pre}))$, the upper bound in the previous display reduces to $\frac{1}{2}(b^{max}(\delta_{pre}) - b^{min}(\delta_{pre}))$. Since $LID(\Delta, \delta_{pre}) = b^{max}(\delta_{pre}) - b^{min}(\delta_{pre})$ and $LID(\Delta, \delta_{pre}) = \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre})$ by assumption, it is then immediate that $\bar{b} \leq \frac{1}{2} \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre})$. The inequality in the opposite direction follows from Lemma B.19.

Finally, substituting in the definition of a and v above and simplifying, we see that $\mathbb{E}_{(\delta, \tau)} \left[a + v' \hat{\beta}_n \right] = l' \beta_{post} - \frac{1}{2}(b^{max}(\delta_{pre}) + b^{min}(\delta_{pre}))$, which from (5) and (6) we see is the midpoint of the identified set. \square

Lemma B.21. *Let χ_α be the $1 - \alpha$ quantile of the $|\mathcal{N}(b, \sigma^2)|$ distribution for $b \geq 0$. Then $b + \sigma z_{1-\alpha} \leq \chi_\alpha \leq b + \sigma z_{1-\alpha/2}$.*

Proof. Since $|\xi| \geq \xi$, we have that $q_{1-\alpha}(|\xi| | \xi \sim \mathcal{N}(b, \sigma^2)) \geq q_{1-\alpha}(\xi | \xi \sim \mathcal{N}(b, \sigma^2)) = b + \sigma z_{1-\alpha}$, which yields the first inequality. For the second inequality, observe that

$$\begin{aligned} q_{1-\alpha}(|\xi| | \xi \sim \mathcal{N}(b, \sigma^2)) &= q_{1-\alpha}(|\xi + b| | \xi \sim \mathcal{N}(0, \sigma^2)) \\ &\leq b + q_{1-\alpha}(|\xi| | \xi \sim \mathcal{N}(0, \sigma^2)) = b + \sigma z_{1-\alpha/2} \end{aligned}$$

where the first inequality uses the triangle inequality, and the final equality uses the fact that a mean-zero normal distribution is symmetric about 0. \square

Lemma B.22. *Suppose the conditions of Proposition A.1 holds. Then there is a unique pair (\bar{a}, \bar{v}) such that $\bar{b}(\bar{a}, \bar{v}) = \frac{1}{2} \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre}) =: \bar{b}_{min}$. Additionally, $\sqrt{\bar{v}' A \Sigma^* A' \bar{v}} = 1/c^*$, for c^* defined in Proposition 4.2.*

Proof. Let $b^{min}(\delta_{pre, A}) := (\min_{\delta \in \Delta, \delta_{pre} = \delta_{pre, A}} l' \delta_{post})$, and define b^{max} equivalently. In the proof to Lemma B.7, we showed that b^{min} is equivalent to the problem

(40). We also showed that Assumption 5 implies that there is a solution δ^{**} to the optimization associated with b^{min} such $A_{(B,post)}$ has rank $|B|$, where B indexes the binding moments. The solution δ^{**} is thus non-degenerate. It follows that in a neighborhood of $\delta_{pre,A}$, $b^{min}(\delta_{pre}) = b^{min}(\delta_{A,pre}) + \bar{\gamma}' A_{(\cdot,pre)}(\delta_{pre} - \delta_{A,pre})$, where $\bar{\gamma}$ is a solution to the dual problem (see, e.g., Section 10.4 of Schrijver (1986)). By the complementary slackness conditions, $\bar{\gamma}_{-B} = 0$. Moreover, we showed in the proof to Lemma B.7 that $\bar{\gamma}_B$ is the unique vector that satisfies $\bar{\gamma}'_B \tilde{A}_{(B,-1)} = 0, \bar{\gamma}'_B \tilde{A}_{(B,1)} = 1$. By an analogous argument, we can show that $b^{max}(\delta_{pre})$ is also linear in a neighborhood of $\delta_{A,pre}$. Moreover, the gradient for b^{max} must be the same as for b^{min} , since $LID(\Delta, \delta_{pre}) = b^{max}(\delta_{pre}) - b^{min}(\delta_{pre})$ and LID is maximized at $\delta_{pre,A}$ and thus LID must have 0 gradient at $\delta_{A,pre}$.

Next, combining the expression for \bar{b} in (55) along with the equalities in (53) and (54) in the proof to Lemma B.20, we see that for any (a, v) ,

$$\bar{b}(a, v) = \max \left\{ \left| \max_{\tilde{\delta}_{pre} \in \Delta_{pre}} a + v'_{pre} \tilde{\delta}_{pre} + b^{max}(\tilde{\delta}_{pre}) \right|, \left| \min_{\tilde{\delta}_{pre} \in \Delta_{pre}} a + v'_{pre} \tilde{\delta}_{pre} + b^{min}(\tilde{\delta}_{pre}) \right| \right\}. \quad (56)$$

This implies that if (\bar{a}, \bar{v}) are such that $\bar{b}(\bar{a}, \bar{v}) = \bar{b}_{min}$, then for all $\tilde{\delta}_{pre} \in \Delta_{pre}$,

$$\bar{b}_{min} \geq \max \left\{ \left| \bar{a} + \bar{v}'_{pre} \tilde{\delta}_{pre} + b^{max}(\tilde{\delta}_{pre}) \right|, \left| \bar{a} + \bar{v}'_{pre} \tilde{\delta}_{pre} + b^{min}(\tilde{\delta}_{pre}) \right| \right\}. \quad (57)$$

Now, note that by the triangle inequality, for any scalars x_1, x_2, x_3 with $x_2 \geq x_3$, $\max\{|x_1 + x_2|, |x_1 + x_3|\} \geq \frac{1}{2}|x_2 - x_3|$, with equality if and only if $x_1 + x_3 = -(x_1 + x_2)$. Further, recall that $b^{max}(\delta_{A,pre}) - b^{min}(\delta_{A,pre}) = LID(\Delta, \delta_{A,pre}) = 2\bar{b}_{min}$. It follows from these two facts along with the expression in the previous display that

$$\bar{b}_{min} = \bar{a} + \bar{v}'_{pre} \delta_{A,pre} + b^{max}(\delta_{A,pre}) = -(\bar{a} + \bar{v}'_{pre} \delta_{A,pre} + b^{min}(\delta_{A,pre})). \quad (58)$$

Displays (57) and (58) together imply that $\nabla_{\tilde{\delta}_{pre}=\delta_{A,pre}} (\bar{a} + \bar{v}'_{pre} \tilde{\delta}_{pre} + b^{max}(\tilde{\delta}_{pre})) \leq 0$, where $\nabla_{\tilde{\delta}_{pre}=\delta_{A,pre}} f$ represents the gradient of the function $f(\tilde{\delta}_{pre})$ evaluated at $\tilde{\delta}_{pre} = \delta_{A,pre}$, since otherwise there is a local perturbation to $\delta_{A,pre}$ that violates (57). Using the local linearity derived early, the previous display implies that $\bar{v}'_{pre} + \bar{\gamma}' A_{(\cdot,pre)} \leq 0$. By an analogous argument, (57) and (58) imply that $\nabla_{\tilde{\delta}_{pre}=\delta_{A,pre}} (\bar{a} + \bar{v}'_{pre} \tilde{\delta}_{pre} + b^{min}(\tilde{\delta}_{pre})) \geq 0$, which implies that $\bar{v}'_{pre} + \bar{\gamma}' A_{(\cdot,pre)} \geq 0$. Hence, we see that $\bar{v}'_{pre} = -\bar{\gamma}' A_{(\cdot,pre)}$. We argued in the proof to Lemma B.19 that \bar{v}_{post} must equal l , so we

have shown that there is a unique value of \bar{v} . Further, (58) uniquely pins down \bar{a} in terms of \bar{v} , and so the pair (\bar{a}, \bar{v}) is unique, as claimed.

Finally, recall from the proof to Lemma B.7 that $-\bar{\gamma}'A_{(\cdot, post)} = l'$. Hence $\bar{v}' = (-\bar{\gamma}'A_{(\cdot, pre)}, -\bar{\gamma}'A_{(\cdot, post)}) = -\bar{\gamma}'A$ and thus $\bar{v}'\Sigma^*\bar{v} = \bar{\gamma}'A\Sigma^*A'\bar{\gamma}$. Since $\bar{\gamma}_{-B} = 0$ and $\bar{\gamma}'_B\tilde{A}_{(B,1)} = 1$, we see that $1/\sqrt{\bar{v}'\Sigma^*\bar{v}}$ corresponds with the formula for c^* given in the footnote to Proposition 4.2.

□

Lemma B.23. *Suppose the conditions of Proposition A.1 hold. Then $\frac{\sigma_{v_n, n}}{\sigma_{\bar{v}, n}} \rightarrow 1$, where the optimal FLCI is based on the affine estimator $a_n + v'_n\hat{\beta}_n$ and \bar{v} is the unique value such that $\bar{b}(\bar{a}, \bar{v}) = \bar{b}_{min}$.*

Proof. It suffices to show that $\frac{\sqrt{n}\sigma_{v_n, n}}{\sqrt{n}\sigma_{\bar{v}, n}} \rightarrow 1$. Note that $\sqrt{n}\sigma_{\bar{v}, n} = \sigma_{\bar{v}, 1} = \sqrt{\bar{v}'\Sigma^*\bar{v}}$. By assumption, Σ^* is positive definite, and we showed in the proof to Lemma B.19 that $\bar{v}_{post} = l$, so $\bar{v} \neq 0$. Hence $\sigma_{\bar{v}, 1} > 0$. Next, observe that $\sqrt{n}\sigma_{v_n, n} = \sqrt{v'_n\Sigma^*v_n}$. It thus suffices to show that $v_n \rightarrow \bar{v}$, since then both the numerator and denominator converge to the same non-zero limit. To do this, we will show that every subsequence of v_n has a convergent subsequence. Consider a subsequence v_{n_m} . We argued in the proof to Proposition 3.2 that $\sigma_{v_n, n} \leq \sigma_{\bar{v}, n}$, which implies that $\sqrt{v'_n\Sigma^*v_n} \leq \sqrt{\bar{v}'\Sigma^*\bar{v}}$. Thus, v_n is bounded in the Mahalanobis norm using Σ^* , which implies that v_n is bounded in the standard euclidean norm since Σ^* positive definite. It follows that v_{n_m} has a convergent subsequence, $v_{n_{m,1}} \rightarrow v^*$. We argued in the proof to Proposition 3.2 that $\bar{b}(a_n, v_n) \rightarrow \bar{b}_{min}$. This implies, however, that $a_{n_{m,1}}$ is bounded. To see why this is the case, note that if there is a divergent subsequence $a_{n_{m,2}}$, then for any fixed $\tilde{\delta}_{pre} \in \Delta_{pre}$, $|a_{n_{m,2}} + v'_{n_{m,2}, pre}\tilde{\delta}_{pre} + b^{max}(\tilde{\delta}_{pre})|$ diverges since $v_{n_{m,2}, pre} \rightarrow v^*_{pre}$. Equation (56) then implies that $\bar{b}(a_{n_{m,2}}, v_{n_{m,2}})$ diverges, which is a contradiction. Thus $a_{n_{m,1}}$ is bounded, and so we can extract a further subsequence such that $(a_{n_{m,2}}, v_{n_{m,2}}) \rightarrow (a^*, v^*)$. To complete the proof, we argue that $\bar{b}(a, v)$ is continuous in (a, v) on the set of values (a, v) such that $\bar{b}(a, v)$ is finite. This suffices for the desired result, since the continuity of \bar{b} along with the convergences shown above imply that $\bar{b}(a^*, v^*) = \bar{b}_{min}$, and Lemma B.22 then implies that $v^* = \bar{v}$. To show the continuity of $\bar{b}(a, v)$ on the set where it is finite, note that (48) implies that $\bar{b}(a, v) = \max\{(\max_{\delta} a + v'\delta \text{ s.t. } A\delta \leq d), (\min_{\delta} -a - v'\delta \text{ s.t. } A\delta \leq d)\}$. However, the results in Section 10.4 of Schrijver (1986) imply that these functions are continuous in v for values of v for which they are finite.

□

Lemma B.24. $\lim_{x \rightarrow \infty} (cv_\alpha(x) - (z_{1-\alpha} + x)) = 0$.

Proof. $cv_\alpha(x)$ solves $\Phi(cv_\alpha(x) - x) - \Phi(-cv_\alpha(x) - x) = 1 - \alpha$. By Lemma B.21, $cv_\alpha(x) \geq x + z_{1-\alpha}$, which diverges as $x \rightarrow \infty$. Thus, $\Phi(-cv_\alpha(x) - x)$ converges to 0 and $\Phi(cv_\alpha(x) - x) \rightarrow 1 - \alpha$, together implying $cv_\alpha(x) - x \rightarrow z_{1-\alpha}$. □

Lemma B.25. *Suppose that Assumption 5 holds at $\delta_{A,pre}$. Then $LID(\Delta, \delta_{A,pre}) > 0$.*

Proof. From (5) and (6), we see that that $LID(\Delta, \delta_{A,pre}) = 0$ if and only if $b^{max}(\delta_{A,pre}) = b^{min}(\delta_{A,pre})$, where $b^{min}(\delta_{pre,A}) := (\min_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = \delta_{pre,A})$, and b^{max} is defined analogously. In the proof to Lemma B.7, we showed that b^{min} is equivalent to the problem (40). Assumption 5 implies that there is a solution δ_{post}^{**} such that

$$A_{(B,post)} \delta_{post}^{**} = d_B - A_{(B,pre)} \delta_{A,pre} \text{ and } A_{(-B,post)} \delta_{post}^{**} < d_{-B} - A_{(-B,pre)} \delta_{A,pre},$$

where $A_{B,post}$ has rank $|B|$. Observe that if $b^{min}(\delta_{A,pre}) = b^{max}(\delta_{A,pre})$, then it must be that $l' \delta_{pre}^{**} = l' \tilde{\delta}_{pre}$ for any $\tilde{\delta}_{pre}$ that is feasible in the problem (40). It thus suffices to construct a feasible value $\tilde{\delta}_{pre}$ such that $l' \tilde{\delta}_{pre} \neq l' \delta_{pre}^{**}$. Since $A_{(B,post)}$ has rank $|B|$, its image is $\mathbb{R}^{|B|}$, so there exists $\bar{\delta}_{post}$ such that $A_{(B,post)} \bar{\delta}_{post} = -l$, for l the vector of ones. Thus, for any $\epsilon_1 > 0$, we have that $A_{(B,post)}(\delta_{post}^{**} + \epsilon_1 \bar{\delta}_{post}) < d_B - A_{(B,pre)} \delta_{A,pre}$. However, since the moments $-B$ are slack at $\delta_{A,pre}$, for ϵ_1 sufficiently small, we also have $A_{(-B,post)}(\delta_{post}^{**} + \epsilon_1 \bar{\delta}_{post}) < d_{-B} - A_{(-B,pre)} \delta_{A,pre}$. If $l' \bar{\delta}_{post} \neq 0$, then we are done. If $l' \bar{\delta}_{post} = 0$, then since all of the moments are slack at $\delta_{post}^{**} + \epsilon_1 \bar{\delta}_{post}$, for $\epsilon_2 > 0$ sufficiently small, $\tilde{\delta}_{post} = \delta_{post}^{**} + \epsilon_1 \bar{\delta}_{post} + \epsilon_2 l$ is also feasible, and by construction $l'(\tilde{\delta}_{post} - \delta_{post}^{**}) = \epsilon_2 l' l > 0$. □

Lemma B.26. *Suppose Δ is convex and centrosymmetric, and δ_A is such that $\delta \in \Delta$ implies $\delta - \delta_A \in \Delta$. Then δ_A satisfies Assumption 4.*

Proof. Recall from the proof to Lemma B.20 that for any $\delta_{pre}^* \in \Delta_{pre}$, $LID(\Delta, \delta_{pre}^*) = b^{max}(\delta_{pre}^*) - b^{min}(\delta_{pre}^*)$, where the functions b^{min} and $-b^{max}$ are convex. Observe that

$$\begin{aligned} b^{min}(\delta_{pre}^*) &= \left(\min_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = \delta_{pre}^* \right) \\ &= - \left(\max_{\delta} l'(-\delta_{post}), \text{ s.t. } \delta \in \Delta, \delta_{pre} = \delta_{pre}^* \right) \\ &= - \left(\max_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = -\delta_{pre}^* \right) = -b^{max}(-\delta_{pre}^*), \end{aligned}$$

where the third equality uses the fact that Δ is centrosymmetric. Hence, $-LID(\Delta, \delta_{pre}^*) = -b^{max}(\delta_{pre}^*) - b^{max}(-\delta_{pre}^*)$. It follows from the subdifferential sum and chain rules for convex functions (e.g., Theorems 8.2 and 9.3 in [Mau Nam \(2019\)](#)) that $\partial -LID(\Delta, \delta_{pre}^*) = \partial(-b^{max})(\delta_{pre}^*) + (-\partial(-b^{max})(-\delta_{pre}^*))$, for $+$ the Minkowski sum. It is then immediate that $0 \in \partial(-LID(\Delta, 0))$, and hence $0 \in \arg \min_{\tilde{\delta}_{pre} \in \Delta_{pre}} -LID(\Delta, \tilde{\delta}_{pre})$. This implies that $LID(\Delta, 0) = \sup_{\tilde{\delta}_{pre} \in \Delta_{pre}} LID(\Delta, \tilde{\delta}_{pre})$.

To complete the proof, we show that $LID(\Delta, \delta_{A,pre}) \geq LID(\Delta, 0)$. We first claim that for any $\delta \in \Delta$, $\delta + \delta_A \in \Delta$. Indeed, by centrosymmetry, $-\delta \in \Delta$. By assumption, this implies that $-\delta - \delta_A \in \Delta$. Applying centrosymmetry again, we see that $\delta + \delta_A \in \Delta$, as desired. Next, suppose that δ^{max} is optimal in the maximization $b^{max}(0) = (\max_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = 0)$. Then $\delta^{max} + \delta_A$ is feasible in the optimization $(\max_{\delta} l' \delta_{post}, \text{ s.t. } \delta \in \Delta, \delta_{pre} = \delta_{A,pre})$, and thus $b^{max}(\delta_{A,pre}) \geq b^{max}(0) + l' \delta_{A,post}$. By analogous argument, we can obtain that $b^{min}(\delta_{A,pre}) \leq b^{min}(0) + l' \delta_{A,post}$. It follows that $LID(\Delta, \delta_{A,pre}) = b^{max}(\delta_{A,pre}) - b^{min}(\delta_{A,pre}) \geq b^{max}(0) - b^{min}(0) = LID(\Delta, 0)$, as needed. \square

Lemma B.27. *Fix Σ^* positive definite, $\delta_A \in \Delta$, and τ_A . Suppose Assumption 5 holds at δ_A , and let $B = B(\delta^{**})$. Let \hat{V}_n denote the set of optimal vertices used in $\psi_{\alpha}^C(\hat{\beta}_n; A, \sqrt{n}d, \theta_n^{ub} + x, \Sigma^*)$, where $\theta_n^{ub} = \sup S(\Delta_n, \sqrt{n}(\delta_A + \tau_A))$, $\Delta_n = \sqrt{n}\Delta$. Then*

$$\lim_{n \rightarrow \infty} \mathbb{P}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} \left(\hat{V}_n = \{c\bar{\gamma}\} \right) = 1,$$

where $c > 0$ and $\bar{\gamma}$ is the vector such that $\bar{\gamma}_{-B} = 0$ and $\bar{\gamma}_B$ is the unique vector such that $\bar{\gamma}' \tilde{A}_{(\cdot, -1)} = 0$, $\bar{\gamma} \geq 0$, $\|\bar{\gamma}\| = 1$.

Proof. Observe that $\hat{V}_n = \arg \min_{\gamma \in V(\Sigma^*)} \gamma' \tilde{Y}_n$, where $\tilde{Y}_n = A\hat{\beta}_n - \sqrt{n}d - \tilde{A}_{(\cdot, 1)}(\theta_n^{ub} + x)$. Since all vertices $\gamma \in V(\Sigma^*)$ satisfy $\gamma' \tilde{A}_{(\cdot, -1)} = 0$ by definition, we have that $\hat{V}_n = \arg \min_{\gamma \in V(\Sigma^*)} \gamma' \check{Y}_n$, for $\check{Y}_n = \tilde{Y}_n - \tilde{A}_{(\cdot, -1)}(\sqrt{n}\tilde{\tau}_1^{ub})$ and $\tilde{\tau}_1^{ub}$ the vector constructed in the proof to Lemma B.8. However, we showed in the proof to Lemma B.8 that $\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\check{Y}_{n,B}] = -\tilde{A}_{(B, 1)}x$ and $\mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\check{Y}_{n,-B}] \rightarrow -\infty$ as $n \rightarrow \infty$.

Lemmas C.1 and D.7 in the supplementary material together imply that there is a unique vector $\gamma^* \in V(\Sigma^*)$ such that $\gamma_{-B}^* = 0$, which satisfies $\gamma_B^* = c\bar{\gamma}$ for $c > 0$. By definition, $\gamma \geq 0$ for all $\gamma \in V(\Sigma^*)$, and thus γ_{-B} has at least one strictly positive element for all $\gamma \in V(\Sigma^*) \setminus \{\gamma^*\}$. It follows that

$$\lim_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\gamma^{*'} \check{Y}_n] = -\gamma_B^{*'} \tilde{A}_{(B, 1)}x \text{ and } \lim_{n \rightarrow \infty} \mathbb{E}_{(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)} [\gamma' \check{Y}_n] = -\infty, \forall \gamma \in V(\Sigma^*) \setminus \{\gamma^*\}.$$

Let P_n denote the sequence of data-generating processes characterized by $(\sqrt{n}\delta_A, \sqrt{n}\tau_A, \Sigma^*)$. Note that for all n , $(\gamma^* - \gamma)\check{Y}_n$ is normally distributed with covariance $(\gamma^* - \gamma)'\Sigma^*(\gamma^* - \gamma)$ under P_n . This combined with the results in the previous display imply that $\gamma^{*\prime}\check{Y}_n - \gamma'\check{Y}_n \xrightarrow{P_n} \infty$ for all $\gamma \in V(\Sigma^*) \setminus \{\gamma^*\}$. Since $V(\Sigma^*) \setminus \{\gamma^*\}$ is finite, this implies that $\min_{\gamma \in V(\Sigma^*) \setminus \{\gamma^*\}} (\gamma^{*\prime}\check{Y}_n - \gamma'\check{Y}_n) \xrightarrow{P_n} \infty$, from which we see that $\gamma^{*\prime}\check{Y}_n = \max_{\gamma \in V(\Sigma^*)} \gamma'\check{Y}_n$ with probability approaching 1 under P_n , which gives the desired result. \square

Appendix References

Ingster, Yuri and I. A. Suslina, *Nonparametric Goodness-of-Fit Testing Under Gaussian Models* Lecture Notes in Statistics, New York: Springer-Verlag, 2003.

Kaido, Hiroaki, Francesca Molinari, and Jörg Stoye, “Constraint Qualifications in Partial Identification,” *arXiv:1908.09103 [econ]*, August 2019. arXiv: 1908.09103.

Lee, Jason D., Dennis L. Sun, Yuekai Sun, and Jonathan E. Taylor, “Exact Post-Selection Inference, with Application to the Lasso,” *The Annals of Statistics*, June 2016, *44* (3), 907–927.

Nam, Nguyen Mau, “Convex Analysis: An introduction to convexity and nonsmooth analysis,” <https://maunamn.wordpress.com/> 2019.

Schrijver, Alexander, *Theory of Linear and Integer Programming*, Wiley-Interscience, 1986.

Sierksma, Gerard, *Linear and Integer Programming: Theory and Practice, Second Edition*, CRC Press, November 2001. Google-Books-ID: jP6H4xrTBkUC.