

1 Homework 1: your basic toolkit

Remember:

- Homework is due on the 10th of September before class.
- Please cite any source/collaboration.
- You do not have to hand it your homework in LaTeX till the 5th Monday but you are encouraged to start early!

Exercise 1.1. Prove the fundamental theorem of arithmetic. You might have to resort to other sources, but please express your solution in your own words. The point of this exercise is to acquaint you with proof techniques. Here's the statement

Theorem 1.2. *Let n be an integer > 1 . Then n can be written as*

$$n = p_1^{\alpha_1} \cdot \dots \cdot p_k^{\alpha_k}$$

where p_i 's are distinct primes and $\alpha_i \geq 0$. This expression is unique up to rearrangement of the factors.

Exercise 1.3. The integers form the collection

$$\mathbb{Z} = \{0, 1, -1, 2, -2, \dots\}$$

Here are the rules of arithmetic in \mathbb{Z} :

1. If $a, b \in \mathbb{Z}$ then $a + b \in \mathbb{Z}$.

2. If $a, b \in \mathbb{Z}$ then $a \cdot b \in \mathbb{Z}$.

3. If $a, b, c \in \mathbb{Z}$ then

$$(a + b) + c = a + (b + c).$$

4. If $a, b, c \in \mathbb{Z}$ then

$$(a \cdot b) \cdot c = a \cdot (b \cdot c).$$

5. If $a, b \in \mathbb{Z}$ then

$$a \cdot b = b \cdot a.$$

6. There exists $0 \in \mathbb{Z}$ such that for all $a \in \mathbb{Z}$

$$a + 0 = 0 + a = a.$$

Such an element is called a **additive identity**.

7. There exists $1 \in \mathbb{Z}$ such that for all $a \in \mathbb{Z}$

$$a \cdot 1 = 1 \cdot a = a$$

Such an element is called a **multiplicative identity**.

8. For all $a \in \mathbb{Z}$ there exists $-a$ such that

$$a + (-a) = 0 = (-a) + a$$

Such an element is called an **additive inverse**.

Prove, using the rules of arithmetic in \mathbb{Z} :

1. Additive and multiplicative identities are unique.
2. Additive inverses are unique.
3. If $a, b, c \in \mathbb{Z}$ and $a + b = a + c$ then $b = c$.
4. If $a \in \mathbb{Z}$ then $(-1) \cdot a = -a$.
5. If $a, b \in \mathbb{Z}$ then $(-a) \cdot b = a \cdot (-b) = -(a \cdot b)$.
6. If $a, b \in \mathbb{Z}$ then $(-a) \cdot (-b) = ab$.

Exercise 1.4. A **permutation on n -letters** is a way of rearranging the set $\underline{n} := \{1, \dots, n\}$. So for example we can rearrange $\{1, 2, 3\}$ as $\{1, 3, 2\}$. Prove that the number of permutation on n -letters is exactly $n!$.

Exercise 1.5. Let X be a set with n elements. Prove that X has 2^n subsets.

Exercise 1.6. Suppose that $n \geq 4$. Prove by induction that $n! > 2^n$.

Exercise 1.7. Let X be a set and \sim be an equivalence relation. Define $C(a) := \{b \in X : b \sim a\}$. Prove

1. $a \in C(a)$.
2. If $a \sim b$ then $C(a) = C(b)$.
3. If a is not $\sim b$ then $C(a) \cap C(b) = \emptyset$.
4. $\bigcup_{a \in X} C(a) = X$.

Illustrate, by means of a picture, what an equivalence relation looks like in light of the above result.

Exercise 1.8. Let A, B, C be sets. Prove

1. $A \subseteq A$.
2. $A = B$ if and only if $A \subseteq B$ and $B \subseteq A$.
3. If $A \subseteq B, B \subseteq C$ then $A \subseteq C$.
4. $\emptyset \subseteq A$ always.
5. $A \cup (B \cap C) = (A \cup B) \cap C$
6. $A \cap (B \cup C) = (A \cap B) \cup C$
7. $A \cup \emptyset = A$
8. $A \cup B = \emptyset$ if and only if $A = \emptyset$ and $B = \emptyset$
9. $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
10. $(A \cup B)^c = A^c \cap B^c$
11. $(A \cap B)^c = A^c \cup B^c$.

Exercise 1.9. Prove that if A has m elements, B has n elements, then $A \times B$ has mn elements.

2 Homework 2: sets and fields

Exercise 2.1. Write down explicitly the addition and multiplication table for $\mathbb{Z}/3$.

Exercise 2.2. Prove that multiplication in \mathbb{Z}/n is well-defined.

Exercise 2.3. True or false: on the set with one element, there is a unique field structure. Justify your answer

Exercise 2.4. Let k be an field. Prove:

1. for any nonzero element $x \in k$ there exists a unique y such that $xy = 1$. We denote this by x^{-1}
2. For any nonzero element $x \in k$, $(x^{-1})^{-1} = x$.

Exercise 2.5. Let k be a field. Consider the set $k \times k$ endowed with the “pointwise” multiplication and addition: $(x, y) + (x', y') = (x + x', y + y')$ and $(x, y) \cdot (x', y') = (x \cdot x', y \cdot y')$ and the additive identity being $(0, 0)$ and the multiplicative one being $(1, 1)$. True or false: this endows $k \times k$ with the structure of a field. Justify your answer.

Exercise 2.6. Let k be a field and $x, y, z \in k$. Then $x = y$ if and only if $x + z = y + z$.

Exercise 2.7. Let p be a prime. Prove that the following equation holds in \mathbb{F}_p :

$$(x + y)^p = x^p + y^p.$$

Exercise 2.8. Let $f : X \rightarrow Y, g : Y \rightarrow Z$ be functions. Prove that

1. if f, g are injective then $g \circ f$ is injective.
2. if $g \circ f$ is injective then f is injective.
3. True or false: if $g \circ f$ is injective then g is injective. Prove or give a counterexample.

Exercise 2.9. Let $f : X \rightarrow Y, g : Y \rightarrow Z$ be functions. Prove that

1. if f, g are surjective then $g \circ f$ is surjective.
2. if $g \circ f$ is surjective then g is surjective.
3. True or false: if $g \circ f$ is surjective then f is surjective. Prove or give a counterexample.

3 Homework 3: Vector spaces, subspaces and spans

Exercise 3.1. Let k be a field and S be a set. Then let k^S denote the set of functions

$$f : S \rightarrow k.$$

For $f, g \in k^S$, define addition $f + g$ to be the function that sends $s \in S$ to $f(s) + g(s)$, i.e. $(f + g)(s) = f(s) + g(s)$. For $a \in k$ and $f \in k^S$, define scalar multiplication af to be the function that sends $s \in S$ to $af(s)$, i.e. $(af)(s) = af(s)$.

1. Prove that with this addition and scalar multiplication, k^S is a vector space over k .
2. What is the additive inverse $-f$ of a function $f \in k^S$?

Exercise 3.2. Which of the following is *not* a vector space over \mathbb{R} ? If it is not, explain why.

1. \mathbb{R} itself, with its ordinary addition and scalar multiplication.
2. \mathbb{C} with ordinary addition and scalar multiplication.
3. The set $\{0\}$, with the only possible addition and scalar multiplication.
4. The empty set \emptyset (which has no elements).
5. The set $\mathbb{R} \cup \{\infty, -\infty\}$ with elements given by the real numbers together with two additional elements ∞ and $-\infty$. The addition and scalar multiplication is as usual on the subset of \mathbb{R} of $\mathbb{R} \cup \{\infty, -\infty\}$. We extend addition to all elements of $(\mathbb{R} \cup \{\infty, -\infty\}) \times (\mathbb{R} \cup \{\infty, -\infty\})$ by

$$\begin{aligned} x + \infty &= \infty = \infty + x && \text{for all } x \in \mathbb{R} \\ x + (-\infty) &= -\infty = (-\infty) + x && \text{for all } x \in \mathbb{R} \\ \infty + \infty &= \infty \\ (-\infty) + (-\infty) &= -\infty \\ \infty + (-\infty) &= 0 = (-\infty) + \infty, \end{aligned}$$

and extend scalar multiplication to all elements of $\mathbb{R} \times (\mathbb{R} \cup \{\infty, -\infty\})$ by

$$a\infty = \begin{cases} \infty & \text{if } a > 0 \\ 0 & \text{if } a = 0 \\ -\infty & \text{if } a < 0, \end{cases} \quad a(-\infty) = \begin{cases} -\infty & \text{if } a > 0 \\ 0 & \text{if } a = 0 \\ \infty & \text{if } a < 0. \end{cases}$$

Exercise 3.3. Let V be a k -vector space. Suppose that 0_k is the additive identity in the field and 0_V is the additive identity of the vector space. Then prove:

1. The additive identity 0_V of V is unique.
2. $0_k v = 0_V$ for all $v \in V$
3. $\alpha 0_V = 0_V$ for all $\alpha \in k$
4. $(-1)v = -v$ for all $v \in V$.

Exercise 3.4. Let k be a field and V a vector space. Consider the function

$$\alpha \cdot : V \rightarrow V \quad v \mapsto \alpha \cdot v.$$

Is the function

1. injective?
2. surjective?
3. bijective?

Prove or give a counterexample.

Exercise 3.5. Consider the vector space from Question 3.1, which we denote by k^S . Consider the subset

$$k_{\text{fin}}^S := \{f : S \rightarrow k : \text{for all but finitely many } s \in S, f(s) = 0\}.$$

Prove or disprove: k_{fin}^S is a subspace of k^S .

Exercise 3.6. Let k be a field and consider

$$\text{Poly}_k := \{p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0 : a_i \in k\},$$

the set of polynomials over k . Take for granted that Poly_k is a vector space over k . Let $m \geq 0$. True or false: the subset

$$\text{Poly}_{k, \leq m} := \{p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0 : a_i \in k, n \leq m\}$$

is a subspace of Poly_k .

Exercise 3.7. Let V be a vector space and let U_1, U_2 be subspaces of V . Prove that

1. $U_1 \cap U_2$ is a subspace of V .
2. Define $U_1 + U_2 := \{u_1 + u_2 : u_1 \in U_1, u_2 \in U_2\}$; this is a subspace of V .

Is the union $U_1 \cup U_2$ a subspace of V ?

Exercise 3.8. Let k be a field with finitely many elements. Prove that

1. Prove that a vector space V over k is finite-dimensional if and only if it has finitely many elements.
2. Prove that if V is a finite-dimensional vector space over k , it has $(\#k)^{\dim(V)}$ elements.

Exercise 3.9. Which of the vector spaces (and subspaces) above are finite dimensional? (No need to give detailed proof)

4 Homework 4: linear transformations

Exercise 4.1. Let V, W be vector spaces over a field k and $f : V \rightarrow W$ a linear transformation. Then prove that $f(0) = 0$ and $f(-v) = -f(v)$.

Exercise 4.2. Suppose that $f : V \rightarrow W, g : W \rightarrow U$ be linear maps, then prove that

1. $g \circ f : V \rightarrow U$ is a linear map,
2. If $h : W \rightarrow U$ is another linear map prove that $(h + g) \circ f = h \circ f + g \circ f$,
3. if $i : V \rightarrow W$ is another linear map prove that $g \circ (f + i) = g \circ f + g \circ i$.

Exercise 4.3. Let $f : V \rightarrow W$ be an injective linear transformation. Suppose that $\{v_1, \dots, v_n\}$ is a linearly independent list of vectors, prove that $\{f(v_1), \dots, f(v_n)\}$ is linearly independent.

Exercise 4.4. Let k be a field, consider the function

$$\sigma_n : k^{\times n} \rightarrow k^{\times n} \quad (x_1, \dots, x_n) \mapsto (x_n, x_1, \dots, x_{n-1})$$

Prove that this is a linear transformation and write down the corresponding matrix.

Exercise 4.5. Prove that the only linear transformations from a 1-dimensional vector space to itself is given by scalar multiplication.

Exercise 4.6. Let $D : \text{Poly}_{\leq d}(k) \rightarrow \text{Poly}_{\leq d-1}(k)$ be “differentiation” as a linear map. Prove that D^{d+1} is the zero transformation.

Exercise 4.7. Let $g, f : V \rightarrow V$ be a linear transformation and suppose that $\dim V = n$. Prove that

1. $\dim(\text{Im}(f \circ g)) \leq \min\{\dim(\text{Im}(f)), \dim(\text{Im}(g))\}$.
2. $\dim(\text{Im}(f)) + \dim(\text{Im}(g)) \leq \dim(\text{Im}(f \circ g)) + n$.

References